



RESEARCH ARTICLE

HR-LCMS Based Metabolites Profiling of *Careya arborea* Roxb.

Priyanka Gupta^{1*}, Darshana Patil², Avinash Patil¹

Abstract

Novel chemicals and bioactive compounds found in medicinal plants can be used in the discovery and development of new drugs. *Careya arborea* Roxb. (Family: Lecythidaceae) is known to be one of the important Ayurvedic medicinal plants which has an illustrious history of use in traditional medicine. This study aims to establish the phytochemical profile of Hydroalcoholic stem bark and leaf extracts of *C. arborea* Roxb. using High Resolution Liquid Chromatography-Mass Spectrometry (HR-LCMS). The results showed the detection of 9 major phytochemical compounds in bark and 28 major phytochemical compounds in leaves. This HR-LCMS study manifested the presence of several compounds of pharmaceutical importance in the bark and leaf extracts. Furthermore, this study confirmed that *C. arborea* Roxb. is a rich source of biologically active phytochemical constituents which can be the reason for its bioactive potential.

Keywords: *Careya arborea*, High resolution liquid chromatography-mass spectrometry (HR-LCMS), Leaves, Phytochemicals, Stem bark.

Introduction

Medicinal plants are a vital source for improving health and overcoming the side effects of modern medicine. Several scientific studies have shown that medicinal plants have many benefits and have biochemical and molecular effects (Gratus *et al.* 2009). Plants are acknowledged as a biosynthetic laboratory due to the presence of numerous chemical compounds that includes primary and secondary metabolites (Kokate *et al.* 2008). It is of vital importance to acquire knowledge about these phytochemical components in order to discover new therapeutic agents and to synthesize new complex chemicals (Mojab *et al.*, 2003). The

development of various analytical methods has enabled the isolation, identification and structure elucidation of these phytochemicals (Feng *et al.*, 2019).

Liquid Chromatography-Mass Spectrometry (LC-MS) or High-Resolution Liquid Chromatography-Mass Spectrometry (HR-LCMS) is an analytical technique that combined the outstanding separation efficacy of liquid chromatography with sensitive qualitative analysis and specific detection of mass spectroscopy (Ardrey, 2003). The development of hyphenated analytical approaches such as HR-LCMS, GC-MS, LC-NMR, GC- IR, etc. has advanced the identification of new molecules in plants (Singh *et al.*, 2018).

HR-LCMS is a much more widely applicable method and facilitates the analysis of large, polar, ionic, thermally unstable and non-volatile compounds that traditionally have been difficult to analyse. HR-LCMS system is helpful in the correct determination and structural elucidation of known and unknown components. It also offers excellent sensitivity within minimum acquisition time and attain high-quality data (Primer, 1998).

HR- LCMS has a wide range of applications in the field of biomedical sciences for structural interpretation, pharmacokinetic analysis, genetic analysis, etc. (Parasuraman *et al.*, 2014). In particular, this hyphenated analytical technique is greatly used in the identification and very sensitive quantification of natural products which are

¹Department of Biotechnology, B. K. Birla College of Arts, Science & Commerce (Autonomous), Kalyan (West), Maharashtra, India.

²Department of Botany, Smt. C.H.M. College, Ulhasnagar-03, Maharashtra, India.

***Corresponding Author:** Priyanka Gupta, Department of Biotechnology, B. K. Birla College of Arts, Science & Commerce (Autonomous), Kalyan (West), Maharashtra, India, E-Mail: priyagupta049@gmail.com

How to cite this article: Gupta, P., Patil, D., Patil, A. (2024). HR-LCMS Based Metabolites Profiling of *Careya arborea* Roxb. *J. Indian bot. Soc.*, Doi: 10.61289/jibs2024.08.30.103

Source of support: Nil

Conflict of interest: None.

present at minute concentrations in complex matrices. It also makes easy to find and identify suspected impurities present at trace levels in the complex sample (Sauvage *et al.*, 2006; Pitt, 2009 and Pawar *et al.*, 2019). Also, it allows the detection of molecular weight to the nearest 0.001 a.m.u (atomic mass units) (Wu *et al.*, 2012).

Careya arborea Roxb. belongs to family Lecythidaceae, is a medium sized deciduous tree of about 8–20-meter height with a spreading crown. It is commonly called Padmaka in Ayurveda. It is distributed throughout India in deciduous forests and grasslands (Anonymous, 1993). It is globally found in India, Ceylon, Malay Peninsula, Cambodia and Australia up to an altitude of 1500 meters (Kirtikar and Basu, 1975 and Anonymous, 2006). In the present study, *Careya arborea* Roxb. bark and leaf extracts were subjected to High Resolution - Liquid Chromatography and Mass Spectrometry (HR-LCMS) analysis for the separation, identification and characterization of the phytoconstituents based on their retention time.

Materials And Methods

Plant Materials- Collection and Authentication

Stem bark and leaves were collected from the forest area of Badlapur, Mumbai (Maharashtra) and authenticated from Agharkar Research Institute, Pune, India. Leaves were washed with running tap water to remove soil particles and air dried. Bark was also air dried. After drying, both the plant parts were ground into fine powder and stored in an airtight container at room temperature for further studies.

Preparation of Plant extracts

Extracts of bark and leaf were prepared by refluxing 10 grams of air-dried powdered materials in 100 ml of Hydro alcoholic (50 % Ethanol) for 6 hours at 70 °C. The extracts were filtered through Whatmann filter paper no. 1. They were evaporated to dryness using a rotary vacuum evaporator at 70 °C and 629 mm of Hg. Plant extracts were reconstituted in HPLC grade ethanol at a concentration of 100 ppm and submitted for HR-LCMS to Sophisticated Analytical Instrument Facility (SAIF), Indian Institute of Technology (IIT), Powai, Mumbai.

High resolution liquid chromatography and mass spectrometry (HR-LCMS) analysis

HR-LCMS studies and accurate mass measurements were carried out by 6550i funnel Q-TOF LCMS (Agilent Technologies, USA) equipped with a dual AIS ESI ion source. The stationary phase (Column) used was Zorbax SB-C18, 2.1x50mm, 1.8 microns (Agilent Technologies, USA) and the mobile phase used were

- 0.1% Formic acid in Water
- 90% Acetonitrile+ 10% Water+ 0.1% Formic acid

The data acquisition and processing were performed using Mass Hunter software.

The HR-LCMS mass spectrum was interpreted by comparing the spectrum of unknown components with the spectrum of known components. For comparison, we have utilized the SAIF -IIT Bombay database, which contains over 62000 patterns of the spectrum. The components of the trial materials were identified by their names, molecular weights, and structures.

Results And Discussion

The phytochemical profile of *Careya arborea* Roxb. bark and leaf were characterized by using HRLC-MS spectra. In a chromatogram, the relative concentrations of various compounds get eluted as a function of retention time. Based on the height of the peak, we could determine the relative concentration of bioactive compounds present. The mass spectrometer analyses the compounds eluted at different times in order to identify the nature and structure of the compounds. In the data library, these mass spectra can be identified as fingerprints of that compound (Figs 1 and 2).

Phytochemical screening of Hydroalcoholic (50 % Ethanolic) extract of *Careya arborea* Roxb. bark and leaf through High Resolution (HR)-LCMS detected several




Plate 1: HR-LCMS chromatogram of Hydroalcoholic (50 % Ethanolic) extract of *Careya arborea* Roxb. bark




Plate 2: HR-LCMS chromatogram of Hydroalcoholic (50 % Ethanolic) extract of *Careya arborea* Roxb. leaves

phytocompounds but in the present study, compounds having Hits (DB) 5 or less than 5 are selected as the most probable compounds present in bark and leaf.

Major known compounds found to be present in *Careya arborea* Roxb. bark are Mebeverine metabolite (Veratric acid glucuronide), Esculetin, Phe His, Convallatoxin, Dihydrodeoxystreptomycin, 11alpha-acetoxylkivorin, Oleamide, Dodecylbenzene and 2-Oxo-4-methylthiobutanoic acid (Table 1).

Esculetin, Oleamide, Dodecylbenzene, Convallatoxin, Dihydrodeoxy streptomycin and 2-Oxo-4-methyl thio butanoic acid found in bark extract possess various biological activities (Table 2). Oleamide shows hypnotic and antidepressant properties (Basile *et al.*, 1999 and Akanmu *et al.*, 2007) which is commercially used in the form of capsules and powder (Nootropics Depot) for promoting relaxation and supporting healthy sleeping habits. Dodecylbenzene is used as a surfactant cleansing agent in its sulphonic acid form (Becker *et al.*, 2010) and commercially available by the name Sulfonax (Kao chemicals, Europe) for laundry and cleaning purposes.

Careya arborea Roxb. leaf showed the presence of 28 major phytocompounds i.e. Triparanol, Anabasamine, m-Salicylic acid, Benzenemethanol, 2-(2-aminopropoxy)-3-methyl-, Demeclocycline, Tuberonic acid, Vanillylmandelic acid, Betaxolol, Ubiquinone, Convallatoxin, Dihydrodeoxystreptomycin, Nostoxanthin sulfate, Undecanedioic acid, Cucurbitacin P, Madecassic acid, Neamine (Neomycin A), Betamethasone dipropionate, 2-methyl-tridecanedioic acid, Hydrocortisone-17- butyrate, ZK 168281, Atocalcitol, Digitoxigenin, Taxa-4(20),11(12)-dien-5alpha-acetoxy-10beta-ol, (22E, 24E)-1alpha, 25 dihydroxy-26, 27-diethyl-22, 23, 24, 24a-tetrahydro-24a homovitamin D3/(22E, 24E), 1-Hexadecanoyl-sn-glycerol, 2 alpha-(3- Hydroxypropyl)-1 alpha, 25- dihydroxy-19-norvitamin D3, Spheroidene and 1, 2-di-(13Z- docosenoyl)-sn-glycerol (Table 3).

Betaxolol, Triparanol, Anabasamine, Neamine (Neomycin A), m-Salicylic acid, Demeclocycline, Tuberonic acid, Madecassic acid, Digitoxigenin, Betamethasone dipropionate, Convallatoxin and Dihydrodeoxy streptomycin found in leaf extract show various biological activities (Table 3).

Table 1: List of compounds identified in the Hydroalcoholic extract of *C. arborea* Roxb. bark

| No. | Compound Name | RT | Mass | Formula | Hits (DB) |
|-----|---|--------|----------|--|-----------|
| 1 | Mebeverine metabolite (Veratric acid glucuronide) | 1.02 | 358.093 | C ₁₅ H ₁₈ O ₁₀ | 3 |
| 2 | Esculetin | 8.175 | 178.0273 | C ₉ H ₆ O ₄ | 3 |
| 3 | Phe His | 10.603 | 302.1393 | C ₁₅ H ₁₈ N ₄ O ₃ | 4 |
| 4 | Convallatoxin | 11.894 | 550.2712 | C ₂₉ H ₄₂ O ₁₀ | 1 |
| 5 | Dihydrodeoxy streptomycin | 11.895 | 567.2961 | C ₂₁ H ₄₁ N ₇ O ₁₁ | 1 |
| 6 | 11alpha-Acetoxykivorin | 12.238 | 644.2737 | C ₃₄ H ₄₄ O ₁₂ | 1 |
| 7 | Oleamide | 17.856 | 281.2743 | C ₁₈ H ₃₅ NO | 2 |
| 8 | Dodecylbenzene | 17.86 | 246.2359 | C ₁₈ H ₃₀ | 1 |
| 9 | 2-Oxo-4-methylthiobutanoic acid | 18.888 | 148.0168 | C ₅ H ₈ O ₃ S | 1 |

Table 2: Biological activities of some important bioactive compounds found in 50 % Hydroalcoholic extract of *C. arborea* Roxb. bark and leaf

| No. | Compound name | Plant part | Biological activities | References |
|-----|-----------------------------------|------------|--|---|
| 1 | Esculetin | Bark | Antioxidant, Liver protecting, Antidiabetic, Antibacterial and Antitumor activity | Kaneko <i>et al.</i> , 2003; Gilani <i>et al.</i> , 1998; Prabakaran and Natarajan, 2012; Lee <i>et al.</i> , 2014 and Jay <i>et al.</i> , 2013 |
| 2 | Oleamide | Bark | Hypnotic properties, Antidepressant property and vasodilation activity | Basile <i>et al.</i> , 1999; Akanmu <i>et al.</i> , 2007 and Hiley and Hoi, 2007 |
| 3 | Dodecylbenzene | Bark | In cosmetics as surfactant cleansing agents | Becker <i>et al.</i> , 2010 |
| 4 | 2-Oxo-4-methyl thio butanoic acid | Bark | poultry nutritional supplement, to treat renal failure disease and anti-microbial properties | Masud <i>et al.</i> , 1994; Tang <i>et al.</i> , 2011 and Dibner and Buttin 2002 |
| 5 | Betaxolol | Leaf | Treatment of glaucoma | Yu <i>et al.</i> , 1999 |
| 6 | Triparanol | Leaf | Lowering of blood cholesterol level | Steinberg <i>et al.</i> , 1961 |
| 7 | Anabasamine | Leaf | Anti-inflammatory activity | Panthong <i>et al.</i> , 1984 |
| 8 | Neamine (Neomycin A) | Leaf | Antibiotic | Waksman and Lechevalier, 1949 |

| | | | | |
|----|----------------------------|---------------|---|--|
| 9 | m- Salicylic acid | Leaf | Peeling agent in skin disorders, Anti-inflammatory activity and Antioxidant activity | Kligman and Kligman, 1998; Lee and Kim, 2003, Randjelović <i>et al.</i> , 2015 and Tasleem, 2015 |
| 10 | Demeclocycline | Leaf | Antibacterial activity and in the treatment of syndrome of inappropriate antidiuretic hormone (SIADH) | De Troyer and Demanet, 1975, Cherrill <i>et al.</i> , 1975 and Pasquale and Tan, 2005 |
| 11 | Tuberonic acid | Leaf | Tuber inducing properties | Shaikh <i>et al.</i> , 2016 |
| 12 | Betamethasone dipropionate | Leaf | Anti-inflammatory, immunosuppressive and antiproliferative activity Treatment of Psoriasis | Charney, 1976 and Alam <i>et al.</i> , 2012 |
| 13 | Madecassic acid | Leaf | Antidiabetic activity, Anti-colitis effect and Anti-inflammatory activity | Hsu <i>et al.</i> , 2015; Xu <i>et al.</i> , 2017b and Won <i>et al.</i> , 2010 |
| 14 | Digitoxigenin | Leaf | Cardiotonic activity and Cytotoxic activity | Cornelius <i>et al.</i> , 2013 and Schneider <i>et al.</i> , 2018a |
| 15 | Convallatoxin | Bark and Leaf | Antitumor effect and inhibit viral infection and replication | Kaushik <i>et al.</i> , 2017; Schneider <i>et al.</i> , 2018b; Schneider <i>et al.</i> , 2017; Anderson and Barton, 2017; Amarelle and Lecuona, 2018 |
| 16 | Dihydrodeoxy streptomycin | Bark and Leaf | Antimicrobial activity | Patil and Lade, 2018 |

Table 3: List of compounds identified in the Hydroalcoholic extract of *C. arborea* Roxb. leaves

| No. | Compound Name | RT | Mass | Formula | Hits (DB) |
|-----|---|--------|----------|--|-----------|
| 1 | Triparanol | 1.026 | 143.0952 | C ₇ H ₁₃ N O ₂ | 2 |
| 2 | Anabasamine | 1.77 | 253.1569 | C ₁₆ H ₁₉ N ₃ | 1 |
| 3 | m-Salicylic acid | 4.327 | 138.0328 | C ₇ H ₆ O ₃ | 4 |
| 4 | Benzenemethanol, 2- (2-aminopropoxy)-3-methyl- | 9.016 | 196.1113 | C ₁₁ H ₁₆ O ₃ | 2 |
| 5 | Demeclocycline | 9.72 | 464.1003 | C ₂₁ H ₂₁ Cl N ₂ O ₈ | 1 |
| 6 | Tuberonic acid | 11.318 | 226.1227 | C ₁₂ H ₁₈ O ₄ | 4 |
| 7 | Vanillylmandelic acid | 11.474 | 198.0494 | C ₉ H ₁₀ O ₅ | 4 |
| 8 | Betaxolol | 11.508 | 307.2172 | C ₁₈ H ₂₉ N O ₃ | 3 |
| 9 | Ubiquinone | 11.765 | 250.1201 | C ₁₄ H ₁₈ O ₄ | 3 |
| 10 | Convallatoxin | 11.845 | 550.2708 | C ₂₉ H ₄₂ O ₁₀ | 1 |
| 11 | Dihydrodeoxy streptomycin | 11.846 | 567.2964 | C ₂₁ H ₄₁ N ₇ O ₁₁ | 1 |
| 12 | Nostoxanthin sulfate | 12.66 | 702.3691 | C ₄₀ H ₅₅ Na O ₇ S | 1 |
| 13 | Undecanedioic acid | 13.334 | 216.1378 | C ₁₁ H ₂₀ O ₄ | 1 |
| 14 | Cucurbitacin P | 13.449 | 520.3473 | C ₃₀ H ₄₈ O ₇ | 2 |
| 15 | Madecassic acid | 13.787 | 504.3525 | C ₃₀ H ₄₈ O ₆ | 1 |
| 16 | Neamine (Neomycin A) | 14.692 | 322.1817 | C ₁₂ H ₂₆ N ₄ O ₆ | 3 |
| 17 | Betamethasone dipropionate | 15.566 | 504.26 | C ₂₈ H ₃₇ F O ₇ | 1 |
| 18 | 2-Methyl- tridecanedioic acid | 15.933 | 258.1865 | C ₁₄ H ₂₆ O ₄ | 4 |
| 19 | Hydrocortisone-17-butyrate | 16.658 | 432.2537 | C ₂₅ H ₃₆ O ₆ | 2 |
| 20 | ZK 168281 | 16.871 | 510.3392 | C ₃₂ H ₄₆ O ₅ | 3 |
| 21 | Atocalcitol | 17.413 | 494.3448 | C ₃₂ H ₄₆ O ₄ | 3 |
| 22 | Digitoxigenin | 17.656 | 374.2474 | C ₂₃ H ₃₄ O ₄ | 2 |
| 23 | Taxa-4(20),11(12)- dien-5alpha-acetoxy-10beta- ol | 17.953 | 346.2522 | C ₂₂ H ₃₄ O ₃ | 5 |

| | | | | | |
|----|---|--------|----------|-------------------|---|
| 24 | (22E,24E)-1alpha,25 dihydroxy-26,27-diethyl 22,23,24,24a-tetrahydro- 24a-homovitamin D3 / (22E,24E) | 18.201 | 482.3761 | $C_{32}H_{50}O_3$ | 2 |
| 25 | 1-Hexadecanoyl-sn-glycerol | 18.297 | 330.2826 | $C_{19}H_{38}O_4$ | 2 |
| 26 | 2Alpha-(3- Hydroxypropyl)1alpha,25- dihydroxy- 19-norvitamin D3 | 19.189 | 462.3774 | $C_{29}H_{50}O_4$ | 1 |
| 27 | Spheroidene | 20.66 | 568.456 | $C_{41}H_{60}O$ | 2 |
| 28 | 1,2-di-(13Z- docosenoyl)-sn-glycerol | 26.292 | 732.653 | $C_{47}H_{88}O_5$ | 2 |

Salicylic acid is used in many face washes and scrubs (Vichy Normaderm, Alba botanica, Clean n Clear, Paula's choice, etc.) for acne prone skin types. Triparanol was the first synthetic cholesterol lowering drug which was patented in 1959 (Blohm and MacKenzie, 1959). Demeclocycline is used in the treatment of bacterial infections and a condition which causes a very low level of sodium in the blood, called syndrome of inappropriate secretion of antidiuretic hormone (SIADH) (De Troyer and Demanet, 1975, Cherrill *et al.*, 1975 and Pasquale and Tan, 2005). It is available in the market in the form of capsules (Ledemycin, Advanz pharma). Betamethasone dipropionate has immunosuppressive and anti-inflammatory properties (Charney, 1976 and Alam *et al.*, 2012). It is available in the form of spray (Sernivo, DPT Lab), cream (Betamil-M, Merck Ltd) and ointment (Diprolene, MSD Lab) to treat discomfort of various skin conditions such as dermatitis, eczema and psoriasis.

Conclusion

The present study investigated and specified the various active metabolites found in the hydroalcoholic bark and leaf extract of *Careya arborea* Roxb. by carrying out HR-LCMS analysis. Using the results of this study, it should be possible to extract, purify, and screen a variety of secondary active metabolites from this traditionally well-known medicinal plant for pharmacological activity.

References

- Akanmu M, Adeosun SO, Ilesanmi OR (2007). Neuropharmacological effects of oleamide in male and female mice. *Behavioural Brain Research* 182 (1): 88-94.
- Alam MD, Ali MS, Alam N, Alam MI, Anwer T, Imam F, Ali MD, Siddiqui MR, Shamim, M (2012). Design and Characterization of nanostructure topical gel of Betamethasone Dipropionate for Psoriasis. *J App Pharm Sci.* 2 (10): 148-158.
- Amarelle L and Lecuona, E (2018). The antiviral effects of Na, K-ATPase inhibition: A Minireview. *Int. J. Mol. Sci.* (19): 8 2154.
- Anderson SE and Barton CE (2017). The cardiac glycoside convallatoxin inhibits the growth of colorectal cancer cells in a p53-independent manner. *Molecular Genetics and Metabolism Reports* 13: 42-45.
- Anonymous (1993). Indian medicinal plants, A compendium of 500 species. Vol. 1, Orient Longman. 344-346.
- Anonymous (2006). The Ayurvedic Pharmacopoeia of India. Part 1(V), Government of India, Ministry of health and family welfare, Department of Ayush, India Pp-110.
- Ardrey RE (2003). Liquid Chromatography - Mass Spectrometry: An Introduction. Jony Wiley and Sons Ltd., England
- Basile AS, Hanus L and Mendelson WB (1999). Characterization of the hypnotic actions of oleamide. *Neuroreport* 10 947-95.
- Becker LC, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, et al. (2010). Amended safety assessment of Dodecylbenzenesulfonate, Decylbenzenesulfonate, and Tridecylbenzenesulfonate salts as used in cosmetics. *International Journal of Toxicology* 29(4): 288S-305S.
- Blohm TR and MacKenzie RD (1959). Specific inhibition of cholesterol biosynthesis by a synthetic compound (MER-29). *Arch. Biochem. Biophys.* 85: 245-249.
- Charney P (1976). Betamethasone Dipropionate cream for the treatment of Psoriasis: A collaborative evaluation. *Arch Dermatol.* 112(5): 681-683.
- Cherrill DA, Stote RM, Birge JR, Singer I (1975). Demeclocycline treatment in the syndrome of inappropriate antidiuretic hormone secretion. *Ann Intern Med.*, 83: 654-656.
- Cornelius F, Kanai R, Chikashi TA (2013). Structural view on the functional importance of the sugar moiety and steroid hydroxyls of cardiotonic steroids in binding to Na,K-ATPase. *The journal of biological chemistry* 288(9): 6602-6616.
- De Troyer A and Demanet, JC (1975). Correction of antidiuresis by demeclocycline. *N Engl J Med.* 293: 915-918.
- Dibner JJ, Buttin P (2002). Use of organic acids as a model to study the impact of gut microflora on nutrition and metabolism. *J Appl Poultry Res.* 11: 453-463.
- Feng W, Li M, Hao Z, Zhang J (2019). Analytical Methods of Isolation and Identification. In: Rao, V., Mans, D., Rao, L., editors. *Phytochemicals in Human Health* [Internet]. London: IntechOpen doi: 10.5772/intechopen.88122.
- Gilani AH, Janbaz KH, Shah BH (1998). Esculetin prevents liver damage induced by paracetamol and CCl_4 . *Pharmacol. Res.* 37: 31-35.
- Gratus C, Damery S, Wilson S, Warmington S, Routledge P, Grieve R, Steven N, Jones J, Greenfield S (2009). The use of herbal medicines by people with cancer in the UK: a systematic review of the literature. *Q J Med.* 102: 831-42. <https://doi.org/10.1093/qjmed/hcp13>
- Hiley CR, Hoi PM (2007). Oleamide: a fatty acid amide signalling molecule in the cardiovascular system? *Cardiovasc Drug Rev.* 25: 46-60.
- Hsu Y Hung Y Lihong Hu Lee Y and Yi M (2015). Anti-Diabetic Effects of Madecassic Acid and Rotundic Acid. *Nutrients* 7: 10065-10075.
- Jay AR, Rijhwani H, Malapati K, Kumar P, Saikia K, Lakhar M (2013). Anticancer activity of esculetin via-modulation of Bcl-2 and NF- κ B expression in benzo[a]pyrene induced lung carcinogenesis in mice. *Biomed. Prev. Nutr.* 3: 107-112.

- Kaneko T, Tahara S, Takabayasi F (2003). Suppression of lipid hydroperoxide-induced oxidative damage to cellular DNA by esculetin. *Biol. Pharm. Bull.* 26: 840–844.
- Kaushik V, Azad N, Yakisich JS, Iyer AKV (2017). Antitumor effects of naturally occurring cardiac glycosides convallatoxin and peruvoside on human ER+ and triple-negative breast cancers. *Cell Death Discovery* 3: 1-9.
- Kirtikar KR and Basu BD (1975). Indian medicinal plants. 2nd ed. Dehradun, India 894-895.
- Kligman D, Kligman AM (1998). Salicylic acid peels for the treatment of photoaging. *Dermatol Surg.* 24: 325–328.
- Kokate C K, Purohit A P, Gokhale S B (2008). Pharmacognosy, 42th ed. Nirali Publication, Pune
- Lee HS, Kim IH. (2003). Salicylic acid peels for the treatment of *Acne vulgaris* in Asian patients. *Dermatol Surg.* 29: 1196–1199.
- Lee JH, Kim YG, Cho HS, Ryu SY, Cho MH, Lee J (2014). Coumarins reduce biofilm formation and the virulence of *Escherichia coli*. *Phytomedicine* 21: 1037–1042.
- Masud T, Young VR, Chapman T, Maroni BJ (1994) Adaptive responses to very low protein diets-the first comparison of ketoacids to essential amino acids. *Kidney Int.* 45: 1182–1192.
- Mojab F, Kamalinejad M, Naysaneh G, Vahidipour H (2003). Phytochemical Screening of Some Species of Iranian Plants. *Iranian Journal of Pharmaceutical Research* 2: 77-82.
- Panthong A, Kanjanapothi D, Thitiponpant Y, Taesotikul T, Arbain D (1984). Anti-inflammatory activity of the alkaloid anabasamine. *Dokl Akad Nauk Uzssr* 198: 45-47.
- Parasuraman S, Anish R, Subramani B, Selvadurai M, Kalaimani J K, Venugopal V (2014). An overview of Liquid Chromatography-Mass Spectroscopy instrumentation. *Pharmaceutical Methods* 5(2): 47–55.
- Pasquale TR, Tan JS (2005). Nonantimicrobial Effects of antibacterial agents. *Reviews of anti-infective agents* 40: 127-135.
- Patil AS, Lade BD (2018). Wound stress induced secondary metabolites in *Passiflora foetida*: Exploration of antimicrobial compounds. *Saudi Journal of Medical and Pharmaceutical Sciences* 4(5): 613-627.
- Pawar DS, Ghodke J, Nasreen S (2019). Antimicrobial activity and HR-LCMS analysis of methanolic extract of *Calotropis gigantea*. *International journal of advanced science and research* 4(1): 19-24.
- Pitt JJ (2009). Principles and Applications of Liquid Chromatography-Mass Spectrometry in Clinical Biochemistry. *Clin Biochem Rev* 30: 19-34
- Prabakaran D, Natarajan A (2012). Antihyperglycemic effect of esculetin modulated carbohydrate metabolic enzymes activities in streptozotocin induced diabetic rats. *J. Funct. Foods* 4: 776–783.
- Primer A (1998). Basics of LC/MS, Hewlett-Packard Company, USA
- Randjelović P, Veljković S, Stojiljković N, Sokolović D, Ilić I, Laketić D, et al. (2015). The beneficial biological properties of salicylic acid. *Acta Fac. Medicae Naissensis* 32: 259–265.
- Sauvage FL, Marcoux FS, Duret B, Deporte D, Lachatre G, Marquet P (2006). Screening of drugs and toxic compounds with liquid chromatography-linear ion trap tandem mass spectrometry. *Clinical Chemistry* 52(9): 1735–1742
- Schneider NF, Geller FC, Persich L, Marostica LL, Padua RM, Kreis W, et al. (2018b). Inhibition of cell proliferation, invasion and migration by the cardenolides digitoxigenin monodigitoxoside and convallatoxin in human lung cancer cell line. *Natural Product Research* 30(11): 1327–1331.
- Schneider NFZ, LaraPersichaRochab C, PachecoRamosb AC, FariaCortesbV, ThaísSilva I, Rodrigo M, et al. (2018a). Cytotoxic and cytostatic effects of digitoxigenin monodigitoxoside (DGX) in human lung cancer cells and its link to Na,K-ATPase. *Biomedicine and Pharmacotherapy* 97: 684–696.
- Schneider NFZ, Silva IT, Perish L, De A, Rocha SC, Marostica L. et al. (2017). Cytotoxic effects of the cardenolide convallatoxin and its Na, K-ATPase regulation. *Molecular and Cellular Biochemistry* 428(1–2): 23–29.
- Shaikh AC, Gupta A, Chaphalkar SR (2016). Quantitative investigation of phytochemicals in Nakshtra plants and distribution analysis of ketones pertaining to LC-MS. *Int. J. Chem, Pharm, Sci.* 4(7): 381-393.
- Singh PA, Desai SD, Singh J (2018). A Review on plant antimicrobials of past decade. *Current Topics in Medicinal Chemistry* 18(10): 812-833
- Steinberg D, Avigan J, Feigelson EB (1961). Effects of triparanol (mer-29) on cholesterol biosynthesis and on blood sterol levels in man. *The journal of clinical investigations* 40(5): 884-893.
- Tang X, Yang YL, Shi YH, Le GW (2011). Comparative in vivo antioxidant capacity of dl-2-hydroxy-4- methylthiobutanoic acid (HMTBA) and dl-methionine in male mice fed a high-fat diet. *J Sci Food Agr.* 91: 2166–2172.
- Tasleem A (2015). Salicylic acid as a peeling agent: a comprehensive review. *Clinical, cosmetic and investigational dermatology* 8: 455–461.
- Waksman SA, Lechevalier HA (1949). Neomycin, a new antibiotic active against Streptomycin-resistant bacteria, including tuberculosis organisms. *Science* 109: 305–307.
- Won JH, Shin JS, Park HJ, Jung H, Koh D, Jo B, Lee J, Yun K, Lee KT (2010). Anti-inflammatory Effects of Madecassic Acid via the Suppression of NF-κB Pathway in LPS-Induced RAW 264.7 Macrophage Cells. *Planta Med* 76(3): 251-257.
- Wu AH, Gerona R, Armenian P, French D, Petrie M, Lynch KL (2012). Role of liquid chromatography-high-resolution mass spectrometry (LC-HR/MS) in clinical toxicology. *Clin Toxicol (Phila)*. 50(8): 733–42. doi: 10.3109/15563650.2012.713108.
- Xu X, Wang Y, Wei Z, Wei W, Zhao P, Tong B, Xia Y, Dai Y (2017). Madecassic acid, the contributor to the anti-colitis effect of madecassoside, enhances the shift of Th17 toward Treg cells via the PPARγ/AMPK/ACC1 pathway. *Cell Death and Disease* 8:1-15.
- Yu DY, Su EN, Cringle SJ, Alder VA, Yu PK, Desantis L (1999). Systemic and ocular vascular roles of the antiglaucoma agents beta-adrenergic antagonists and Ca²⁺ entry blockers. *Surv. Ophthalmol.* 43(Suppl 1): 214–222.