



RESEARCH ARTICLE

PGPR attributes and molecular identification of non-rhizobial Endophytes for growth enhancement of *Sesbania sesban* (L) Merr.

Prashant Kumar, Harshita Vashistha, Sandeep Kumar*

Abstract

The ongoing research explores the isolation, characterization, identification and plant growth promoting (PGP) functions of selected plant growth-promoting rhizobacteria (PGPR) isolated from the root nodules of a wild leguminous plant *Sesbania sesban* (L) Merr., growing in a plain area adjacent to the Shivalik hill regions of Uttarakhand. From an initial pool of 70 bacterial isolates, 16 gram-negative strains were selected and screened for their potential PGP traits, such as phosphate solubilization, siderophore production, hydrogen cyanide (HCN) and indole-3-acetic acid (IAA) production. Notably, bacterial strains SSP1 and SSP3, demonstrated strong efficacy in these traits. Ribotyping analysis of 16S rRNA genes identified them as *Enterobacter xiangfangensis* SSP01 and *Acinetobacter pittii* SSP03, showcasing their taxonomy as non-rhizobial endophytes. Moreover, these bacterial strains demonstrated significant enhancements in biomass accumulation and vegetative growth parameters in *S. sesban*. In pot trial experiments, the SSP01 and SSP03 consortia displayed better results, highlighting their potential in agricultural applications. By exploiting the beneficial properties of these bacterial strains, our findings offers a promising avenue for sustainable agriculture, presenting an eco-friendly alternative to conventional chemical fertilizers.

Keywords: *Sesbania sesban*, Traditional green manure, PGPR attributes, Molecular identification, Vegetative growth, Pot trial.

Introduction

Excessive utilization of pesticides, fungicides and chemical fertilizers in modern agronomy poses significant risks to animals, humans, the environment, and microbial life forms. The advent of the Green Revolution in 1967 brought about a remarkable transformation in Indian agriculture, resulting in unprecedented levels of crop production. Over the course of the past few years, the rate of growth in crop production has been decreasing continuously. This decline can be attributed to the worsening of soil quality and health, primarily caused by the immoderate

utilization of agrochemicals, pesticides, fungicides, and chemical fertilizers in modern farming systems. Pesticides are regarded as an important tool for crop protection and growth, but they are toxic to the environment as well as for all living organisms. overuse of pesticides induces biodiversity loss and destruction. (Ahmed and Turchi 2021, Khan *et al.* 2023, Shafiq *et al.* 2024). In India, soils typically exhibit poor fertility due to nutrient leaching resulting from intensive cultivation, limited agricultural knowledge, imbalanced use of chemical fertilizers, and inadequate fertilization which have been used more frequently over the past few decades, due to which soil efficiency deteriorates. These factors not only support a decline in crop productivity but also lead to greenhouse gas emissions and water contamination (Alemineu and Alemayehu 2020). The diminishing fertility of the soil also poses a serious challenge to restoring vegetation on degraded lands. One of the key nutrient deficiencies observed in these soils is nitrogen, which performs an imperative role in maintaining soil fecundity. However, nitrogen fertilizer is expensive, scarce, and prone to rapid depletion through leaching, runoff, and evaporation, further contributing to water and environmental contamination. Addressing the increasing reliance on chemical fertilizers necessitates the development

Department of Botany and Microbiology, Gurukula Kangri (Deemed to be University), Haridwar, Uttarakhand, India

***Corresponding Author:** Sandeep Kumar, Department of Botany and Microbiology, Gurukula Kangri (Deemed to be University), Haridwar, Uttarakhand, India, E-Mail: sandeepchokar@gmail.com

How to cite this article: Kumar, P., Vashistha, H., Kumar, S. 2024. PGPR Attributes and Molecular Identification of non-rhizobial Endophytes for Growth Enhancement of *Sesbania sesban* (L) Merr. J. Indian bot. Soc., Doi: 10.61289/jibs2024.05.15.123

Source of support: Nil

Conflict of interest: None.

of sustainable agricultural strategies (Prasad *et al.* 2021, Timmis and Ramos 2021). Previously, the conventional understanding was that rhizobia were exclusively responsible for symbiotic biological nitrogen fixation (BNF) within the microenvironments of root nodules. Several investigations suggest that the root nodules harbor a more diverse array of bacteria associated with nodulation (De Meyer *et al.* 2015). Various non-rhizobial endophytes (NREs) from distinct groups have been documented to play a role in facilitating the root nodulation approach and promoting plant health and growth (Preyanga *et al.* 2021). NREs infiltrate plant tissues and establish residence in the inner compartments of the plant, as documented by Hardoim *et al.* (2008). Several investigations suggest that most of the NREs have the capability to boost plant growth, elevate productivity, and provide protection against pathogens. Consequentially, divergent genera of NREs were recorded from various research, including *Agrobacterium*, *Acinetobacter*, *Bacillus*, *Enterobacter*, *Klebsiella*, *Phyllobacterium*, *Pseudomonas*, *Pantoea*, and *Serratia* (Saharan and Nehra 2011, Debnath *et al.* 2023). In terms of genera and species, PGPR displays significant diversity, with each plant species hosting a specific set of PGPR in its rhizosphere. PGPR are soil bacteria linked to plant root nodules that confer useful benefits on their hosts. Certain PGPRs are able to establish endophytic populations by penetrating the interior of the root. It is essential to identify the appropriate species of PGPR for artificial inoculation in the rhizosphere of a particular plant spp. to ensure compatibility. Researchers can introduce suitable PGPR species into the rhizosphere to supplement the existing population of naturally occurring PGPRs (Kenneth *et al.* 2019). By identifying

and characterizing these new PGPRs, we can unlock their beneficial attributes and explore their use in sustainable agricultural practices. The continuous discovery of PGPRs among different bacterial genera highlights the relevance of this study as well as the microorganisms in the rhizosphere and their potential contributions to soil health and the producibility of agriculture (Chandran *et al.* 2021, Chauhan *et al.* 2021). *Sesbania sesban* (L.) Merrill, frequently known as “Egyptian Sesban” or “Jayanti,” belongs to the leguminous plant family Fabaceae. *S. sesban* is an annual crop that can reach heights of 4 to 8 meters and achieve a growth rate of 4-5 meters within six months (Mahamat *et al.* 2021). Choosing this plant for bacterial isolation offers several advantages. *Sesbania* species, known for rapid growth and high biomass production, provide an abundant source of root nodules for successful bacterial isolation. These plants have a cooperative friendship with bacteria that fix nitrogen, offering the opportunity to isolate strains with beneficial traits. *Sesbania*'s adaptation to tropical and subtropical regions ensures the discovery of bacteria adapted to local agricultural conditions. *Sesbania* plants serve as an excellent choice for exploring bacterial diversity and their possible applications within environmentally friendly farming (Nohwar *et al.* 2019, Singh *et al.* 2021; Kumar *et al.* 2024). The aims of the present study was to isolate and screen PGPRs from *Sesbania sesban*, expanding knowledge of their bacterial diversity and potential applications, and identifying novel PGPR species for enhancing plant growth and development by effectively colonizing the rhizosphere and establishing symbiotic relationships with plant roots, while also exploring potential synergistic effects within bacterial consortia.




Figure1: Geographical representation of the bacterial isolation experimental sites

Materials and Methods

Collection of Root Nodules

Healthy, fresh, and young nodules grew at different locations in the district of Haridwar, India, and were detached from the root of *S. sesban*. Nodules were collected from three different regions: (i) Near Patanjali flyover (29.905804N, 78.002941E); (ii) Village Dhandhedi Near Roorkee bypass (29.877135N, 77.959460E); (iii) Department of Pharmaceutical Sciences, Gurukula Kangri (D.U.), Haridwar (29.91884N, 78.11771E) (Figure 1). Nodules were surface sterilized to eliminate debris, soil particles, and microbes that had attached to the outer surface. During the collection, plants at location 1 were not in the flowering stage, while those at locations 2 and 3 were flowering. Plant height and the number of nodules were recorded and are presented in Figure 2.

Isolation and Screening of Rhizobia

Root nodules were first cleaned with distilled water (DW) that had been sterilized by dipping in 4% sodium hypochlorite (NaClO) for 5 minutes, after which they were rinsed multiple times with sterile DW. After rinsing the nodules 4-5 times with sterilized distilled water to banish any remaining material, they were blot-dried and kept for later use (Jildeh *et al.* 2020). Sterilized nodules were transferred into a test tube, that had 10 ml of sterile DW. To produce a cloudy bacteroid suspension, the nodules were crumbled by using a sterile glass rod. These were streaked on Congo Red YEMA medium plates. The streaked plates were tightly sealed by using parafilm to avoid contamination, and they were incubated for a duration of 24 to 72 hours at 28 to 30°C. After 24-72 hours, bacterial colonies appear white, translucent, elevated, and mucilaginous, after being picked up, the unique colonies were moved to the YEMA plate for purification and further evaluation. The isolation of pure culture was facilitated by additional streaking, spreading, and visual analysis of colony morphology. In order to preserve and conduct further investigations, pure isolates were streaked on YEMA slants. All experiments were conducted in triplicate (Dhiman *et al.* 2019(a); Kumar and Dubey, 2022).




Figure 2: Graph represents plant height and no of nodules of different locations

Colony Morphology

The handpicked bacteria were streaked on nutrient agar medium (NAM) petri plates using the pour plate procedure and stored in an incubator at 30°C for 24 hours. Subsequently, different bacterial samples were examined for colony characteristics, including color, shape, surface features, and opacity (Kalaivanan and Mohan, 2017).

Gram Stains

Gram staining was performed by preparing smearing on the glass slide. The culture was 24h old, after it had been fixed with heat. Then after rinsing with running water, crystal violet solution was dropped on the glass slide and stood for 2 minutes. Followed by aquadest and the alcohol 95% wash for 30 seconds, it was dipped in Gram iodine solution for 2 minutes. We poured safranin over it, rinsed it with aquadest, air-dried it, and investigated it under the microscope. Only Gram-negative isolates were used for further research work (Layly *et al.* 2021).

Motility Test

Agar slants containing the preserved bacterial isolates were used for inoculation on a cover glass with the aid of an inoculum loop. The cover glass is secured onto a concavity glass slide and brought into focus under 45x objectives to examine the motility of bacterial isolates (Legesse, 2017).

Effect of Temperature, pH and Salt (NaCl) Concentrations

Different studies have been carried out that included temperature, pH, and NaCl (sodium chloride) concentrations to maximize the growth conditions of selected rhizobacterial strains. All isolates had been incubated at temperatures of 15°C, 30°C and 40°C. The medium had been prepared in a buffer to measure the effects of different pH levels (4, 7, and 9). The NAM medium was supplemented with different concentrations of salt 1%, 2%, 4%, and 5% (Basbuga *et al.* 2021).

Biochemical Assessment of a Selected Isolate

Each of the isolates undergoes a battery of biochemical tests, including, catalase, citrate, oxidase, nitrate MR, VP, hydrolysis of gelatin, starch and urea.

Catalase Test

Heme enzyme catalase breaks down the hydrogen peroxide (H_2O_2) into oxygen and water. 48-hour-old bacterial culture was placed on a slide with 1 ml of a 3% H_2O_2 solution. Bubbles signify a positive response (Wadhwa *et al.* 2017).

Citrate Test

Modifies Simmon's citrate broth media by substituting the carbon base by sodium citrate ($Na_3C_6H_5O_7$) and bromothymol blue (pH indicator). Streak the plates containing modified media with log phase culture of isolated root nodulating

bacterial strains and incubate for 48 h. Observe the plate after incubation to see that the color changes from green to blue (Hayward, 1960; Wadhwa *et al.* 2017).

Oxidase Test

The oxidase test was conducted on 24h old bacterial cultures grown on nutrient agar, where a 1 percent glucose smear was applied to Whatman's No. 1 filter paper saturated with a newly prepared tetramethyl-p-phenylenediamine dihydrochloride ($C_{10}H_{18}C_{12}N_2$) 1% aqueous solution. A positive outcome is denoted by the emergence of a purple hue within 5-10 seconds. A response between 10-60 seconds is categorized as a delayed favorable result, while any response after 60 seconds or the absence of a reaction is labeled as negative (Singh and Singh, 2015).

Nitrate Reduction Test

For nitrate reduction, the bacterial inoculated suspension into the nitrate broth at 30°C for 24h. After incubation look for nitrogen gas first then add 6-8 drops of nitrite reagents A (sulfanilic acid) and 6-8 drops of reagent B (α -naphthylamine). Observe color change within a minute, the appearance of a red color upon the add-on reagents A and B signifies the existence of nitrite, if color is no change means lack of nitrite can imply no reduction, and this can be illustrated by adding reagent C (zinc powder) to generate a red colour in the presence of nitrate. If there is no color change when reagents A, B, and C are present, it means that the nitrate has been reduced beyond nitrite and changed into different nitrogen compounds (Bhusal and Muriana, 2021).

Methyl Red-Voges Proskauer (MR-VP) Test

The selected isolates were added to sterile MR-VP broth tubes, which were afterward placed in the incubator for a duration of 2 days at 30°C. Five drops of methyl red ($C_{15}H_{15}N_3O_2$) were appended to the tubes after incubation, and any changes in color were noted. Within 10 to 15 seconds, the color changes from yellow to red, indicating a favorable result (Mazotto *et al.* 2010). After autoclaving, the Voges-Proskauer broth was cold at 25-30°C temperature. A sterilized loop was applied to inoculate the 24 hours old cultures of the selected isolates, and the medley was stored for 48 hours at 30°C. After the incubation duration, 1 ml alpha-naphthol was assorted to the broth and shaken, and then 0.5 ml of 40% potassium hydroxide (KOH) was appended and agitated. A positive outcome was considered to be the development of a red color within one hour after the reagents were added (Han *et al.* 2012; Van Thuoc *et al.* 2019).

Gelatin Hydrolysis Test

Inoculate the solid medium plates with 0.4% (w/v) gelatin using a drastically grown culture of isolated bacterial strains. Petri-plates were cultured at 28-30°C for the duration of 7 days to facilitate optimal growth. Ice bath is given to tubes

and immersed tilted in an ice bath and exposed to cold treatment, liquification of gelatin occurs which is a positive indicator of gelatin hydrolysis (Bhattacharya *et al.* 2013).

Starch Hydrolysis Test

Starch agar media pour in petri plate and inoculate spot of suspended bacterial culture on media and incubate at 24-48h at 35±2°C. After proper incubation, gram iodine solution is flooded on media, immediately blue-black color formed with starch may fade. A distinct zone surrounding bacterial colonies signifies a positive starch hydrolysis and the absence of a clear zone recorded as a negative result (Wadhwa *et al.* 2017).

Urea Hydrolysis Test

Christensen's urea agar (urea 20g, dextrose 1g, NaCl 5g, peptone 1g, phosphate monobasic 2g, phenol red 0.012g + 100 ml, DW) filter with 0.45 mm pore size filter paper and add (agar 20g+900ml, DW) prepared slants and stored at 4-8°C. Use 18-24h old fresh bacterial culture streak on media incubate at 35°C. Observe color change from 6 hours to 24 hours and every day up to 6 days. Color change 1-6 hours is rapidly positive, color change up to 6 days delayed positive and no color change is negative (Rai and Sen, 2015).

PGPR Attributes of Selected Isolates

Indole Acidic Acid (IAA) Production

Fresh nutrient broth with 24 hours old bacterial isolates (2ml) supplemented with L-tryptophan (0.2%), two blobs of orthophosphoric acid (H_3PO_4) and 4ml Salkowski reagent incubated at 25°C in the dark as long as 30 minutes. IAA production was evidenced due to the emergence of a pink colour on the basis of the incubation timing (Bhutani *et al.* 2018).

Siderophore Production

The chromium azurol-S (CAS) agar medium was utilized to production of siderophores as follows: universal method by Schwyn and Neilands, (1987). All selected isolates were growing in a succinate medium and incubated for 24 hours at 37°C with 120 rpm in a shaking incubator. The bacterial sample was centrifuged for ten minutes, at 4°C temperature, and 10,000 rpm. CAS (0.5 ml) solution (cell-free extract) was blended with an amount of 0.5 ml supernatant. The UV-Vis spectrophotometer (UV-1601) was used to measure the results at 470 nm after 20 minutes of incubation. The relation of (D/d) is the average diameter (of orange (D) to the colony (d)) was measured. The following formula was used for estimation of the percent siderophore unit (PSU):

$$\text{Siderophore Production (psu)} = \left\{ \frac{Ar - As}{Ar} \right\} 100 = \% \text{ siderophore}$$

where Ar represents the absorbance of the sample (cell free supernatant of the sample and CAS solution), and As is the absorbance of the reference (uninoculated broth and CAS solution) (Payne, 1994).




Table 1: Growth of isolates on NaCl concentration, pH, and temperature tolerance (+=less growth, ++=good growth, +++=best growth, and “-” = no growth)

Characteristics	Physiological Characters									
	Growth in NaCl (%)				Growth at pH			Growth at temp (°C)		
Isolates	1%	2%	4%	5%	4	7	9	15	30	40
SSP01	+++	+++	+++	++	++	+++	++	+++	+++	+++
SSP02	+++	+++	++	+	+	+++	+	++	+++	+++
SSP03	+++	+++	+++	++	++	+++	+++	+++	+++	+
SSP04	+++	+	-	-	+	+	-	++	++	+
SSP05	+++	+++	++	++	+	+++	-	+++	+++	+++
SSP07	+++	+	+	+	-	+	+	++	+++	++
SSP08	+++	++	+	+	+	+	+	+	+++	+
SSP09	+++	++	-	+	+	+	-	+	++	+
SSP10	+++	++	++	+	-	++	-	+	++	+
SSP12	+++	+	+	-	+	++	+	++	+++	++
SSP18	+++	++	++	+	-	++	-	+	++	+
SSP21	+++	+++	++	-	+	+	+	+	+	+
SSP22	+++	++	+	+	-	+	-	+	++	+
SSP23	+++	+	++	-	+	+	+	++	+++	++
SSP25	+++	++	+	+	+	+	-	+	++	+
SSP30	+++	+	++	-	-	+	-	+	+	+

Pot Trial Experiment

Pot trial was conducted in triplicate of each treatment. The experiment was carried out using sandy clay soil, and pots were maintained in the Department of Botany and Microbiology. Treatments involving bacterized seeds were arranged and planted in pots according to the following: T1 (control), T2- (SSP01), T3- (SSP02), T4- (SSP03), T5- (SSP05), T6- (SSP01 + SSP03), T7- (SSP02 + SSP03), T8- (SSP03 + SSP05) and T9- (SSP01 + SSP02 + SSP03). The growth parameters, such as root and shoot length and weight (both fresh and dry), were assessed by oven-drying root and shoot samples at temperatures ranging from 25 to 30°C. The treatments were formulated through microbe-microbe interactions. However, during these interactions, combinations such as SSP01+SSP05 and SSP01+SSP03+SSP05 did not exhibit synergistic growth. The effect of each isolate and consortia on seed germination was measured after 15 days of sowing (Dhiman *et al.* 2019b).

Result and Discussion

Isolation of Rhizobia

Rhizobia were detached from the root nodules of *S. sesban* grown under wild conditions. There was a significant diversification in the rhizobial spp. colonizing the roots of *S. sesban*. The current study sought to identify PGP rhizobacteria from the root nodule of *S. sesban*. Root nodulating bacteria living in symbiotic relationships can produce valuable bioactive metabolites that are likely used to improve plant

growth and fitness. In a single nodule a variety of nodulating bacteria were associated. (Diagrammatic representation of research work given in Figure 3.)

Physiological Characterization of Selected Isolates

Table 1 demonstrates the growth capabilities of selected isolates across various pH levels (ranging from 4.0 to 9.0), temperatures (15 to 40 degrees Celsius), and NaCl concentrations (1% to 5%). While the isolates generally exhibited growth within the pH range of 4.0 to 9.0. SSP01, SSP02, SSP03, SSP07, SSP08, SSP10, SSP021 and SSP23 isolates displayed particular adaptability to extreme pH conditions, thriving at both pH 4.0 and pH 9.0. Notably, bacterial isolates SSP04, SSP05, SSP07, SSP09, SSP10, and SSP18 were unable to grow at pH levels 4.0 and 9.0. Additionally, bacterial isolates SSP04, SSP09, SSP12, and SSP21 failed to grow under NaCl concentrations of 4% and 5%. Bacteria grew at temperatures ranging from 15 to 40°C. Pawar *et al.* (2014) found that at 36°C and pH 7.0, nodulating bacteria that were isolated from soybean root nodules grew well. Datta *et al.* (2015) discovered that *Rhizobium* strains isolated from *Vigna radiata*, *Glycine max* (soybean), and *Lens culinaris* (lentil) root nodules grew well at pH values of 6.0 and 7.0, a temperature of 34°C, and a concentration of 4% salt. Earlier various researches reported the morphological, physiological, and biochemical properties of rhizobial spp. and these characteristics are confirmed by Bergey's Manual of Systematic Bacteriology (Claus and Berkeley, 1986; Holt

et al. 1994; Lalitha and Immanuel, 2013; Rasool et al. 2015).

Validation of Rhizobia Isolated from Root Nodules

Seventy rhizobial isolates were isolated from *S. sesban* root nodules, all the isolates were sorted and tested to a series of authentication, such as the Congo red test, catalase, citrate, oxidase, MR-VP, nitrate reductase, and hydrolysis of gelatin, starch, and urea. On Congo red YEMA medium, 16 isolates tested, which were indicate negative results for Congo red medium (Mahmood and Athar, 2008).

Biochemical Characterization and Gram's Reaction of Selected Rhizobial Isolates

Results of Gram staining, and cell shape, of particular rhizobial isolates. It was determined that the isolates were rhizobia spp. by the production of opaque, gummy colonies, nearly round, with smooth surfaces and raised elevation. They are Gram negative rods. The rhizobial isolate passed through the catalase, citrate, oxidase, nitrate reductase, methyl red test, voges proskauer test and hydrolysis of starch, gelatin, and urea (Table 2).

Nitrogen Fixation Activity

Selected isolates were spot inoculated on the Jensen agar plate sequentially of 24 hours old culture. On Jensen's media (Nitrogen free) SSP01, SSP02, SSP03, SSP05, SSP07, SSP8 and SSP10 were able to grown and SSP04, SSP09, SSP12, SSP18, SSP21, SSP22, SSP23, SSP25 and SSP30 do not able to grow on same media.




Figure 4: Shows HCN production (A), Phosphate production (B), IAA production (C), Siderophore production (D)

Indole Acetic acid (IAA) Production

IAA synthesis was positively performed with 5 endophytic bacterial isolates (SSP01, SSP02, SSP03, SSP05, and SSP10) shows in Figure 4 (C). After completing 120 hours of incubation, the isolate SSP01 produce the highest amount of IAA (90.8 µg/ml), followed by SSP03 producing 78.6 µg/ml while SSP10 synthesize the least quantity of IAA (46.3 µg/ml). It has previously been reported that the native isolates can produce IAA when tryptophan is present (Da Silva et al. 2020). By increasing in the formation of IAA, *Enterobacter cloacae*, which was reported from *Ocimum sanctum*, additionally stimulate plant growth (Panigrahi et al. 2020). *E. xiangfangensis* (BWH6) and *E. asburiae* (STY10) synthesize IAA

Table 2: Biochemical characterization, Gram's reaction, shape and motility (R = rod, C = cocci, M = motile, NM = non motile, + = less growth, ++ = good growth, and - = no growth)

Isolates	Gram's Reaction	Shape	Motility test	Catalase	Citrate	Oxidase	Nitrate reductase	MR	VP	Hydrolysis of		
										Gelatin	Starch	Urea
SSP01	-	R	M	++	++	-	+	+	+	-	-	++
SSP02	-	R	M	++	++	++	-	-	-	-	-	-
SSP03	-	R	NM	++	+	-	-	-	-	-	-	-
SSP04	-	C	NM	-	-	+	-	-	-	++	++	++
SSP05	-	R	NM	++	++	++	-	+	+	-	++	-
SSP07	-	R	NM	+	+	-	-	-	-	+	-	-
SSP08	-	C	NM	+	+	-	-	-	+	-	-	+
SSP09	-	C	M	-	+	+	+	+	-	+	-	+
SSP10	-	R	M	+	+	+	-	+	+	-	+	-
SSP12	-	C	NM	+	-	-	+	-	-	++	+	+
SSP18	-	C	NM	-	++	+	-	-	+	-	+	-
SSP21	-	R	M	-	+	-	-	+	-	+	-	++
SSP22	-	C	NM	++	-	-	+	-	-	-	+	-
SSP23	-	C	NM	-	+	-	-	+	-	+	-	+
SSP25	-	C	NM	+	-	++	-	-	+	-	-	-
SSP30	-	R	M	+	-	-	+	-	-	+	++	-

Table 3: Qualitative and quantitative analysis of selected isolates (- = absence, + = presence, ++ = good, +++ = best)

Qualitative analysis				Quantitative analysis			
Isolates	Siderophore	HCN	IAA	Phosphate	Phosphate (mg/ml)	IAA ($\mu\text{g/ml}$)	Siderophore %
SSP01	+++	++	+++	+++	19.3	90.8	64
SSP02	++	+	++	++	13.6	70.3	45
SSP03	++	+	++	+++	14.8	78.6	43
SSP05	++	+	+	++	15.7	60.6	51
SSP07	+	-	-	-	-	-	39
SSP08	-	++	-	++	13.1	-	-
SSP10	+	++	+	+	11.9	46.3	41

in the existence of L-tryptophan were also reported (Zang *et al.* 2021). *E. hormaechei* also promote plant growth by synthesizing the IAA, ammonia, N_2 fixation, and solubilization of phosphate and potassium (Ranawat *et al.* 2021).

Siderophore Production

Six isolates (SSP01, SSP02, SSP03, SSP05, SSP07, and SSP10) produced siderophore, proved by the halo zone that has developed around the colonies (Figure 4(D)). During the growth phase of incubation, siderophore production commenced in all isolates between 24 and 120 hours. In qualitative terms, the most substantial siderophore ring formation was exhibited by SSP01 and SSP03 at 120 hours of incubation. In quantitative terms, isolate SSP01 displayed the highest siderophore production at 64%, followed by SSP03 at 51%. Isolate SSP07 exhibited the lowest siderophore production, recorded only 39%. *Enterobacter* Sp. PR14 was previously publish for synthesis of siderophore (Sagar *et al.* 2020). The production of siderophores by microbes' aids plant growth by chelating iron and facilitating the absorption of soluble compounds by the plants. *Erwinia*, *Enterobacter*, *Pseudomonas*, *Rhizobium*, and *Serratia*, has been documented as a siderophore forming bacteria (Kour *et al.* 2020). Endophytic *Bacillus*, *Enterobacter*, and *Klebsiella* were also reported to synthesize siderophore (Mowafy *et al.* 2021).

Phosphate solubilization

Six isolates (SSP01, SSP02, SSP03, SSP05, SSP08, and SSP10) exhibited phosphate solubilization, creating a halo zone with solubilization indices ranging between 1.4 to 2.0 cm. The most substantial phosphate utilization was achieved by SSP01, recording 19.3 mg/ml, followed by SSP05 with 15.7 mg/ml (Figure 4 (B), Table 3 and Figure 5). In our investigation, six isolates (SSP01, SSP02, SSP03, SSP05, SSP08, and SSP10) demonstrated phosphate solubilization. Insoluble phosphate could be dissolved in the medium by the endophytic bacteria SSP01, which is the member of *Enterobacter* genus. (Aminu *et al.* 2019). It has been documented that phosphate-solubilizing bacteria constitute a beneficial group capable of converting insoluble phosphate in the soluble form, which plants can absorb as a nutrient for

their better growth. Additionally, previous reports indicate that *Enterobacter* spp. also possess the ability for phosphate solubilization at wide range of pH and temperature (Borham *et al.*, 2017). *Acinetobacter pittii* also help in the enhancement of soybean by solubilizing phosphate in both inorganic and organic p-cycling (He and Wan, 2021).

Hydrogen Cyanide Production

For the production of volatile HCN, six isolates (SSP01, SSP02, SSP03, SSP05, SSP08, and SSP10) were cultivated in exponential growth cultures (Table 3). The isolate's color changed from yellow to light brown following a full




Figure 5: Graph shows quantitative analysis, A=phosphate solubilization, B=IAA production, and C=Siderophore production

incubation period at 30°C. The synthesis of substances that aid host plants in preventing infections through the formation of hydrogen cyanide, a secondary metabolite created when glycine is decarboxylated, is one indirect method of promoting plant growth. It works well as a biological control agent to prevent plant diseases (Admassie *et al.* 2020).

Molecular Identification of the Isolates

The most auspicious bacterial isolates were chosen to be sequenced for the 16S rRNA gene. According to Sambrook and Russell, (2001), the DNA was isolated. PCR was used in the amplification of the 16S rRNA gene by using primers 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'-TACGGYTACCTTGTTACGACTT-3'). The sequence was subjected to analysis through the BLAST within the GenBank database (National Center for Biotechnology Information), and alignment was performed via the Clustal W program. To confirm the identification of the isolates, an evolutionary tree was generated by operating MEGA 11 (Kumar *et al.*, 2018). Accession numbers *Enterobacter xiangfangensis* (SSP01) OM846632 and *Acinetobacter pittii* (SSP03) OM866163 were given to the identified bacteria. All of the chosen isolates' phylogenetic trees are displayed in Figures 6 and 7. A common species of soil bacteria called *Enterobacter* encourages the growth of plant root systems and shields host plants from harmful infections (Jha *et al.* 2011). *Enterobacter* sp. PR14 exhibits a diverse array of biocontrol traits, producing NH₃, chitinase, siderophore, HCN, and ACC deaminase. Moreover, its ability to tolerate elevated levels of various heavy metal ions positions it as a promising biocontrol agent, an excellent candidate for stress tolerance, and a valuable tool for the bioremediation of heavy metal in contaminated soil. Consequently, it serves as an ideal sustainable alternative to persistent inorganic fertilizers and fungicides (Sagar *et al.* 2020). *Acinetobacter* spp. can be found across various places and frequently occurs in the rhizosphere in numerous plants. It is important in agriculture because it acts as PGPR activities like, phosphate, potassium, siderophore, ammonia, phytohormone production, catalase assay, antibiotics, biosurfactants. Similarly, *A. pittii* has been evaluated and concluded to exhibit PGP potential and can be used alone or in combination in the sustainable cultivation of agricultural crops (Kumari *et al.* 2018; Mujumdar *et al.* 2023).

Seed Germination and Pot Trial Experiment

The maximum seed germination percentage was recorded in T 6 (69.25%), followed by T 8 (61.11%), T 9 (58.61) while minimum was recorded in T 7 (21.94%), shown in (Figure 8). Seeds bacterized with isolates SSP01 (T 2), SSP03 (T 4), SSP05 (T 5), SSP01+SSP03 (T 6), SSP03+SSP05 (T 8), and SSP01+SSP02+SSP03 (T 9) exhibited significant improvements in various plant growth parameters. The following percentage increases were recorded in contrast to the control group; The most substantial increase was




Figure 6: Phylogenetic tree of SSP01 isolate





Figure 7: Phylogenetic tree of SSP03 isolate



Photoplate 1: Pot trial experiment

observed in root length is T 6 (40%), followed by T 5 (35%), T 4 (23%), T 8 (16%), T 2 (14%) and T 9 (11%). Shoot Length: T 6 demonstrated the highest increase (41.5%), followed by T 5 (37.5%), T 4 (25.5%), T 2 (19%), T 8 (14%), and T 9 (10%). Fresh weight of root and shoot: The seeds inoculated with T 6 (39.82%) exhibited the maximum fresh weight of root and shoot, followed by T 5 (27.67%), T 4 (20.53%), T 8 (18.75%), T 2 (16.07%), and T 9 (12.5%) respectively, in comparison to

the control. Dry weight of plant root and shoot: Similar to fresh weight, the highest dry weight of root was recorded




Figure 8: Graph effect of bacterial isolates with consortium on seed germination % parameters of *S. sesban*




Figure 9: Graph represents pot trial data of all six parameters at 7th week

in seeds inoculated with T 6 (44.44%) with the order of effectiveness being T 5 (38.88%), T 4 (27.77%), T 8 (22.22%), T 2 (16.66%) and T 9 (11.11%), in comparison to the control. The highest dry weight of shoot was recorded in seeds




Figure 10: Graph show, length (A), fresh weight (B), dry weight (C) of root and shoot

Table 4: Impact of bacterial isolates with consortium on vegetative growth parameters of *S. sesban*

Treatments	Root Length (cm)	Shoot Length (cm)	Fresh Root Weight (g)	Fresh Shoot Weight (g)	Dry Weight Root (g)	Dry Weight Shoot (g)
T1 (Control)	10 ± 0.28	20 ± 0.29	0.112 ± 0.004	0.487 ± 0.0121	0.018 ± 0.0011	0.07 ± 0.0115
T2 (SSP01)	11.4 ± 0.69**	23.8 ± 0.75**	0.13 ± 0.003**	0.581 ± 0.0012*	0.021 ± 0.0023**	0.0863 ± 0.0002**
T3 (SSP02)	6.5 ± 0.23*	15.5 ± 0.20 ^{ns}	0.089 ± 0.006*	0.354 ± 0.0031*	0.012 ± 0.0019*	0.052 ± 0.0016**
T4 (SSP03)	12.3 ± 0.24**	25.1 ± 0.25**	0.135 ± 0.0023**	0.602 ± 0.0027**	0.023 ± 0.0023**	0.088 ± 0.0023*
T5 (SSP05)	13.5 ± 0.27**	27.3 ± 0.24*	0.143 ± 0.0012*	0.647 ± 0.0035**	0.025 ± 0.0023*	0.096 ± 0.0046**
T6 (SSP01+SSP03)	14 ± 0.11**	28.3 ± 0.80**	0.157 ± 0.0017**	0.698 ± 0.0058*	0.028 ± 0.0035**	0.1 ± 0.0061*
T7 (SSP02+SSP03)	6 ± 0.17*	14.8 ± 0.29**	0.099 ± 0.004*	0.282 ± 0.0017**	0.012 ± 0.0021*	0.036 ± 0.0023**
T8 (SSP03+SSP05)	11.6 ± 0.30*	22.8 ± 0.51*	0.138 ± 0.005*	0.585 ± 0.0035**	0.022 ± 0.0029**	0.082 ± 0.0036**
T9 (SSP01+SSP02+SSP03)	11.1 ± 0.40**	22 ± 0.69*	0.126 ± 0.0029*	0.553 ± 0.0116**	0.02 ± 0.0064 ^{ns}	0.079 ± 0.0052*
CD at 1%	1.81	1.23	1.51	1.62	0.08	0.05
CD at 5%	1.13	0.89	1.23	1.17	0.73	0.65

Control (non-bacterized seeds) values are mean of ten plant samples from each treatment; significance at 0.01 and 0.05 levels of analysis of variance (ANOVA),

CD, critical difference, ^{ns} Not Significant at 0.05 level of LSD as compared to control

** Significant at 0.01 level of LSD in comparison to control

* Significant at 0.05 level of LSD in comparison to control

inoculated with T6 (42.85%) with the order of effectiveness being T5 (37.14%), T4 (25.71%), T2 (18.53%), T8 (17.14%), and T9 (12.85%), as compared to the control. Notably, the most significant enhancement was observed in SSP01+SSP03 (T6) while the least effect was seen in SSP02+SSP03 (T7). Pot trial experiment is shown in photoplate 1, Figures 9 and 10 (A, B, and C) and statistically in Table 4. *Enterobacter* sp. and *Acinetobacter* sp. were previously reported from *S. sesban* (Jobby *et al.* 2016; Fanuel *et al.* 2022) both bacterial spp. show PGP attributes (Kumar and Dubey, 2022; Kour and Yadav, 2023). Microbial consortiums have been shown to be as effective as versus chemical fertilizers (e.g., urea (NH₂CONH₂), diammonium phosphate (DAP, (NH₄)₂HPO₄), muriate of potash (MoP), and so on) in raising crops under abiotic stress situations of soil salt concentration (Maheshwari *et al.* 2023).

Conclusion

In conclusion, the selected bacteria in both forms, individual and in consortium, had a weighty sequel on the growth of *S. sesban*. The outcome of the pot trial shows that the plant height and weight (fresh & dry) of root & shoot of *S. sesban* increased observable due to inoculation with the multi-strain bacterial consortium. The potential of the microbial consortium to exert diverse biological effects and trigger the growth of *S. sesban* was recorded. Treatment 6 (SSP01+SSP03) possesses the potential to effect plant growth by various PGP mechanisms. The findings of our research also suggest valuable insights into the diversity and potential applications of beneficial bacteria from *S. sesban* root nodules. We observed, that bioinoculants enhance plant growth and can reduce reliance on chemical fertilizers. However, further investigations were needed to elucidate the specific mechanisms by which these bacteria promoted plant growth and optimize their application techniques in different agricultural systems.

Acknowledgement

The authors wish to thank the head of the Department of Botany and Microbiology, Gurukula Kangri (Deemed to be University), Haridwar, for providing the necessary laboratory facilities.

Author contribution

P.K. executed the experiments and drafted the manuscript. H.V. collected data and planned research work and S.K. corrected, and finalized the manuscript, and read and approved the final manuscript.

Conflict of interest

The authors declare no competing interests.

References

Adhikari P and Pandey A 2019 Phosphate solubilization potential of endophytic fungi isolated from *Taxus wallichiana* Zucc. roots. *Rhizosphere*, **9**: 2–9.

- Admassie M, Assefa F and Alemu T 2020 Different mechanism, application, methods and properties of plant growth promoting rhizosphere bacteria. *Appl. Microbiol.*, **6**:177-188.
- Ahmed N and Turchini G M 2021 The evolution of the blue-green revolution of rice-fish cultivation for sustainable food production. *Sustain. Sci.*, **16**:1375–1390.
- Aleminew A and Alemayehu M 2020 Soil fertility depletion and its management options under crop production perspectives in Ethiopia: A review. *Agric. Rev.*, **41**(2): 91-105.
- Aminu Y, Maryam M G and Kabiru S A 2019 Hormonal response of gibberellin (Ga3), grafting and seasonal variations on growth and yield parameters on Okra (*Abelmoschus esculentus*). *Am. Int. J. Biol. Life Sci.*, **1**(2): 33-39.
- Bakker A W and Schippers B O B 1987 Microbial cyanide production in the rhizosphere in relation to potato yield reduction and *Pseudomonas* spp.-mediated plant growth-stimulation. *Soil Biol. Biochem.*, **19**(4): 451–457.
- Basbuga S, Basbuga S, Yayla F, Mahmoud A M and Can C 2021 Diversity of rhizobial and non-rhizobial bacteria nodulating wild ancestors of grain legume crop plants. *Int. J. Microbiol.*, **24**: 207-218.
- Bhattacharya C, Deshpande B and Pandey B 2013 Isolation and characterization of rhizobium sp. from root of legume plant (*Pisum sativum*) and its antibacterial activity against different bacterial strains. *J. Agric. Sci.*, **3**(4):138-141.
- Bhusal A and Muriana P M 2021 Isolation and characterization of nitrate reducing bacteria for conversion of vegetable-derived nitrate to 'natural nitrite'. *Appl. Microbiol.*, **1**(1):11-23.
- Bhutani N, Maheshwari R, Negi M and Suneja P 2018 Optimization of IAA production by endophytic *Bacillus* spp. From *Vigna radiata* for their potential use as plant growth promoters. *Isr. J. Plant Sci.*, **65**: 83–96.
- Borham A, Belal E and Metwaly M 2017 Phosphate solubilization by *Enterobacter cloacae* and its impact on growth and yield of wheat plants. *J. Sustain. Agric. Sci.*, **43**(2): 89-103.
- Chandran H, Meena M and Swapnil P 2021 Plant growth-promoting rhizobacteria as a green alternative for sustainable agriculture. *Sustain*, **13**(19):1-30.
- Chauhan A, Saini R and Sharma J C 2021 Plant growth promoting rhizobacteria and their biological properties for soil enrichment and growth promotion. *J. Plant Nutr.*, **45**(2): 273-299.
- Claus D and Berkeley R C W 1986 Genus *Bacillus*. In: P.H.A. Sheath (Ed.), *Bergey's Manual of Systematic Bacteriology*, Vol. III. Williams and Wilkins, Baltimore, 1105-39.
- Da Silva, MJC Junior, SFP Junio, KF Nascimento, VX de Medeiros, AS da Silva, SJC and Sant'Ana, AEG 2020 IAA production of indigenous isolate of plant growth promoting rhizobacteria in the presence of tryptophan. *Aust. J. Crop Sci.*, **14**:537-544.
- Datta A, Singh R K and Kumar S 2015 Isolation, characterization and growth of *Rhizobium* strains under optimum conditions for effective biofertilizer production. *Int. J. Pharmacol.*, **32**:199-208.
- De Meyer S E, De Beuf K, Vekeman B and Willems A 2015 A large diversity of non-rhizobial endophytes found in legume root nodules in Flanders (Belgium). *Soil Biol. Biochem.*, **83**: 1–11.
- Debnath S, Chakraborty S, Langthasa M, Choure K, Agnihotri V, Srivastava A and Pandey P 2023 Non-rhizobial nodule endophytes improve nodulation, change root exudation pattern and promote the growth of lentil, for prospective

- application in fallow soil. *Front. Plant Sci.*, **14**(1152875): 1-19.
- Dhiman M, Dhiman V K, Rana N and Dipta B 2019(a) Isolation and characterization of *Rhizobium* associated with root nodules of *Dalbergia sissoo*. *Int. J. Curr. Microbiol.*, **8**:1910-1918.
- Dhiman S, Dubey R C, Baliyan N, Kumar S and Maheshwari DK 2019(b) Application of potassium-solubilising *Proteus mirabilis* MG738216 inhabiting cattle dung in improving nutrient use efficiency of *Foeniculum vulgare* Mill. *Environ. Sustain.*, **2**: 401-409.
- Dubey R C and Maheshwari D K 2012 *Practical Microbiology*. S. Chand and Co. New Delhi.
- Fanuel K, David M and Asenath A 2022 Diversity of Culture dependent endophytic bacteria isolated from leguminous agroforestry trees in western Kenya. *Res. Sq.*, 1-13.
- Han M, Luo W, Gu Q and Yu X 2012 Isolation and characterization of a keratinolytic protease from a feather-degrading bacterium *Pseudomonas aeruginosa* C11. *Afr. J. Microbiol. Res.*, **6**(9): 2211-2221.
- Hardoim P R, van Overbeek L S and Van Elsland J D 2008 Properties of bacterial endophytes and their proposed role in plant growth. *Trends Microbiol.*, **16**(10): 463-471.
- Hayward A C 1960 A method for characterizing *Pseudomonas solanacearum*. *Nature*, **186**(4722): 405-406.
- He D and Wan W 2021 Phosphate-solubilizing bacterium *Acinetobacter pittii* gp-1 affects rhizosphere bacterial community to alleviate soil phosphorus limitation for growth of soybean (*Glycine max*). *Front. Microbiol.*, **12**(737116):1-12.
- Holt J G, Krieg N R, Sneath P H, Staley J T and Williams S T 1994 *Bergey's manual of determinative bacteriology*. 9th. Baltimore: William and Wilkin
- Jensen H L 1951 Notes on biology of *Azotobacter*. *J. Appl. Microbio.*, **14**: 89-94.
- Jha C K, Aeron A, Patel B V, Maheshwari D K and Saraf M 2011 *Enterobacter*: Role in Plant Growth Promotion. In: Maheshwari, D. (eds) *Bacteria in Agrobiolgy: Plant Growth Responses*. Springer, Berlin, Heidelberg, 159-182.
- Jildeh Z B, Kirchner P, Oberlander J, Vahidpour F, Wagner P H and Schoning M J 2020 Development of a package-sterilization process for aseptic filling machines: a numerical approach and validation for surface treatment with hydrogen peroxide. *Sens. Actuator A Phys.*, **3**(111691): 1-11.
- Jobby R, Shah K, Shah R, Jha P and Desai N 2016 Differential expression of antioxidant enzymes under arsenic stress in *Enterobacter* sp. *Environ. Prog. Sustain.*, **35**(6): 1642-1645.
- Kalaivanan S D and Mohan V 2017 Screening and molecular characterization of salt tolerant bio-control bacterial isolates from *Casuarina equisetifolia* rhizosphere soil. *Asian J. Plant Pathol.*, **11**(4):156-166.
- Kenneth O C, Nwadike E C, Kalu A U and Unah U V 2019 Plant growth promoting rhizobacteria (PGPR): a novel agent for sustainable food production. *Am. J. Agric. Biol. Sci.*, **14**(35):1-20.
- Khan B A, Nadeem M A, Nawaz H, Amin M M, Abbasi G H., Nadeem M and Ayub M A 2023 Pesticides: Impacts on agriculture productivity, environment, and management strategies. In: Aftab, T. (eds) *emerging contaminants and Plants. Emerging contaminants and associated treatment technologies*. Springer, Cham., 109-134.
- Kour D, Kaur T, Devi R, Rana K L, Yadav N, Rastegari A A and Yadav A N 2020 Biotechnological applications of beneficial microbiomes for evergreen agriculture and human health. *I. J. Biotec. Bioeng.*, 255-279.
- Kour D and Yadav A N 2023 Alleviation of cold stress in wheat with psychrotrophic phosphorus solubilizing *Acinetobacter rhizosphaerae* EU-KL44. *Braz. J. Microbiol.*, **54**(1): 371-383.
- Kumar P, Vashistha H and Kumar S 2024 Phytochemistry, pharmacological, soil amelioration, PGPR attributes, traditional, modern and future prospects of *Sesbania sesban* (L) Merr. *J. Indian Bot. Soc.*, 104(01): 1-12.
- Kumar N and Dubey RC 2022 Plant growth-promoting attributes of an endophyte *Enterobacter roggenkampii* BL502 isolated from *Barleria lupulina* Lindl. *Org. Agric.*, **12**(1): 137-145.
- Kumar S, Stecher G, Li M, Knyaz C and Tamura K 2018 MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.*, **35**(6):1547-1549.
- Kumari P, Meena M and Upadhyay R S 2018 Characterization of plant growth promoting rhizobacteria (PGPR) isolated from the rhizosphere of *Vigna radiata* (mung bean). *Biocatal. Agric. Biotechnol.*, **16**:155-162.
- Lalitha S and Immanuel SP 2013 Biochemical characterization of *Rhizobium* and its impact on black gram and green gram plants. *Int. J. Curr. Sci.*, **9**:1-6.
- Layly IR, Meryandini A, Helianti I and Astuti RI 2021 Identification of lipase producing bacteria from palm oil sewage sludge processing plant at Malimping, Banten, Indonesia. *Biodiversitas*, **22**(10): 4512-4524.
- Legesse D Y 2017 Optimization and partial characterization of bacillus protease isolated from soil and agro-industrial wastes. *Int. J. Nutr. Food Sci.*, **6**(1): 31-38.
- Mahamat O B, Younes S, Otchom B B, Franzel S, Mahamat A D O and Fadoua A 2021 Ecology, morphology, distribution, and use of *Sesbania tchadica* (*Sesbania sesban*) from the Republic of Chad: A review. **1**:1-26.
- Maheshwari D K, Kumar S, Kumar P, Kumar S and Dheeman S 2023 Effect of Microbial Consortium Vs. Perfected Chemical Fertilizers for Sustainable Crop Growth. In: Maheshwari D K and Dheeman S (Eds) *Design and development of microbial consortia, Sustainable Agrobiolgy. Microorganisms for Sustainability*. Springer, Singapore. **(43)**: 319-337.
- Mahmood A and Athar M 2008 Cross inoculation studies: response of *Vigna mungo* to inoculation with rhizobia from tree legumes growing under arid environment. *Int. J. Environ. Sci. Technol.*, **5**:135-139.
- Mazotto A M, Lage-Cedrola S M, Lins U, Rosado A S, Silva K T, Chaves J Q, Rabinovitch L, Zingali R B and Vermelho A B 2010 Keratinolytic activity of *Bacillus subtilis* AMR using human hair Lett. *Appl. Microbiol.*, **50**(1): 89-96.
- Mowafy A M, Fawzy M M, Gebreil A and Elsayed A 2021 *Endophytic Bacillus, Enterobacter, and Klebsiella* enhance the growth and yield of maize. *Acta agriculturae scandinavica, section B. Soil and Plant Sci.*, **71**(4): 237-246.
- Mujumdar S, Bhoyar J, Akkar A, Hundekar S, Agnihotri N, Jaybhay P and Bhuyan S 2023 *Acinetobacter*: A versatile plant growth-promoting rhizobacteria (PGPR). In *plant-microbe interaction-recent advances in molecular and biochemical approaches*, *Academic Press*, **(1)**: 327-362.
- Nohwar N, Khandare R V and Desai N S 2019 Isolation and characterization of salinity tolerant nitrogen fixing bacteria from *Sesbania sesban* (L) root nodules. *Biocatal. Agric. Biotechnol.*, **21**: 1-6.

- Panigrahi S, Mohanty S and Rath C C 2020 Characterization of endophytic bacteria *Enterobacter cloacae* MG00145 isolated from *Ocimum sanctum* with Indole Acetic Acid (IAA) production and plant growth promoting capabilities against selected crops. *S. Afr. J. Bot.*, **13**:17–26.
- Pawar V A, Pawar P R, Bhosale A M and Chavan S V 2014 Effect of *Rhizobium* on seed germination and growth of plants. *J. Acad. Ind. Res.*, **3**:84-88.
- Payne S M (1994) Detection, isolation, and characterization of siderophores. *Methods Enzymol.*, **235**: 329–344.
- Prasad S, Malav L C, Choudhary J, Kannojiya S, Kundu M, Kumar S and Yadav AN 2021 Soil microbiomes for healthy nutrient recycling. Current trends in microbial biotechnology for sustainable agriculture, *Springer* Singapore, 1-21.
- Preyanga R, Anandham R, Krishnamoorthy R, Senthilkumar M, Gopal N O, Vellaikumar A and Meena S 2021 Groundnut (*Arachis hypogaea*) nodule *Rhizobium* and passenger endophytic bacterial cultivable diversity and their impact on plant growth promotion. *Rhizosphere*, **17**(100309): 1-11.
- Rai R and Sen A 2015 Biochemical characterization of french bean associated rhizobia found in north Bengal and Sikkim. *J. Acad. Ind. Res.*, **4**(1):10-18.
- Ranawat B, Bachani P, Singh A and Mishra S 2021 *Enterobacter hormaechei* as plant growth-promoting bacteria for improvement in *Lycopersicum esculentum*. *Curr. Microbiol.*, **78**:1208-1217.
- Rasool S, Sharma B and Rasool S 2015 Isolation and characterization of *Rhizobium* sp. from a wild legume from BGSBU campus of District Rajouri, J&K. *Int. J. Agric. Sci.*, **5**: 407-413.
- Sagar A, Riyazuddin R, Shukla P K, Ramteke P W and Sayyed R Z 2020 Heavy metal stress tolerance in *Enterobacter* sp. PR14 is mediated by plasmid. *Ind. J. Exp. Biol.*, **58**:115-121.
- Saharan B S and Nehra V 2011 Plant growth promoting rhizobacteria: a critical review. *Life Sci. Med. Res.*, **21**(1): 1-30.
- Sambrook J and Russell D 2001 Molecular Cloning: A Laboratory Manual, 3rd edn. Cold Spring Harbour Laboratory Press: Cold Spring Harbour, NY
- Schwyn B and Neilands J B 1987 Universal chemical assay for the detection and determination of siderophores. *Anal. Biochem.*, **160**(1): 47–56.
- Shafiq S, Haq M Z U, Shahbaz A, Shafique S, Riaz M, Hanif M T and Hassan G Z 2024 Agrochemicals and climate change. In water-soil-plant-animal nexus in the era of climate change, IGI Global, 49-77.
- Singh A K and Singh G 2015 A study of multiple heavy metal tolerance in root nodulating bacteria. *Int. J. Res. Dev. Pharm. Life Sci.*, **4**(5):1713-1721.
- Singh K, Gera R, Sharma R, Maithani D, Chandra D, Bhat M A and Bhatt P 2021 Mechanism and application of *Sesbania* root-nodulating bacteria: an alternative for chemical fertilizers and sustainable development. *Arch. Microbiol.*, **203**(4):1259-1270.
- Timmis K and Ramos J L 2021 The soil crisis: the need to treat as a global health problem and the pivotal role of microbes in prophylaxis and therapy. *Microb. Biotechnol.*, **14**(3): 769-797.
- Van Thuoc D, Hien T T and Sudesh K 2019 Identification and characterization of ectoine-producing bacteria isolated from can gio mangrove soil in Vietnam. *Ann. of Microbiol.*, **69**(8): 819-828.
- Wadhwa Z, Srivastava V, Rani R, Kanchan T and Jangra S 2017 Isolation and characterization of *Rhizobium* from chickpea (*Cicer arietinum*). *Int. J. Curr. Microbiol. App. Sci.*, **6**(11): 2880-2893.
- Weller D M and Cook R J 1983 Suppression of take-all of wheat by seed treatments with fluorescent pseudomonads. *Phytopathol.*, **73**(3): 463-469.
- Zhang B X, Li P S, Wang Y Y, Wang J J, Liu X L, Wang X Y and Hu X M 2021 Characterization and synthesis of indole-3-acetic acid in plant growth promoting *Enterobacter* sp. *RSC Adv.*, **11**(50):31601-31607.