



RESEARCH ARTICLE

Corynandra felina (L.f.) Cochrane and Iltis (Cleomaceae), a new record to the angiospermic flora of Kerala

Sangeetha Suran^{1,2*} Sunilkumar T. Pillai²

Abstract

Corynandra felina (L.f.) Cochrane and Iltis, native to the range of peninsular India, is reported here as a new distributional record to the flora of Kerala from Chinnar Wildlife Sanctuary, Idukki district.

Keywords: *Cleome felina*, Chinnar Wildlife Sanctuary, Kerala, New record.

Introduction

Cleome L. sensu lato is the largest genus of the family Cleomaceae, with 200–250 species, distributed all over the world (POWO, 2022, Mabberley 2017). In 1753, Linnaeus established the genus with eight species under the family Capparidaceae, which was followed by subsequent researchers (De Candolle 1824, Bentham and Hooker 1862, Boissier 1867, Oliver, 1868, Hooker 1875, Hedge and Lamond, 1970, Jafri 1973). Meanwhile, Pax (1891) divided the family Capparidaceae into two subfamilies viz. Capparidoideae and Cleomoideae, and placed the genus *Cleome* under the subfamily Cleomoideae. Further, Airy Shaw (1965) raised the status of Cleomoideae to family level, Cleomaceae and which is supported by the recent report of Hall *et al.*, (2002) and Angiosperm Phylogeny Group System of Classification (2016).

Recently, the genus *Cleome* was segregated into number of genera, and the genus *Corynandra* Schrad. ex Spreng. has

been reinstated based on the species *Cleome chelidonii* L.f. (Iltis and Cochrane, 2007, Cochrane and Iltis, 2014). The genus *Corynandra* is delineated from its other allied genera by means of the closed aestivation of the flowers, large number of stamens, the adaxial (if not all) filaments of which are more or less apophysate, typically erect capsules with persistent valves, that dehisce from the apex and open seed cleft (Cochrane and Iltis 2014). In India, the genus *Cleome* sensu lato is represented by seventeen taxa (Sundara Raghavan, 1993) and out of which, ten were recorded from Kerala part of Western Ghats (Sasidharan, 2010).

Materials and Methods

As a part of first author's Ph.D. work on south Indian *Cleome* sensu lato, several field exploration trips were conducted during 2019-2021. While exploring the Chinnar Wildlife Sanctuary regions of Idukki district in Kerala, the authors came across an unknown population of *Cleome* grown in the rocky places, which was morphologically distinct from all other known species recorded so far from Kerala. On critical examination and perusal of relevant literature (Hooker, 1875, Gamble, 1921; Sundara Raghavan 1993; Sanchez-Acebo 2005), the specimen was identified as *Corynandra felina* (L.f.) Cochrane and Iltis (basionym: *Cleome felina* L.f.), a species native of peninsular India, which is yet to be figured in the floristic account of Kerala state.

During the study, it is also noticed that, one online database (<https://indiabiodiversity.org>) reported the distribution of *C. felina* from Kerala. Further studies perusal of the detailed literature survey on published floras (Manilal and Sivarajan, 1982; Manilal, 1988; Ramachandran and Nair, 1988; Sasidharan and Sivarajan, 1995; Sivarajan and Mathew, 1997; Sasidharan, 1998; 1999; 2002; 2004; 2013; Mohanan

¹Devaswom Board Pampa College, Parumala, Pathanamthitta District, Kerala, India.

²Department of Postgraduate studies and Research in Botany, Sanatana Dharma College Alappuzha, Kerala, India.

***Corresponding Author:** Sangeetha Suran, Devaswom Board Pampa College, Parumala, Pathanamthitta District, Kerala, India, E-Mail: sangeethas0808@gmail.com

How to cite this article: Sangeetha S and Sunilkumar T (2023). *Corynandra felina* (L.f.) Cochrane and Iltis (Cleomaceae), a new record to the angiospermic flora of Kerala. J. Indian bot. Soc. 103(2):143-145. Doi: 10.5958/2455-7218.2023.00013.X

Source of support: Nil

Conflict of interest: None.

Table 1: Comparison of distinguishing characters of *Corynandra chelidonii* and *C. felina*

Characters	<i>Corynandra chelidonii</i>	<i>Corynandra felina</i>
Stem	Erect	Procumbent
Leaves	Dimorphic, 5 – 9 foliate	Monomorphic, trifoliate
Texture of leaves	Membraneous	Coriaceous
Filament	White	Violet
Capsules	Cylindrical, 6–9 cm long	Flat, 3.5 – 4.2 cm long
Seeds	Without elaiosome	With elaiosome

and Sivadasan, 2002; Nayar, 2006; Sunil and Sivadasan 2009;; Nayar *et al.*, 2014) and herbarium consultation in different herbaria (CAL, MH, CALI, TBGT, KFRI, KUBH, CMPR, CATH, RHT) confirmed that, there is no specimen of this species is either mentioned in any floras or deposited in any of these herbaria from Kerala. The detailed description, distribution, field photographs (Figure 1) and dissected photo plate (Figure 2) are provided here for the easy identification of the species.

Result and Discussions

Corynandra felina (L.f.) Cochrane and Iltis, Novon 23 (1): 2014.

Type: Sri Lanka, J.G. Koenig 1777, (LINN 850:9, digital image!). Basionyms: *Cleome felina* L.f., suppl. 300. 1781. *Polanisia felina* (L.f.) DC; Prodr. 1: 242. 1824.

Description

Annual herb, 15–50 cm high. Stem erect or procumbent, branched, sub-cylindrical, green, appressed white hairy. Leaves exstipulate, palmately compound, trifoliate, petiolate; petiole sub-cylindrical with dorsal groove, 0.8–1.3




Figure 2: *Corynandra felina* (L.f.) Cochrane and Iltis. A. Habit; B. Flowering twig; C. Leaf adaxial; D. Leaf abaxial; E. Single flower; F. Sepal; G. Petal; H. Stamens; I. Single stamen; J. Gynoecium; K. Capsule; L. Opened capsule with seed containing elaiosome; M. seed.

cm long, pubescent; petiolules absent; leaflets obovate or oblanceolate, upper leaflet large, 1.0–1.8 × 0.5–0.9 cm, lateral leaflets small, 0.8–1.5 × 0.3–0.6 cm, apex round, sub-acute, obtuse or retuse, margin ciliated, base cuneate, pubescent on both sides, coriaceous. Flowers axillary, bisexual, zygomorphic, tetramerous, pedicellate; pedicel cylindrical, 1–1.5 cm long, green, hairy. Sepals 4, free, lanceolate, ca. 0.4 × 0.1 cm, apex acute, margin ciliated, green, hairy, deciduous. Petals 4, free, oblong, not clawed, 0.8 – 0.9 × ca. 0.4 cm, apex acute, margin entire, pink or purple, abaxial surface hispid, adaxial surface smooth, glabrous. Stamens numerous, 45–55; filaments apophysate, ca. 0.7 cm long, purple, glabrous; anther ca. 0.1 cm long, ovoid, yellow, curved after anthesis. Gynoecium flat, oblong; ovary ca. 0.3 × 0.2 cm across, green, glabrous; style short, ca. 0.2 cm long, purple, glabrous; stigma entire, white, glabrous. Capsule flat, oblong 3.5–4.2 cm long, green, drooping, beaked at apex. Seed 14–16 seeds per capsule, horse shoe shaped, tubercled, brown or black, cleft open, elaiosome present.

Flowering & fruiting: May–September.

Specimens examined: India: **Andra Pradesh:** Parchur Prakasam, Vengankunta 29.09.2008, *B. Sadasivaiah and Khadar Basha s. n.* (BSID0002827); Bhumayyagiri Palli, 15.10.2009, *K. Rajakullai Swamy and Rajasekhar s.n.* (BSID002829). **Kerala:** Idukki district, Chinnar Wildlife Sanctuary, 10.3068° N, 77.2060° E, 31.05.2021, *Sangeetha S. and T. Sunilkumar* 10594 (KUBH) (Kerala University Botany Herbarium, Kariavattom, Thiruvananthapuram). **Tamil Nadu:** Coimbatore, Forest College Compound, 468 m, 07.05.1968, *M. Chandrabose* 29880 (MH); Dharmapuri, Addappullam forest, 700 m, 25.07.1978, *E. Vajravelu* 57969 (MH); Madras Presidency,




Figure 1: *Corynandra felina* (L.f.) Cochrane and Iltis. A. Habit; B. Flower; C. Plant with Fruit; D. Location of collection.

Trichinopoly, Satnur 09.05.1944, *James Sinclair* 3520 (E00693752); Ramanadapuram, Virudanagar, 09.09.1949, *Ramamurthy* 93792 (MH); Salem, Pennagaram, s.d 400 m, *E. Vajravelu* 20731 (MH); Virudanagar, 17.09.1956, *Joseph* 59 (KUBH).

Notes: *Corynandra felina* is differed from its closely resembled species *Corynandra chelidonii* (L.f.) Cochrane & Iltis, distinguishing characters are provided in Table 1. The present collection of *C. felina* from Chinnar Wildlife Sanctuary is a new record to the angiosperm flora of Kerala. Rama Rao (1914) mentioned the name *Cleome felina* in his publication 'Flowering Plants of Travancore'. But presently most of the areas previously considered as Travancore is coming under the state of Tamil Nadu and there is no scientific evidence from Kerala to justify his statement in any of the herbaria. Hence it is confirmed that, the occurrence and distribution of *C. felina* from the state of Kerala is a wrong statement without any scientific evidence. Perusal of this data, it is clear that the present collection *Corynandra felina* from Chinnar Wildlife Sanctuary is a new record to the angiosperm flora of Kerala.

Acknowledgement

Authors are thankful to the Department of Botany, Kerala University, Kariavattom for providing the support. Authors are also express sincere gratitude to The Manager, Principal and staff of Postgraduate Studies and Research Department of Botany S.D. College, Alappuzha for providing the facilities and Dr. K.M. Prabhukumar, Senior Scientist, CSIR-National Botanical Research Institute, Lucknow for confirming the identity of the taxa and improving the quality of the manuscript.

References

- Airy shaw H K 1965 Diagnosis of new families, new names etc. for the seventh edition of Willis's "Dictionary" Kew Bulletin 18 256.
- APG (Angiosperm Phylogeny Group) IV (2016). An update of the angiosperm phylogeny group for the orders and families of flowering plants. Bot. J. Linn. Soc. 181: 1–20.
- Bentham G & J D Hooker 1862 Genera plantarum ad exemplaria imprimis in herbariis kewensibus servata, Reeve.
- Boissier E 1867 Flora Orientalis. vol. 1-5 Genevae et Basileae.
- Cochrane T S and Iltis H H 2014 Studies in the Cleomaceae VII five new combinations in *Corynandra*, an earlier name for *Arivela*. Novon, A Journal for Botanical Nomenclature 23(1) 21–26.
- De candolle A P 1824 Prodromus systematics Naturalis regni vegetabilis. Sumptibus Sociorum Treuttel et Wurtz, Paris.
- Gamble J S 1919 Flora of the Presidency of Madras. Newman and Adlard, London.
- Hall J C, Sytsma K J and Iltis H H 2002 Phylogeny of Capparaceae and Brassicaceae based on chloroplast sequence data. American Journal of Botany 89 1826–1842.
- Hedge I and Lamond J 1970 Flora Iranica. Akademische Druck-U- verlagsanstalt Graz Austria.
- Hooker J D 1875 Flora of British India. L Reeve & Co. Ltd., London.
- Iltis H H and Cochrane T S 2007 Studies in the Cleomaceae V: a new genus and ten new combinations for the Flora of North America. Novon A Journal for Botanical Nomenclature 17(4) 447–451.
- IPNI 2022 International Plant Names Index. Published on the Internet <http://www.ipni.org>, The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Botanic Gardens. Retrieved 05 October 2022.
- Jafri S M H 1973 Flora of Pakistan. Dept. of Botany University of Karachi, Pakistan.
- Linnaeus C 1753 Species Plantarum. British Museum (Natural History), London.
- Mabberley D J 2017 A portable dictionary of the Vascular plants, their classification and uses. Cambridge University Press, Cambridge.
- Manilal K S. and Sivarajan V V 1982 Flora of Calicut. Bishen Singh Mahendra Pal Singh Dehra Dun.
- Manilal K S 1988 Flora of Silent Valley. Department of Science & Technology, (Government of India), Calicut.
- Mohan N and Sivadasan M 2002 Flora of Agasthyamala. Bishan Singh Mahendra Pal Singh Dehra Dun.
- Nayar T S, Beegam A R, Mohanan N and Rajkumar G 2006 Flowering Plants of Kerala. TBGRI, Thiruvananthapuram, Kerala.
- Nayar T S, Beegam A R and Sibi M 2014 The Flowering plants of Western Ghats in India. TBGRI, Thiruvananthapuram, Kerala.
- Oliver D 1868 Flora of Tropical Africa. L Reeve and Co. Ltd. Ashford, Kent.
- Pax F 1891 Capparidaceae. Die naturlichen Pflanzenfamilien 2 209–236.
- POWO 2022 "Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://www.plantsoftheworldonline.org/> Retrieved 26 October 2022.
- Ramachandran V S and Nair V J 1988 Flora of Cannanore. Botanical Survey of India, Calcutta.
- Rama R 1914 Flowering Plants of Travancore. Govt. press, Thiruvananthapuram, Kerala.
- Sanchez-Acebo, L 2005 A phylogenetic study of the new world Cleome (Brassicaceae, Cleomoideae). Annals of the Missouri Botanical Garden 92 179–201.
- Sasidharan N and Sivarajan V V 1995 Flowering plants of Thrissur forests. Scientific Publishers, Jodhpur.
- Sasidharan N 1998 Studies on the flora of Periyar Tiger Reserve. Kerala Forest Research Institute Peechi, Thrissur.
- Sasidharan N 1999 Study on the flora of Chinnar Wildlife Sanctuary. KFRI Research Report No. 167. Kerala Forest Research Institute Peechi, Thrissur.
- Sasidharan N 2002 Floristic studies in Parambikulam Wildlife Sanctuary. KFRI Research Report 246:165–167.
- Sasidharan N 2004 Biodiversity documentation for Kerala Part 6 Flowering Plants. Kerala Forest Research Institute, Kerala.
- Sasidharan N 2013 Flowering plants of Kerala CD-ROM ver 2.0. Kerala Forest Research Institute, Kerala.
- Sivarajan V V and Philip Mathew 1997 Flora of Nilambur. Bishen Singh and Mahendra Pal Singh, Dehra Dun.
- Sundara Raghavan R 1993 Flora of India Vol. 2. Botanical Survey of India, Howrah.
- Sunil C N and Sivadasan M 2009 Flora of Alappuzha District Kerala India. Bishen Singh and Mahendra Pal Singh, Dehra Dun.