https://doi: 10.5958/2455-7218.2022.00114.0

**ORIGINAL RESEARCH ARTICLE** 



# Account of the marine macroalgae from the northern coastal district of Andhra Pradesh, India

Aron Santhosh Kumar Y1\*, Palanisamy M2

### Abstract

The reports on the occurrence of the marine macroalgae from Andhra Pradesh are sporadic and merely reported from the middle part (Visakhapatnam district) of the state and the rest of the southern and northern parts of the coastal areas were not yet be covered. Hence, the present attempt was subjected to instigate the details on the occurrence of marine macroalgae from the northern coastal districts (Vizianagaram and Srikakulam) of Andhra Pradesh during March 2017 - August 2019. A sum of 57 taxa of marine macroalgae belongs to 27 genera of 20 families under 15 orders collected from the 22 localities. Among them, class Rhodophyceae (46%) is the predominant group in species composition followed by Chlorophyceae (35%) and Phaeophyceae (19%). Also, the present study highlights the correlation of ecological parameters of the coastline with the distribution and diversity of seaweeds in Andhra Pradesh. Further, it reveals the occurrence of 16 taxa reported for the first time from the coastline of Andhra Pradesh.

Keywords: Andhra Pradesh, Additions, Diversity, Northern districts, Physio-chemical parameters.

## Introduction

Marine macroalgae (seaweeds) are the key components and regulatory elements of the aquatic ecosystem with significant ecological and economic roles. They are exclusively found in marine habitats including estuaries, lagoons, brackish water, salt marshes, and mangroves ranging from tropical to temperate counties. They have nonvascular plant bodies differentiated into holdfast, stipe, and fronds with various forms of unicellular to multicellular macroscopic filamentous or thallus (Lightowler and Davies 2002). They constitute about 11, 000 species, of which includes 7, 200 taxa of Rhodophyceae, 2, 000 taxa of Phaeophyceae, and

<sup>1</sup>Botanical Survey of India, Southern Regional Centre, TNAU Campus,Coimbatore – 641 003, Tamil Nadu, India,

2Central National Herbarium, Botanical Survey of India, Botanic Garden, Howrah – 711 103, West Bengal, India

\*Corresponding Author: Aron Santhosh Kumar Y, Botanical Survey of India, Southern Regional Centre, TNAU Campus, Coimbatore – 641 003, Tamil Nadu, India, E-Mail: santhosharon@gmail.com

**How to cite this article:** Kumar YAS, Palanisamy M (2023). Account of the marine macroalgae from the northern coastal district of Andhra Pradesh, India. *J. Indian bot. Soc.* Doi: 10.5958/2455-7218.2022.00114.0

Source of support: Nil

Conflict of interest: None.

1,800 taxa of Chlorophyceae (http://www.seaweed.ie/). The occurrence, distribution, and diversity of seaweeds differ greatly according to their ecological and hydrological factors. The coastline of India is endowed with 865 taxa of seaweeds reported from the various maritime states. Rhodophyceae is the predominant class encompasses 138 genera of 33 families with 442 (51%) taxa, Chlorophyceae with 212 taxa (25%) belonging to 46 genera of 19 families and Phaeophyceae covers 50 genera of 13 families with 211 taxa (24%) (Rao and Gupta 2015). The marine macroalgal diversity is always enormous on the west coast of India than the east coast of India due to continuous starches of rocky plateaus, hydrological parameters, and other environmental factors. Tamil Nadu coast shows the maximum number of seaweeds (426 taxa), followed by Andaman & Nicobar Islands (244 species); Maharashtra (240 species); Gujarat (198 species); Kerala (147 taxa); Goa (145 taxa); Lakshadweep islands (114 species); Karnataka (108 species); Diu Island (70 species); Odisha (21 species) and West Bengal (14 species).

Though the Indian coastline is endowed with an enormous diversity of seaweeds, most of the states are having the lacuna on the knowledge of seaweeds. Perusal of the literature survey indicate the occurrence of seaweeds on the east coast especially of Andhra Pradesh is limited and poorly studied. Sreeramulu (1952, 1953), Rao and Sreeramulu (1964, 1970), Anonymous (1984), Murthy and Rao (2003), Lakshmi and Rao (2009), Ghosh and Keshri (2010), Rao *et al.*, (2011), Kaliaperumal and Chennubhotla (2015) Sowjanya and Sekhar (2017), Kumar and Palanisamy (2021) were inspected the distribution, diversity, occurrence, and other prospects of seaweeds from Andhra Pradesh in different decades. They merely surveyed Visakhapatnam, the midst coastline of Andhra Pradesh. Information on seaweeds from the northern (Srikakulam and Vizianagaram) and southern coastline (East Godavari, West Godavari, Krishna, Guntur, Prakasam, and Nellore) of Andhra Pradesh are sporadic and poorly known. Hence, this present study is subjected to furnish the data on the occurrence of the marine macroalgae from the northern coastal districts Srikakulam and Vizianagaram of Andhra Pradesh in different seasons.

# **Materials and Methods**

Andhra Pradesh is geographically positioned between 12°41' - 19° 07'N latitude and 69°37' - 84°44'E longitude at the southern part of India. The boundaries of the state are restricted by Tamil Nadu to the south, Karnataka to the southwest and west, Bay of Bengal to the east, Telangana to the northwest and north, and Odisha to the northeast. Among the other coastal states, Andhra Pradesh has got a coastline of around 973 km, running from Pulicat Lake (Nellore district) in the south to Donkuru (Srikakulam District) in the North. Vizianagaram and Srikakulam are the northern coastal districts of Andhra Pradesh with coastal length of 222 km (Vizianagaram - c. 28 km & Srikakulam - c. 194 km). The boundaries of the northern districts limited by Odisha in the northwest, Visakhapatnam in the south, Bay of Bengal in the East. The length of the Nagavali, Vegavathi, Gomukhi, Suvarnamukhi, Champavathi and Gostani are the major rivers flowing in Viziangaram District. Similarly, the rivers Nagavali, Vamsadhara, Mahendratanaya, Champavati, Bahuda, Kumbhikota Gedda, Suvarnamukhi, Vegavati, Gomukhi are stretched in Srikakulam District. The habit of the coast is sandy along with eroded coarse boulders, calcareous rocks, mosaic Rocks, granites and rocky beds support the luxuriant growth of seaweeds (Figures 1-5).

A sum of 22 localities was fixed for the collection of marine macroalgae during low tides in different seasons from March 2017-August 2019. The collected samples were preserved using standard methods of Wet (Liquid preservation) and Dry (Herbarium) adopted by Srinivasan (1969). To avoid the misperception of species, the external and internal morphology of specimens were examined using optical microscopes (Nikon Eclipse 50i; Carl Zeiss. Axio Lab. A1) coupled with a computer attached digital sight DS–Fil camera. Also referred the standard books and monographs such as *Catalogue of the Benthic Marine Algae of the Indian Ocean* (Silva et al., 1996), *Algae of India and neighbouring countries: Chlorophyta* (Krishnamurthy 2000), *A Revised Checklist of Indian Marine Algae* (Oza and Zaidi 2001), *Seaweeds of India: The diversity and distribution of seaweeds in Gujarat* 



Figure 1: Map of study area Northern coastal districts of Andhra Pradesh, India



**Figure 2 (a-h):** Habit and habitats of marine macroalgae from Northern coastal districts. a. Baruva; b. Yeramukkam; c. Bandaruvanipeta; d. Mukkam; e - f. Mixed marine macroalgal diversity at Yerramukkam and Bandaruvanipeta; g. Monospecific diversity of Ulva flexuosa on granite rocks at Lakshimipuram; h. Juvenile stage of red and green algae at Chintapalli.



Figure 3 (a-l): Chlorophyceae. a. Ulva linza; b. U. rigida; c. U. prolifera; d. Chaetomorpha antennina; e. C. linum; f. C. spiralis; g. Acrosiphonia orientalis; h. Bryopsis plumosa; i. Acrocladus herpesticus; j. Valoniopsis pachynema; k. Caulerpa racemosa; l. C. scalpelliformis.



**Figure 4 (a-i):** Phaeophyceae. a. Dictyopteris australis; b. Dictyota dichotoma; c. Padina boergesenii; d. P. boryana; e. P. tetrastromatica; f. P. pavonica; g. Stoechospermum polypodioides; h. Chnoospora minima i. Sargassum cinereum.



Figure 5 (a-l): Rhodophyceae. a. Gelidiella acerosa; b. Gelidium micropterum; c. Gracilaria corticata; d. Grateloupia lithophila; e. Jania rubens; f. Jania spectabile; g. Chondracanthus acicularis; h. Hypnea cervicornis; i. Champia compressa; j. Bostrychia tenella; k. Ceratodictyon repens; l. Acanthophora spicifera.

coast (Jha et al., 2009), A Checklist of Indian Marine Algae: Algae of India Vol. 3 (Rao and Gupta 2015) and Pictorial Guide to Seaweeds of Gulf of Kachchh, Gujarat (Kamboj et al., 2019);

Further, consulted the herbarium specimens housed at Central National Herbarium (CAL), Industrial Section in the Indian Museum (BSIS) Kolkata; Madras Herbarium (MH), Coimbatore; National Facility for Marine Algae Herbarium (NFMAH), Mandapam, Ramanathapuram. The systematic treatment and nomenclature uncertainties were clarified with the aid of online resources such as Macroalgal Herbarium Portal (http://macroalgae.org/), Algae Base (http://www.algaebase.org/) and Marine Biological Association of the UK (http://www.mba.ac.uk/).

# **Results and Discussion**

The current attempt reveals the occurrence of 57 taxa of marine macroalgae belong to 27 genera of 20 families under 15 orders collected from the 22 localities of Srikakulam (18 localities) and Vizianagaram (4 localities) during March 2017 to August 2019. Rhodophyceae is the predominant class represents 26 taxa (46%); Chlorophyceae shows moderate diversity with 20 taxa (35%) and Phaeophyceae shows minimum diversity with11 taxa (19%) reported from the study area. The class Chlorophyceae represents with 6 genera belonging to 5 families under 4 orders. Whilst, Rhodophyceae denotes with 15 genera of 12 families fall under 8 orders. Phaeophyceae comprise 6 genera with 3 families under 3 orders (Table 1). The order Ulvales (Ulva) and Dictyotales (Dictyopteris, Dictyota, Padina, Stoechospermum) signified with 8 taxa in each class. The order Cladophorales includes 6 taxa belong to the family Cladophoraceae with 2 genera (Chaetomorpha and Acrocladus). Order Gracilarials has encountered 5 taxa under 2 genera (Gracilaria & Hydropuntia). Bryopsidaceae (2 taxa) and Caulerpaceae (3 taxa) are characterised with 5 taxa. The rest of the orders Ulotrichales, Ectocarpales, Bangiales contain single taxon. Fucales (2 taxa of Sargassaceae), Gelidiales (3 taxa belong to Gelidiaceae & Gelidiellaceae), Cryptomaniales (2 taxa of Halymeniaceae), Corallinales (3 taxa of Corallinaceae), Rhodymeniales (4 taxa of 2 families) and Ceramiales (2 taxa of Ceramiaceae & Rhodomelaceae) are the orders of moderate values. (Tables 1 and 2).

The occurrence of Ulva linza, U. prolifera, Chaetomorpha linum, Acrocladus herpesticus, Bryopsis plumosa, Dictyopteris australis, Padina boergesenii, P. boryana, P. pavonica, Gelidium micropterum, Gracilaria corticata var. cylindrica, Hypnea cervicornis, H. esperi, J. spectabile, Chondracanthus acicularis and Ceratodictyon intricatum are reported for the first time from the coastline of entire and treated them as the new additions to the marine macroalgal flora of Andhra Pradesh, India. Ulva compressa, U. fasciata, U. flexuosa, U. lactuca, U. prolifera, Acrosiphonia orientalis, Chaetomorpha antennina, Acrocladus herpesticus, Caulerpa taxifolia, Padina tetrastromatica, Stoechospermum polypodioides, Gracilaria corticata, Gracilaria corticata var. cylindrica, Grateloupia filicina, Amphiroa fragilissima, Jania rubens, and Chondracanthus acicularis are the common species find in all the three different seasons. Bryopsis plumosa, Caulerpa racemosa, Dictyopteris australis, Padina pavonica, Sargassum cinereum, Phycocalidia suborbiculata, Gelidium pusillum, Gracilaria foliifera, Hydropuntia edulis, Grateloupia lithophila, Ceratodictyon variabile and Acanthophora spicifera are the mutual species found both in premonsoon and post monsoon seasons. Ulva intestinalis, Chaetomorpha aerea, C. brachygona, C.liniodes, Bryopsis pennata, Gelidium micropterum, Gelidiella acerosa, Hypnea cervicornis, Hypnea esperi, Hypnea musciformis, Champia compressa, Ceratodictyon intricatum and Ceratodictyon repens are the seasonal species found during post monsoon seasons (Table 2).

The Northern coastal districts of Andhra Pradesh gained the maximum atmospheric temperature of 37.6°C in pre-monsoon (February to May); it was quite moderated (28–29.5°C) in monsoon (June to September) and minimum

 Table 1: Numerical data on marine macroalgae from northern

 coastal regions of Andhra Pradesh

Class	Order	Family	Genus	Таха
Chlorophyceae	04	05	06	20
Pheaophyceae	03	03	06	11
Rhodophyceae	08	12	15	26
Total	15	20	27	57

(26.5–27.5°C) in post monsoon (October to January). Likewise, water temperature ranges from 29–32°C in premonsoon; 26.5–28°C in monsoon and 25–27°C in post monsoon. The seawater reaches extreme level of salinity (31.5–32.5 ppt) during pre-monsoon; it was average (30–31.5 ppt) in post monsoon and less (29.5–30.5 ppt) in monsoon. The salinity level of the shoreline varies from 28.0 to 29.5 ppt near to the backwater and estuarine areas due to fresh water entry. The pH level of sea water along the coastline of Andhra Pradesh is around 7.4 to 8.2; but the level of pH in post monsoon is quite greater (7.8–8.2) and it decreasing in monsoon (7.4–7.6). Dissolved oxygen content in the sea waters varied from 7.0 to 7.8 ml/l accordance with the spells of pre-monsoon (7.4–7.8 ml/l), monsoon (7.3–7.6 ml/l) and post monsoon (7.4–7.8 ml/l).

The ecological parameters of the coastline have the great influence towards the distribution and diversity of seaweeds in Andhra Pradesh (Table 3). This study indicates that, spells of post monsoon with the average level of water temperature (26.5–27.5 °C), atmospheric temperature (25–27 °C), salinity (30–31.5 ppt), pH (7.5–7.8), and D.O (7.4–7.8 ml/l) favour the optimized periods in the direction of the enormous diversity of seaweeds (47 taxa). The period of pre-monsoon expressed the moderate diversity of seaweeds (35 taxa) due to the increase of water temperature (29-32°C), atmospheric temperature (28.5-37.6°C), salinity (31.5-32.5 ppt), pH (7.8-8.2). In monsoon, the number of seaweeds was not as much of post monsoon and spotted the occurrence of 26 species account to the reduction in the water temperature (26.5-28°C), atmospheric temperature (28-29.5°C), salinity (29.5-30.5 ppt), pH (7.4-7.6). Further, the present study reveals maximum diversity of seaweeds than the investigation carried out by various researchers such as Sreeramulu (1952, 1953), Rao and Sreeramulu (1964, 1970), Anonymous (1984), Murthy and Rao (2003), Lakshmi and Rao (2009), Ghosh and Keshri (2010), Rao et al. (2011), Kaliaperumal and Chennubhotla (2015) Sowjanya and Sekhar (2017), from Visakhapatnam. The findings of the research from the study area are relatively similar to the coastline of Visakhapatnam based on the physiochemical parameters premeditated by Rao & et al. (2011). It is highly sensitive and susceptible to both natural and anthropogenic threats that lead to the deterioration and extinction of seaweeds. Hence, it is essential to extend the attention on conservation of

Genus	Binomials of the taxa	Pre-monsoon	Monsoon	Post Monsoon
CHLOROPHYCEAE ORDER: ULVALES FAMILY: ULVACEA	E			
	Ulva compressa L.	+	+	+
	Ulva fasciata Delile	+	+	+
	Ulva flexuosa Wulfen	+	+	+
Jlva	Ulva lactuca L.	+	+	+
	Ulva intestinalis L.	-	-	+
	Ulva linza L. *	-	-	+
	Ulva prolifera O.F. Muell. *	+	+	+
	Ulva rigida C. Agardh	+	+	-
Drder: Ulotrici Amily: Ulotrici	HALES HACEAE			
Acrosiphonia	Acrosiphonia orientalis (J. Agardh) P.C. Silva	+	+	+
ORDER: CLADOPH AMILY: CLADOPH	IORALES IORACEAE			
Chaetomorpha	Chaetomorpha aerea (Dillwyn) Kuetz.	-	-	+
	Chaetomorpha antennina (Bory) Kuetz.	+	+	+
	Chaetomorpha brachygona Harv.	-	-	+
	Chaetomorpha linoides Kuetz.	-	-	+
	Chaetomorpha linum (O.F. Muell.) Kuetz. *	-	+	-
crocladus	Acrocladus herpesticus (Montagne) Boedeker *	+	+	+
Drder: Bryopsie Amily: Bryopsie	DALES DACEAE			
Bryopsis	Bryopsis pennata J.V.Lamour.	-	-	+
	Bryopsis plumosa (Huds.) C. Agardh*	+	-	+
DRDER: BRYOPSIC AMILY: CAULERP	ACEAE			
Caulerpa	Caulerpa racemosa (Forssk.) J. Agardh	+	-	+
	Caulerpa scalpelliformis (R. Br. ex Turner) C. Agardh	-	-	+
	Caulerpa taxifolia (M.Vahl) C.Agardh	+	+	+
PHAEOPHYCEAE ORDER: DICTYOTA AMILY: DICTYOTA	ALES			
Dictyopteris	Dictyopteris australis (Sonder) Askenasy*	+	-	+
Dictyota	Dictyota dichotoma (Huds.) J.V.Lamour.	+	+	-
Padina	Padina boergesenii Allender & Kraft*	-	+	+
	Padina boryana Thivy*	-	-	+
	Padina gymnospora (Kuetz.) Sonder*	-	-	+
	Padina pavonica (L.) Thivy*	+	-	+
	Padina tetrastromatica Hauck	+	+	+
toechospermum	Stoechospermum polypodioides (J.V.Lamour.) J.Agardh	+	+	+
ORDER: ECTOCAR AMILY: SCYTOSIP	PALES HONACEAE			
hnoospora	Chnoospora minima (Hering) Papenfuss	-	+	+
ORDER: FUCALES AMILY: SARGASS	ACEAE			
argassum	Sargassum cinereum J.Agardh	+	-	+

Table 2: Seasonal distribution of marine macroalgae from northern coastal district of Andhra Pradesh

	Sargassum polycystum C.Agardh	-	-	+
RHODOPHYCEAE ORDER: BANGIALES FAMILY: BANGIACEA	E			
Phycocalidia	Phycocalidia suborbiculata (Kjellman) Santiañez & M.J.Wynne	+	-	+
ORDER: GELIDIALES FAMILY: GELIDIACEA	E			
Gelidium	Gelidium micropterum Kuetz. *	-	-	+
	Gelidium pusillum (Stackh.) Le Jolis	+	-	-
ORDER: GELIDIALES FAMILY: GELIDIELLAG	CEAE			
Gelidiella	Gelidiella acerosa (Forssk.) Feldmann & Hamel +			
ORDER: GRACILARIA FAMILY: GRACILARIA	LES CEAE			
Gracilaria	Gracilaria corticata (J. Agardh) J. Agardh	+	+	+
	Gracilaria corticata var. cylindrica M.U. Rao*	+	+	+
	Gracilaria foliifera (Forssk.) Boergesen	+	-	-
	Gracilaria textorii (Suringar) Hariot	+	+	+
Hydropuntia	Hydropuntia edulis (S.G.Gmelin) Gurgel & Fredericq	+	-	+
ORDER: CRYPTOMAN FAMILY: HALYMENIA	NIALES CEAE			
Grateloupia	Grateloupia filicina (J.V. Lamour.) C.Agardh	+	+	+
	Grateloupia lithophila Boergesen	+	-	+
ORDER: CORALLINALES FAMILY: CORALLINACEAE				
Amphiroa	Amphiroa fragilissima (L.) J.V. Lamour.	+	+	+
Jania	Jania spectabilis (Harv.) J.H.Kim, Guiry & HG.Choi*	-	+	-
	Jania rubens (L.) J.V. Lamour.	+	+	+
ORDER: GIGARTINAL FAMILY: GIGARTINAC	ES EAE			
Chondracanthus	Chondracanthus acicularis (Roth) Fredericq	+	+	+
FAMILY: CYSTOCLON	IACEAE			
Hypnea	Hypnea cervicornis J.Agardh*	-	-	+
	Hypnea esperi Bory*	-	-	+
	Hypnea musciformis (Wulfen) J.V. Lamour.	-	-	+
	Hypnea valentiae (Turner) Mont.	+	+	-
ORDER: RHODYMEN FAMILY: CHAMPIACE	IALES AE			
Champia	Champia compressa Harv.	-	-	+
FAMILY: LOMENTARI	ACEAE			
Ceratodictyon	Ceratodictyon intricatum (C.Agardh) R.E.Norris *	-	-	+
	Ceratodictyon repens (Kuetz.) R.E.Norris	-	-	+
	Ceratodictyon variabile (J.Agardh) R.E.Norris	+	-	-
ORDER: CERAMIALES	S AE			
Centroceras	Centroceras clavulatum (C. Agardh) Mont.	+	+	-
FAMILY: RHODOMEL	ACEAE			
Acanthophora	Acanthophora spicifera (M.Vahl) Boergesen	+	-	+
Bostrychia	Bostrychia tenella (J.V. Lamour.) J. Agardh	+	+	-
Total		35	26	47

Note: (\*) - Distributional additions to the state

Parameters	Pre-monsoon (February to May)	Monsoon (June to September)	Post monsoon (October to January)
Atmospheric Temperature (°C)	28.5–37.6	28–29.5	26.5–27.5
Water Temperature (°C)	29–32	26.5–28	25–27
Salinity (ppt)	31.5–32.5	29.5–30.5	30–31.5
рН	7.8–8.2	7.4–7.6	7.5–7.8
D.O (ml/L)	7.0–7.3	7.3–7.6	7.4–7.8

Table 3: Physico-chemical parameters of seawater from study area

these unique habitats to protect these natural treasures from habitat destructions. The particulars furnished in present study would be the doorway to scientific community to understand the basic facts of the marine macroalgae from the northern parts of Andhra Pradesh.

#### Acknowledgement

Authors express their sincere gratitude to Dr. A.A. Mao, Director, BSI, Kolkata and Dr. M.U. Sharief, Scientist 'E' & Head of Office, BSI, SRC, Coimbatore, for providing facilities and support. They are thankful to MoEF & CC, New Delhi for financial assistance under AICOPTAX Scheme (Sanctioned order No. 22018/22/2015-RE (Tax) Dt. 05.12.2016).

### References

- Anonymous (1984). A Report on the Survey of Marine Algal Resources of Andhra Pradesh. CSMCRI, Bhavnagar.
- Ghosh S and Keshia JP (2010). The genus *Enteromorpha* link (Ulvales: Chlorophyta) at Visakhapatnam Coast, India. *J. Appl. Biosci.* **36:** 186-190. '

Guiry MD and Guiry GM (2022). *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. https://www.algaebase.org; searched on February 17, 2022.

Jha B, Reddy CRK, Thakur MK and Rao MU (2009).

Seaweeds of India: The Diversity and Distribution of Seaweeds of the Gujarat Coast. Springer, Dordrecht.

- Kaliaperumal N and Chennubhotla V S K 2015 *Studies on marine algae of Andhra Pradesh: A Review. Seaweed Res. & Utiliz.* **37** (2) 1-13.
- Kamboji R D, Das L and Palanisamy M 2019 *Pictorial Guide to Seaweeds of Gulf of kachchh, Gujarat.* Geer Foundation, Gujarat.
- Krishnamurthy V 2000 Algae of India and neighbouring countries I. Chlorophyta. Enfield, NH: Science Publishers, USA.

Kumar A S and Palanisamy M 2021 Investigation on the occurrence of Genus *Ulva* from the shoreline of Andhra Pradesh. *J. Indian Bot. Soc.* **101**(4) 332-340.

- Lakshmi K P and Rao G M N 2009 Some numerical studies on marine algae of Visakhapatnam Coast. J. Algal Biomass Utln. 1 60–85.
- Lightowler H J and Davies G J 2002 Assessment of iodine intake in vegans: weighed dietary record vs duplicate portion technique. *Eur. J. Clin. Nutr.***56** 765–770.
- Murthy S B K and Rao M U 2003 Effects of industrial wastes on the growth and reproductive stages of macroalgae of Visakhapatnam coastline. *Indian J. Mar.Sci.* **32** 57-66.
- Oza R M and S H Zaidi 2001 A Revised Checklist of Indian Marine Algae. CSMCRI, Bhavnagar, India.
- Rao K S, Murty P and Rao G M N 2011 Seasonal studies on marine algae of the Bhimili coast, East coast of India. *Seaweed Res.* & *Utiliz.* **2**(2) 69–82.
- Rao M U and Sreeramulu T, 1964 An ecological study of some intertidal algae of the Visakhapatnam coast. J. Ecol. **52** 595-616.
- Rao M U and Sriramulu T 1970 An annotated list of the marine algae of Visakhapatnam (India). *Bot. J. Linn. Soc.* **63** 23–45.
- Rao P S N and Gupta R K 2015 Algae of India, Vol 3: A checklist of Indian Marine Algae (Excluding Diatoms & Dinoflagellates). Botanical Survey of India, Kolkata.
- Silva P C, Basson P W, Moe R. L 1996 *Catalogue of the Benthic Marine Algae of the Indian Ocean*. University of California press, London.
- Sowjanya V and Sekhar R 2017 Ecology of marine macro algal flora of Visakhapatnam coastal areas, Bay of Bengal, India. *J. Threat. Taxa* **9** 9911–9919.
- Sreeramulu T 1952 On a Porphyra from Waltair Coast. *Sci. Cult.* **18** 285-286.
- Sreeramulu T 1953 A Codium from the Coromandel Coast. *J. Indian Bot. Soc.*, **32** 67-69.
- Srinivasan K S 1969 Phycologia Indica: Icons of Indian Marine Algae, Volume-1. BSI Publication, Calcutta.