https://doi:10.61289/jibs2023.27.09.1136

ORIGINAL ARTICLE



Identification of five species of *Beilschmiedia* using leaf architecture – an anatomical tool

Ulka Chodankar, Meenakshi Vaidya*

Abstract

The genus *Beilschmiedia* belongs to family Lauracaeae. For the present study five species of genus *Beilschmiedia* namely *Beilschmiedia assamica, B. gammieana, B. roxburghiana, Beilschmiedia brandisii* and *B. fagifolia* were collected. All five species of genus *Beilschmiedia* studied all show Pinnate Camptodromous festooned brochidodromous type of venation. The highest vein order is 6°. Based on the secondary veins angle of divergence and angle of origin of tertiary veins, the species can be further separated. There is very little work done on the the leaf architecture studies in this genus and so this present study was undertaken. The species are so intimate by their leaf morphological characters, it is often difficult to determine their nomenclature types. Hence attempt has been made to recognise the taxonomic value of laminar architecture of the species growing in India.

Keywords: Leaf architecture, Beilschmiedia, Pinnate Camptodromous, festooned brochidodromous.

Introduction

The family Lauraceae aromatic erect, trees or shrubs. Hooker (1883) separated Lauraceae generically on the character of 2 and 4 celled anthers into tribe Perseaceae which had shrubs or trees. Third row of stamens, if present, with the anthercells opening outwards by valves. *Beilschmedia* belongs to it. The genus has evergreen trees or shrubs. Leaves are alternate or opposite and flowers are small, bisexual borne in fascicles or panicles or solitary. For the present study the species collected are *Beilschmiedia assamica*, Meissn., *Beilschmiedia brandisii*, HK. f., *Beilschmiedia fagifolia*, Nees., *Beilschmiedia*

S.V. K. M's Mithibai College of Arts, Chauhan Institute of Science and Amrutben Jivanlal College of Commerce and Economics (Autonomous) Affiliated to University of Mumbai, Vile Parle - West Mumbai, India.

*Corresponding Author: Meenakshi Vaidya, S.V. K. M's Mithibai College of Arts, Chauhan Institute of Science and Amrutben Jivanlal College of Commerce and Economics (Autonomous) Affiliated to University of Mumbai, Vile Parle - West Mumbai, India, E-Mail: meenakshi.vaidya@mithibai.ac.in

How to cite this article: Chodankar U, Vaidya M (2023). Identification of five species of beilschmiedia using leaf architecture - an anatomical tool. *J. Indian bot. Soc.*, Doi: 10.61289/jibs2023.27.09.1136

Source of support: Nil

Conflict of interest: None.

gammieana, King. and Beilschmiedia roxburghiana, Nees. Beilschmiedia assamica has opposite leaves which are elliptic ovate or lanceolate obtusely acuminate shining and reticulate on both surfaces. The leaves of *B. brandisii* are alternate, elliptic-oblong acuminate glabrous. While the leaves of *B. fagifolia* are elliptic or elliptic-lanceolate, obtusely acuminate. Those of *B. gammieana* are opposite, elliptic-oblong or ovate obtusely acuminate shining and reticulate on both sides. *B. roxburghiana* has ovate or ovate lanceolate or elliptic oblong, obtusely acuminate, shining and finely reticulate on both surfaces (Hooker, 1883).These species are difficult to identify morphologically and so leaf venation study could be an important tool in the separation of the species.

Materials and methods

The plant material for this work was collected from Shillong-Meghalaya. The duplicates of herbarium were collected from the herbarium section of B.S.I. Eastern Circle and A.R.I., Pune. The plant material which was collected was identified, checked and authenticated with the help of Standard Herbaria from A.R.I. Herbarium, Pune and B.S.I. Eastern Circle, Shillong.

For leaf clearing the method used was as described by Payne, (1969) and Mohan Ram and Nayyar, (1978) in which the mature leaves either fresh, dried or preserved were first cleared by keeping them in 5% sodium hydroxide solution at room temperature for 1-2 days. The decoloured leaves



Figure 1: a-primary vein; b-secondary vein; c- tertiary vein, d-areole, e-margin, f-secondary vein, g-tertiary vein, h-quarternary vein, i-quinternary vein, j- 60 vein

a-primary vein; b-secondary vein; c-tertiary vein, d-areole, e-margin, f-secondary vein, g-tertiary vein, h quarternary vein, i-quinternary vein, j- 6º vein

were washed and transferred to 5% sodium hypochlorite till they were transparent. They were then thoroughly washed to remove acid traces and were stained with aqueous Saffranine by keeping them in it for 10-15 minutes. The leaves were then mounted in glycerine jelly. The terminology used in is in accordance with Hickey (1973, 1979), Hickey and Wolfe (1975), Dilcher (1974) and Melville (1976).

Observations and Results

Out of the five species of genus Beilschmiedia studied all show Pinnate Camptodromous festooned brochidodromous type of venation. The highest vein order is 5° in Beilschmiedia assamica, B. gammieana and B. roxburghiana. Secondary vein angle of divergence is acute moderate and pattern of tertiary veins is random reticulate in B. gammieana and B. roxburghiana. These two species can be further separated on the basis of angle of origin of tertiary veins OO/OR/ OA/RO/RR in B. roxburghiana and OR/AR/RR/OO/RA in B. gammieana. The secondary veins angle of divergence are acute narrow and tertiary veins pattern is orthogonal reticulate with angle of origin of tertiary veins RR/RA/AR/AA/ RO in B. assamica. In B. assamica the pattern is orthogonal reticulate. The higher order venation forming a reticulum in which vein orders are distinct. Quarternary veins (4^o) are thin; its course is orthogonal. Quinternary veins (5°) are thin; its course is orthogonal. The highest vein order of leaf is 5°. The marginal ultimate venation is looped. Areoles are well developed; arrangement is random reticulate and shapes are triangular, quadrangular, pentagonal polygonal. Veinlets are present and are branched three times.

Tracheoids present at the vein endings.

The highest vein order is 6°, tertiary veins pattern is random reticulate and observed in *Beilschmiedia brandisii* and *B. fagifolia*. These two species are separated on the basis of angle of origin of tertiary veins which is AO/AR/RR/RA/ RO/OO in *B. fagifolia* and OO/OR/RR/RO/AO in *B. brandisii*. In *B. brandisii* the pattern is random reticulate. The higher order venation forming a reticulum in which vein orders are distinct. Quarternary veins (4°) are thin; its course is orthogonal. Quinternary veins (5°) are thin; its course is orthogonal. The highest vein order of leaf is 6°. The marginal ultimate venation is looped. Areoles are well developed; arrangement is random reticulate and shapes are triangular, quadrangular, pentagonal, polygonal. Veinlets are present and are branched three times. Tracheoids present at the vein endings.

Key to the species

Pinnate Camptodromous, with festooned brochidodromous venation

Highest vein order 5°

Secondary veins angle of divergence acute narrow Pattern of tertiary veins orthogonal reticulate Angle of origin of tertiary veins RR/RA/AR/AA/RO......B. assamica

Secondary veins angle of divergence acute moderate Pattern of tertiary veins random reticulate Angle of origin of tertiary veins is OR/AR/RR/OO/RAB. roxburghiana

Angle of origin of tertiary veins is OO/OR/OA/RO/RRB. gammieana

Highest vein order 6°

Secondary veins angle of divergence acute narrow

Pattern of tertiary veins random reticulate

Angle of origin of tertiary veins is OO/OR/RR/RO/AOB. brandisii

Secondary veins angle of divergence acute moderate

Pattern of tertiary veins random reticulate

Discussion

Leaf architectural study is found to be useful for taxonomic purpose. Kerner and Oliver (1895) had worked out systems of classification based on foliar venation. Later Foster (1953) published paper dealing with foliar venation. Hall and Melville (1954) proposed veinlet termination number as a technique for testing the purity of fragments of a particular leaf type for pharmacognostical properties. Meyerhoff (1952) and Klucking (1962) worked on venation features of angiosperm leaves. Rao and Bhupal (1973) have studied the sclereids. Rao and Das (1979) presented a resume on the morphological features of tracheoids. The use of cuticular features in taxonomy have been described (Metcalfe and Chalk, 1950; Kasapligil, 1962; Avita and Inamdar, 1980; Ravindran et al, 1991,1993; Christophel *et al.* 1996; Baruah and Nath, 1997).

Vaidya & Agharia (2015) have completed the pharmacognostic studies of *Calophyllum* in which the venation study was an important part.

Chodankar and Vaidya (2021 a, b, c) have already studied the leaf architecture in *Machilus, Phoebe* and *Cinnamomum.* Vaidya (2014, 2015 a & b) has studied the venation and trichomes and stomata (2016a & b) in the species of *Litsaea* which also belongs to family Lauraceae.

Conclusion

The species of *Beilschmiedia* can be separated morphologically based on leaves being pubescent or tomentose, coriaceous and clustered or not clustered. Inflorescence being glabrous, or pubescent and so it is difficult to identify the correct species. Even though all the five species showed pinnate camptodromous, with festooned brochidodromous venation they can be separated with the help of venation pattern, and correctly identified. Thus, anatomy will be a beneficial tool in the identification of the species.

References

- Avita S and JA Inamdar (1982). Stomatal Complex in Lauraceae. Acta Bot. Indica, 9 (1):50-56.
- Baruah A and SC Nath (1997). Foliar epidermal characters in 12 species of *Cinnamomum* Sch. Lauraceae. Phytomorphology, **47 (2):** 1127-134.

- Chodankar Ulka and Meenakshi Vaidya (2021a). Anatomy A Useful Tool In The Identification Of Ten Species Of Genus *Machilus* Of Family Lauraceae in JIBS Vol. **101 (4)** : 349-355
- Chodankar Ulka and Meenakshi Vaidya (2021 b). Patterns Of Leaf Architecture In Six Species Of *Phoebe* From Family Lauraceae In WJPR, **10 (3):** 2021: 1772- 1778
- Chodankar Ulka and Meenakshi Vaidya (2021 c): Study Of Leaf Architecture In Eleven Species Of *Cinnamomum* Blume. Of Family Lauraceae In WJPR Volume **10(2):** 1190- 1199.
- Christophel DC, R Kerrigan and A Rowett (1996). The use of cuticular features in the taxonomy of Lauraceae. Ann. Missouri. Bot. Gard., **83 (3):** 419-432.
- Dilcher D L (1974), The Study of Angiosperm Leaf remains. *The Botanical Rev.*, **40** 1-157.
- Foster AS (1953). Techniques for the study of venation pattern in the leaves of angiosperms. Proc. 7th Int. Bot. Con. Stockholm (1950): 586 -587.
- Hall J P and Melville C (1951). Veinlet termination number a new character for the differentiation of leaves. J. Pharm. Pharmac., **3: 934- 941.**
- Hickey L J (1979). A revised classification of the architecture of dicoty ledonous leaves In: Metcalfe and Chalk. Anatomy of dicotyledons. Clarendon Press, Oxford.
- Hickey LJ (1973). Classification of the architecture of dicotyledonous leaves. Amer. J. Bot., **60:** 17-35.
- Hickey LJ and JA Wolfe (1975). The basis of angiosperm Phylogeny: venation. Ann. Missouri Bot. Gard., **62:** 538-589.
- Hooker J D 1883 Flora of British India Vol-**5:** 116-189. Reeve and Co., London.
- Kasapligil B (1962). An anatomical study of the Secondary tissue in roots and stems of *Umbellularia californica* Nutt. and *Laurus nobilis* L. L. Madrono 16: 205-224.
- Kerner A J and Oliver FC (1895). The Natural History of Plants Vol. I. New York. Holt. Rinehait and Winston.
- Klucking E P (1962). An Oligocene flora from the Western Cascades, with an analysis of leaf structure, Ph. D. Thesis, University of California, Berkeley, Dept. of Geology.
- Melville R (1976). The terminology of leaf architecture of Apocynaceae. Taxon, **25**:549-561.
- Metcalfe C R and Chalk L (1950). Anatomy of Dicotyledons Vol. I. Clarendon Press.
- Meyerhoff AA (1952). A study of leaf venation of Betulaceae with its applications to Paleobotany, Ph. D. Thesis, Stanford University.
- Mohan Ram, HY and Nayyar Vijaylaxmi (1978) Leaf clearing technique with a wide range of applications; Proc. Indian Acad. Sci. (Plant Sci.) B., 87: 125-127.
- Payne, WW (1969). A quick method for clearing leaves. Ward's Bulletin, **8(61):** 4,5.
- Rao T A and Das Silpi (1979). Leaf sclereids-occurrence and distribution in the angiosperms. *Bot. Not.*, **132:** 319-324.
- Rao TA and OP Bhupal (1973). Typology of the sclereids, Proc Indian. Acad. Sci. (Sect.B)., 77: 41-55.
- Ravindran Shylaja, Balakrishnan R, Manilal KS and PM Ravindran (1991). Study in *Cinnamomum* species. Feddes Repert., **102(3/4):** 167-175.
- Ravindran Shylaja, Manilal KS and PM Ravindran (1993). Dermal morphology of 8 species of *Cinnamomum*, J. Plant Anatomy and morphology(Jodhpur), **6(1)**: 42-50.
- Vaidya M (2014). Anatomy- Leaf Venation: A tool for identification of some species of genus *Litsaea* Lamk. of family Lauraceae. *J. Indian Bot. Soc.*, **93 (3 & 4):** 103-108

Vaidya M (2015a). Role of Anatomy in identification of twelve species of genus *Litsaea* Lamk. in Asian Journal of Biochemical and Pharmaceutical Research, 5(1): 93-100.

Vaidya M (2015b). Study of leaf architecture in some species of *Litsaea* Lamk. Of family Lauraceae in Journal of anatomy. Photon 115: 182-184.

Vaidya M (2016 b). Study of Trichomes in Some Species of Litsaea

Lamk. in WJPR, 5 (2):1069-1077.

- Vaidya Meenakshi (2016a). Study of Stomatal Complexes in Somev Species of the Genus *Litsaea* Lamk. in WJPR, **5 (1):** 851-863.
- Vaidya Meenakshi & E R Agharia (2015). Pharmacognostic studies of the leaves of *Calophyllum inophyllum* Linn. *International Journal of Green and Herbal Chemistry*. **4(3):** 215 – 221.