

https://doi: 10.61289/jibs2024.11.08.152

# SHORT COMMUNICATION

# Securidaca L. (Polygalaceae): A New Generic Record To The Flora Of Andhra Pradesh, India

Polimati Jimmy Carter<sup>1</sup>, Gollamandala Ravi<sup>2\*</sup>

#### **Abstract**

The genus *Securidaca* L., is reported here as a new record for the flora of Andhra Pradesh. Detailed description, location map, ecological information and photographs are provided for its easy identification.

Keywords: Alluri Sitharama Raju district, Andhra Pradesh, Eastern Ghats, Flora, Securidaca.

#### Introduction

The Polygalaceae family showcases a broad distribution, with its members spanning various regions across the America and Africa. This family encompasses a diverse array of plant forms, including trees, shrubs, herbs, and woody climbers, comprising approximately 21 genera and around 900 taxa (Chadat, 1980; Banerjee, 1993; Eriksen, 1993; Eriksen & Persson, 2006). In India, the representation of the Polygalaceae includes four genera, namely *Epirixanthes* Blume, *Polygala L., Salomonia Lour.*, and *Securidaca L.* The genus *Securidaca* was first defined by Linnaeus in 1754, using specimens from the America. Its distinctive features include a keel with or without a small fringed crest, eight stamens with fused filaments at the base, and a fruit divided into two compartments, each winged. The genus includes 80

species distributed throughout the tropics, predominantly in Australia, with only one species found in India (Willis, 1973). Securidaca inappendiculata Hassk. was first collected by Bakar in 1933 from the riverbank of Beaufort at mile 18, situated at an altitude of 200 meters above sea level (amsl), and deposited under Securidaca tavoyana Wall. ex A.W. Benn. Subsequently, a fossil leaf of S. inappendiculata was discovered from the Siwalik sediments of Seria Naka at the Indo-Nepal border in the Balrampur district of Uttar Pradesh (Prasad et al., 1997)). In the present study, we document Securidaca inappendiculata, a large woody climber thriving in the moist deciduous forests of Alluri Sitharama Raju district in Andhra Pradesh as a new generic addition to the flora of Andhra Pradesh (Fig. 1).

#### **Material and Methods**

While exploring the flora of Alluri Sitharama Raju district, Andhra Pradesh, five mature individuals of a Securidaca species were observed in Akumamidikota Beat, Pullangi Section, Maredumilli Range, Alluri Division. Few specimens were collected from the existing populations by the first author. The voucher specimens were studied critically and identified as Securidaca inappendiculata. The identification was further confirmed by matching the specimens with electronic images (CAL0000207583, CAL0000207584) are available at Indian Virtual Herbarium of Botanical Survey of India (https://ivh.bsi.gov.in/phanerogams general collection/ en?search\_bar=Securidaca+inappendiculata&selection) and also the review of pertinent literature (Pullaiah & Chennaiah, 1997; Reddy et al., 2008) revealed that this genus had not been previously reported from the Andhra Pradesh. Therefore, this discovery marks a new addition to the flora

<sup>1</sup>Godavari Wildlife Society, Rajahmundry, Andhra Pradesh, India. <sup>2</sup>Bharatiya Engineering Science and Technology Innovation University, Mandrol lake, Gownivaripalli, Gorantla Mandal, Sri Satya Sai, Andhra Pradesh, India.

\*Corresponding Author: Gollamandala Ravi, Bharatiya Engineering Science and Technology Innovation University, Mandrol lake, Gownivaripalli, Gorantla Mandal, Sri Satya Sai, Andhra Pradesh, India, E-Mail: gollamandala.ravi2@gmail.com

**How to cite this article:** Carter, P.J., Ravi, G. (2025). *Securidaca* L. (Polygalaceae): A New Generic Record To The Flora Of Andhra Pradesh, India. *J. Indian bot. Soc.*, 105(1):91-93 Doi: 10.61289/jibs2024.11.08.152

Source of support: Nil
Conflict of interest: None.




Fig. 1: Map showing the location of Securidaca inappendiculata Hassk. in Alluri Sitarama Raju District, Andhra Pradesh

of the Andhra Pradesh in terms of genus representation. A detailed description, location map, notes, and accompanying photographs have been provided to aid in the identification of this species. Voucher specimens have been deposited at the Botanical Survey of India, Deccan Regional Centre (BSID), Hyderabad, Telangana.

#### **Observations and Results**

## Taxonomic Treatment

Securidaca inappendiculata Hassk., Pl. Jav. Rar.:295. 1848. Securidaca bracteata A.W.Benn. in J.D.Hooker, Fl. Brit. India 1: 208. 1872. Securidaca tavoyana Wall. ex A.W.Benn. in J.D.Hooker, Fl. Brit. India 1: 208. 1872.

# Description

Robust woody climbers, to 6 m high, with dark brown stem, young branches pink, fibres strong, silky; branchlets slightly puberulous. Leaves elliptic, obovate, oblong or oblanceolate; cuneate at base, entire along margin, cuspidate or abruptly acuminate at apex,  $6.8-14.3\times4-7.5$  cm long, coriaceous, glabrous and deep green above, puberulous and pale beneath; midvein pinkish, raised abaxially, impressed adaxially, lateral nerves 10-12 pairs, oblique; petioles 5-7 mm long. Panicles terminal or axillary, 13-15 cm. Flowers 10-20 cm long, sub corymbose panicles; pedicels filiform, 2-5 mm long, lower pedicels longer than upper, up to 15 mm

long; bracts ovate, caducous, minute, hairy, outer sepals 3, ovate-elliptic to rounded, slightly oblique at base, ca  $3 \times 2$  mm, ciliate; inner wing sepals 2, elliptic, patent, alternate at base, rounded at apex, concave,  $5-6 \times 3-5$  mm. Petals 3; middle petal 5-6 mm long, keeled, crested; lateral petals 2, adnate at base to keel, truncate. Stamens 8, 4–5 mm long. Ovary sessile, sub globose, 2 mm long; style ca. 5 mm long, geniculately bent at top; stigma bilobed. Samara globose;  $6-10 \times 1.5-2.5$  cm long, strongly corrugate; wings oblique, entire, oblong to obovate, straight on one side, rounded or obtuse at apex, rugose, closely strongly arcuate-nerved; seeds suborbicular, compressed, ca 7 mm long. (Fig. 2).

## Habitat and Ecology

During our field visit, we observed five matured individuals growing along the banks of a small stream adjacent to the roadside, found in association with *Macaranga peltata* (Roxb.) Müll.Arg., *Sida acuta* Burm.f., *Lantana camara* L., *Sida rhombifolia* subsp. *rhombifolia*, *Dioscorea* sp., and *Ipomoea* sp.

## Phenology

August - October.

# **Global Distribution**

India, Nepal, Bhutan, Bangladesh, Myanmar, Vietnam, China, Malaysia, Thailand, Indonesia, Cambodia, Laos, Philippines.



**Fig. 2:** Securidaca inappendiculata Hassk., A – Habitat; B, C, D, & E- Leaf both sides and closeup of leaf lower surface and stem portion; F – Flowering twig; G & H – Inflorescence and flower closeup; I – Anthers; J – Ovary; K– Fruiting twig.

#### **Distribution** in India

West Bengal, Uttar Pradesh, Assam, Tripura, Meghalaya and Orissa. Present report from Andhra Pradesh (Alluri Sitharama Raju District).

## Specimens Examined

INDIA: Eastern Ghats, Andhra Pradesh, Alluri Sitharama Raju district, Alluri Division, Maredumilli Range, Pullangi Section, Akumamidikota Beat, Latitude: 17.716444° N; Longitude: 81.769361° E at 534 mt. Jimmy Carter. P 6693; 02.09.2023 (BSID).

Additional specimens examined: Securidaca cumingii Hassk. ex Miq. Cuming, H. Philippines. MNHN P P02547029. Securidaca inappendiculata Hassk. Zollinger H Indonesia. Botany L 0016592 and Indian virtual Herbarium (IVH) BSI, CAL0000207583, CAL0000207584).

# Acknowledgements

The authors express their gratitude to the Andhra Pradesh Forest Department for necessary permission and logistic support in conducting field studies, and thanks are also extended to Mr. P. Bharath Simha Yadav, Research scholar (The Madura College, Madurai, Tamil Nadu) and Mr. Ashutosh Sharma, Research scholar (The University of Trans-Disciplinary Health Sciences and Technology (TDU), Yelahanka, Bengaluru, Karnataka), for his inputs during manuscript preparation. The second author extends gratitude to Dr D. Veeranjaneyulu, EIACP Division, EPTRI, Telangana for his unwavering support.

#### References

Banerjee RN (1993). Polygalaceae. In: *Flora of India* (Sharma, B.D. & N.P. Balakrishnan, eds.) **2**: 450 – 493. Botanical Survey of India, Calcutta.

Chadat R (1980). *Monographia Polygalacearum*. Mem. Soc. Phys. Hist. Nat. Geneva (Suppl.) **7**:1 –143.

Eriksen B (1993). Taxonomical studies in the Polygalaceae and Valerianaceae. (Ph.D. Thesis) University of Goteborg, Goteborg, Sweden.

Eriksen B and Persson C (2006). Polygalaceae. In: Families and genera of vascular plants (Kubitzki, K., ed.) *Springer*, **9**: 345 – 363.

Prasad M, Antal JS & Tiwari V D (1997). Investigation on plant fossils from Seria Naka in the Himalayan foot hills of Uttar Pradesh, India. *Palaeobotanist*, **46** (3), 13 – 30.

Pullaiah T & Chennaiah E (1997). *Flora of Andhra Pradesh* (India) Vol. I. Scientific Publishers, Jodhpur, India.

Reddy CS, Reddy KN & Raju VS (2008). Supplement to *Flora of Andhra Pradesh*, India. Bishen Singh Mahendra Pal Singh, Dehradun.

Willis JC (1973). A dictionary of the flowering plants and ferns (8<sup>th</sup> edition). Cambridge University Press, Cambridge.