



Plant profile, phytochemistry, pharmacology and genetic diversity of *Saraca asoca*: a vulnerable medicinal tree of India

Shukla Saluja*, Yogendra K. Gautam

Abstract

Saraca asoca, commonly known as 'Ashoka', is an important ancient and indigenous medicinal plant, found throughout India. The word 'Ashoka' means 'remover of sorrow' in Sanskrit, with reference to its bark keeping women healthy and youthful. It is also an important sacred and religious tree. Although the bark is most useful, every part of the plant including flower, leaves, root and seeds has medicinal properties and has been used in Ayurvedic medicinal preparations for centuries. The bark contains glycoside, tannins, flavanoids, saponins, esters and primary alcohols which contribute to its medicinal properties. Overexploitation, unethical harvesting and deforestation of natural habitats have reduced its population in the wild, and consequently the species is being listed in the 'globally vulnerable' category by the IUCN. Increased market demand has resulted in adulteration with substitutes. This review is an attempt to consolidate the current knowledge of scientific information on the therapeutic properties, phytochemistry, pharmacology, relationship among related taxa and genetic diversity of the species.

Keywords: Alternative medicine, Ashoka, genetic variation, medicinal plant, pharmacology, phytoconstituents.

Introduction

Medicinal plants have been a part of our lives and utilized in traditional medicines from time immemorial. A large number of herbal medicines and plant-based drug formulations of immense therapeutic value are described in Indian folk medicines. Even today the plant-based drugs are popular over modern synthetic drugs mainly due to their negligible side-effects, low-cost and easy availability. *Ayurveda*, a traditional medicinal practice based on human physiology, employs a large number of medicinal plants and herbs, either exclusively or in combination for overall health. According to WHO (2005), in developing countries like India, almost 70% of the population is dependent on traditional herbal medicines for their primary health care needs.

Department of Botany, Sri Venkateswara College, University of Delhi, New Delhi, India

*Corresponding Author: Shukla Saluja, Department of Botany, Sri Venkateswara College, University of Delhi, New Delhi, India, E-Mail: shuklasaluja@svc.ac.in

How to cite this article: Saluja, S., Gautam, Y.K. 2023. Plant profile, phytochemistry, pharmacology and genetic diversity of Saraca asoca: a vulnerable medicinal tree of India. J. Indian bot. Soc., 103(2):109-118. Doi: 10.5958/2455-7218.2023.00018.9

Source of support: Nil

Conflict of interest: None.

Saraca asoca (Roxb.) W. J. deWilde belonging to family Fabaceae, is an indigenous, ancient, venerated and an important medicinal tree of India. It is very popular and has been used in many traditional medical therapies for its extra-ordinary properties, with many health benefits and cures especially for women related disorders (Sharma et al., 2000). The Ashoka tree finds mention in the Charka Samhita as anodynes in gynecological disorders (Mohan et al., 2016). The tree is also known as 'Sita Ashoka', which literally means 'without sorrow' in Sanskrit language owing to its manifold health benefits especially on keeping women healthy and youthful. It is an important sacred tree of Hindus and Buddhists. The Ashoka tree is referred in the legendary Ramayana in connection with the 'Ashoka Vatika', where Hanuman first met Sita, and Lord Buddha was said to be born under this tree species (Pradhan et al., 2009). It is a popular tree with many vernacular names - Ashoka, Sita-ashoka, Aganpriya (Hindi), Kankeli, Gandhpushp (Sanskrit), Oshok (Bengali), Jashundi (Marathi), Asupala, Ashopalav (Gujarati), Asokada, Kankelimara (Kannada) and Asokam (Tamil and Malayalam) (Yadav et al., 2019).

The genus *Saraca* comprises about 20 species (Santapau &Henry 1998). Of these, *S. asoca* (Roxb.) W. J. deWilde, *S. declinata* Miq. (Red Saraca), *S. indica* L. and *S. thaipingensis* Prain (Yellow Saraca) have been reported from India (Begum *et al.*, 2014). Whereas *S. asoca* is found in the wild besides

in gardens and avenues, the other three species are found only in botanical gardens (Begum *et al.*, 2014). Though they have immense medicinal value, these are also grown for ornamental purpose (Sil *et al.*, 2018).

The most popularly used part of the Saraca asoca is the bark, even though all the other parts such as leaves, roots, flowers, and seeds exhibit medicinal properties with extensive use in anti-microbial, anti-cancer, anti-inflammatory, antiarthritic and many more medications (Yadavet al., 2019). The bark is useful in treating excessive menstrual bleeding, irregular mensuration, pain, dysmenorrhea, congestion, uterine spasms, and leucorrhoea and as a uterine tonic with a stimulatory effect on the ovarian tissue endometrium (Baranwal 2014). Ashokarishta, Ashokaghritha, Ashoka kwath, Asoka arishtam and Madhukadya are some of the well-known indigenous preparations commonly available, which uses S. asoca as the primary raw constituent (Smitha & Thondaiman 2016). The plant parts contain many secondary metabolites including tannins, flavonoids, alkaloids, steroids, saponins, glucosides, terpenoids, triterpenoids, phenolics, catechol and other phytochemicals which contribute to its medicinal properties (Ahmed et al., 2016).

Ashoka flowers are used to decorate Buddhist monasteries and Hindu temples (Boroker & Pansare 2017). The two important rituals associated with the tree are 'Ashoka tri-ratri' and 'Asokasathi', which represents its cultural significance (Pradhan *et al.*, 2009). As a ritual in traditional culture, the flower buds are eaten by married Hindu women on 'Ashok Shashti day' to seek protection for children against negativity, grief and sorrowand to invoke the deities (Pradhan *et al.*, 2009). Many tribal womenfolk from Chhattisgarh drink a decoction of the bark in cow's milk and sugar for three days after every three months to prevent gynecological problems and diabetes (Boroker & Pansare 2017).

Unethical harvesting, indiscriminate use of the plant parts, increasing demand and unscientific management practice have resulted in severe shortage of the genuine plant product which in turn has resulted in an increased cost and rampant substitution and adulteration (Smitha & Thondaiman 2016). Over-exploitation due to its high medicinal value, increasing market demand, habitat loss due to deforestation, habitat fragmentation, narrow ecological niche, illegal encroachments of its natural habitats and poor seed viability has reduced its natural population in the wild (Devan & Warrier 2021). An assessment based on category B i.e., based on extent of occupancy and area of occupancy, no sub populations have been reported in Sri Lanka in the forest surveys conducted during 1991 and 1996 for National Conservation Reviews indicating Saraca to be extremely rare or possibly extinct (www.iucnredlist. org/species/34623/9879360#population). The species is presently listed as 'a globally vulnerable species' in the

threatened species list of IUCN (https://www.iucnredlist. org/species/34623/9879360). This review is an attempt to compile different research works on the phytochemical constituents and pharmacological activities of the various plant parts, relationship among related taxa and molecular marker studies for its molecular characterization which may help in understanding the genetic diversity within the population that may be beneficial for future conservation and improvement strategies of the tree. A brief account of distribution and botanical description /morphology is also included. The study may provide a comprehensive view for the management of the species.

Morphology

Saraca asoca is a small to medium-size evergreen tree with a dense and spreading crown (Figure 1). The tree can grow up to 7 - 10 m in height. The bark is grey or dark brown covered with a warty surface. It is uneven and rough to touch, due to the presence of projecting lenticels and circular lenticelsand is channeled or sometimes cracked (Boroker & Pansare 2017). The leaves are alternate, paripinnate, 15-20 cm long, copper red when young and green on maturity. The leaves have 4 - 6 pair of leaflets which are oblong-lanceolate, obtuse or acute, glabrous with rounded base; petioles are short (Boroker & Pansare 2017). Flowers are fragrant, and borne in dense axillary corymbs. They are bisexual, aromatic, staminate and astringent in taste. Young flowers are yellow which gradually turn to orange and scarlet red on maturity (Mishra & Vijaykumar 2015). Pods are linear, oblong, compressed, dehiscent, tapering to both ends and are green and leathery which turns to brownish purple or black on maturity. Each pod contains 4 - 8 seeds which are ellipsoid, oblong, compacted and covered with a seed-coat. The seeds turn black on maturity and are recalcitrant type (Chauhan 2019).



Figure 1: Saraca asoca. A. Tree in flowering stage. B. A close view of the twig. C. An enlarge view of inflorescence. D. Pods. E. Pod with seeds. (Photo credit: Prof. S. R. Yadav).

Distribution

Saraca asoca is native to the Indian sub-continent, though also found in Indo-Malaysian region and Sri Lanka. In India, it is mainly found in peninsular region extending into the Western Ghats, Eastern Ghats and the sub-Himalayan tracks, growing at an elevation of approximately 750 m (Patwardhan *et al.*, 2014). Its original distribution was in the Deccan Plateau and distributed in Central and Eastern Himalaya, Burma, East Bengal, Western Peninsula, and Sri Lanka (Preeti *et al.*, 2012). It is also found in semi-evergreen and in moist deciduous forests along the river streams (Patwardhan *et al.*, 2016). The most suitable growth condition is red laterite alluvial soil with an annual rainfall of 2000 - 4000 mm (Yadav *et al.*, 2019).

Phylogeny

Phenetic study based on morphological characters and ternary plot analysis has shown S. declinata to be closer to S. thaipingensisas compared to S. indica (Sil et al., 2018). Traits such as ovary texture, the pattern of distribution of trichomes on anther and hooded anther, exine surface, bract, stigma morphology and shape and size of the seed have been suggested to be crucial morphological markers for delimiting the three species of Saraca. A molecular analysis of the intra and inter specific diversity of six species of Saraca using matK sequences showed Saraca as a diverse genus with different species. The phylogenetic interrelationships have indicated S. declinata to be the most evolved and S. dives with the foremost evolution. Saraca indica and S. asoca showed firm molecular resemblance yet their molecular diversity has placed them as separate species while S. declinata and its synonym S. palembanica exhibit distinct molecular identities (Sil et al., 2021).

Phytochemistry

The biological activities of the medicinal plants are generally due to the presence of secondary metabolites. Ashoka bark, leaves and flowers which show therapeutic importance have a variety of phytochemical constituents including flavonoids, alkaloids, triterpenoids, phenolics, tannins, steroids, glycosides, quinones and volatile organic compounds (Singh *et al.*, 2015, Ahmad *et al.*, 2016) (Table 1). Minerals such as sodium, potassium, calcium, aluminum, iron, magnesium, strontium and phosphate are present in the bark (Saha*et al.*, 2012). The wax from the bark also contains n-alkanes, primary alcohol, esters and n-octacasanol (Boroker & Pansare 2017).The plant seeds and fruits contain a lectin named Saracin (Ghosh *et al.*, 1999). The chemical structures of some of its phytochemicals drawn on software ChemSketch 2015are shown in Figure 2.

Adulterants

Overexploitation, destructive extraction and non-availability of Ashoka trees due to increased market demand have caused increased adulteration or substitution of the drug. The commonly used adulterants are the barks of Polyalthia longifolia, Shorea robusta, Trema orientalis, Bauhinia variegata, Afananexis polystakis and Caesalpinia pulcherrima(Khatoon et al., 2009, Urumarudappa et al., 2016). Adulteration and/ or substitution with Polyalthia longifolia is common as both have the same vernacular name 'Ashoka' resulting in misidentification and confusion (Khatoon et al., 2009). A comparison of transverse and longitudinal sections of stem bark of Saraca asoca and Polyalthia longifolia which has smaller epidermal cells along with specialized cells in the cortical region can help in identification of the two barks by microscopy (Khatoon et al., 2009). Some of the identifying features of S. asoca, S. declinata, and Polyalthia longifolia are listed in Table 2 (Devan & Warrier 2021) (Table 2). Adulteration or substitution can adversely affect the health and safety of the consumer and reduce the therapeutic efficacy of the traditional medicine. The NMR spectroscopic analysis to validate market samples with S. asoca, has shown high level of adulteration in the market samples (Urumarudappa et al., 2016). Kumar (2016) used RAPD derived SCAR markers to identify and differentiate S. asoca from P. longifolia. Hedge et al., (2017) isolated and used Start Codon Targeted (SCoT) and Inter Simple Sequence Repeat (ISSR) markers and phytochemical profiling by RP-HPLC analysis to identify bark samples for validation.

Biological and pharmacology activities of Saraca asoca

Saraca asoca has multiple pharmacological activities attributed to its rich phytochemicals (Table 3). Some of the activities are discussed below.

Anti-mennorhagic and uterine tonic

The dried bark, root and flowers are especially useful for the treatment of gynecological problems, mainly associated with absence of periods (amenorrhea), irregular periods, painful periods (dysmenorrhea), excessive menstrual bleeding (menorrhagia), premenstrual syndrome, leucorrhoea and endometriosis. It has an estrogenic stimulatory effect to the endometrium and ovarian tissues acting as a uterine tonic (Baranwal 2014). Swar *et al.*, 2017 treated ovariectomized female albino Wistar rats with ethanolic flower extract and observed that the flowers has estrogenic potency which may be due to phytoestrogens such as β -sitosterol, kaempferol, luteolin and quercetin present in the flower.

Effect on menstrual cycle

An investigation to study the restoring capacity of the bark extract on reproductive cycle, Progesterone (P4), Estrogen (E2) levels over disruptions caused due to the synthetic contraceptive Levonorgestrel (LNG) in female albino rats, showed that the methanol bark extract significantly reduced the hormone levels to normal (Hazarika *et al.*, 2021).

Table 1: Various phytoconstituents from different plant organs.

Class	Compound	Plant organ	Reference
Flavonol	Catechin	Bark, seed, pod	Mukhopadhyay et. al., 2017
	Epicatechin		Pradhan <i>et. al.</i> , 2009
	Leucocyanidin		Ahmad <i>et. al.</i> , 2016
	Leucopelargonidin	Bark, leaves	Ahmad <i>et. al.</i> , 2016,
	Epigallocatechin	Bark	Ahmad <i>et. al.,</i> 2016
	Gallocatechin		Ahmad <i>et. al.</i> , 2016
	epiafzelechin-(4 β $ ightarrow$ 8)- epicatechin		Pradhan <i>et. al.</i> , 2009
	procyanidin B2		Pradhan <i>et. al.,</i> 2009
	3',5 -dimethoxy epicatechin		Ahmad <i>et. al.</i> , 2016
	Deoxyprocyanidin B		Singh <i>et. al.,</i> 2015
	Haematoxylin		Singh <i>et. al.</i> , 2015
Flavonol glycosides	Quercetin-3-O-P-D-glucoside	Flowers	Pradhan <i>et. al.</i> , 2009
	Cyanidine-3,5-diglucoside		Pradhan <i>et. al.,</i> 2009
	Pelargonidin-3,5-diglucoside		Pradhan <i>et. al.,</i> 2009
	3'-deoxycatechin-3-O-α-L-rhamnopyranoside	Bark	Ahmad <i>et. al.,</i> 2016
	3'-deoxyepicatechin-3-O-β-D-glucopyranoside		Ahmad <i>et. al.</i> , 2016
Lignin glycosides	Lyoniside	Bark, leaves, flowers	Ahmad <i>et. al.,</i> 2016
	5-methoxy-9-β-xylopyranosyl isolariciresinol	Bark	Singh <i>et. al.,</i> 2015
	Icariside E		Sadhu <i>et. al.,</i> 2007
	Nudiposide		Sadhu <i>et. al.</i> , 2007
	Schizandriside		Sadhu <i>et. al.</i> , 2007
Phenolic acid	Gallic acid	Bark, leaves, flowers	Saha <i>et. al.,</i> 2013
	Quercetin	Bark, flowers	Saha <i>et. al.</i> , 2013, Swar <i>et. al.</i> , 2017
	Ellagic acid	Bark	Saha <i>et. al.,</i> 2013
	Acetyl salicylic acid	Seed	Gupta <i>et. al.</i> , 2014
Sterol	β-sitosterol	Bark, leaves, flower	Swar et. al., 2017
Steroidal glycosides	β-sitosterol glucoside	Bark, leaves	Sadhu <i>et. al.</i> , 2007
Fatty acid	Oleic acid	Pod, seed	Singh <i>et.al.</i> ,2015, Yadav <i>et.a</i> l., 2019
	Stearic acid		Singh <i>et.al.</i> ,2015, Yadav <i>et.al.</i> , 2019
	Linoleic acid		Singh <i>et.al.</i> ,2015, Yadav <i>et.al.</i> , 2019

Anti-keratinizing

Shahid *et al.*, (2015) investigated the effect of methanolic barkextract in artificially estrogen induced proliferation of endometrial thickening and keratinized metaplasmia in rat uterus. The methanol extract of bark reduced the endometrial thickening to 2 - 4 layers of stratified epithelium and significantly lowered serum estrogen levels. It also reduced the formalin-and carrageenan-induced paw edema, lipopolysaccharides induced COX-2 enzyme in rat uterus, thus indicating it's anti-proliferative and anti-keratinizing

potential manifested through an anti-estrogenic and antiinflammatory activity.

Anti-oxidative and Anti-cancer

Several reports have elaborated the presence of different anti-oxidant compounds in thebark, leaves, and flowers of *S. asoca* which are effective against a number of disorders including cancer (Ahmad & Ghosh 2022). Methanol bark extract of the trees has been shown to reduce growth in HeLa cells (human cervical cancer cell lines) showing important

Trait	S. asoca	S. declinata	Polyalthia longifolia
Stem bark	channeled	curved	curved
Distribution of fibers in phloem region	group of 3-24	solitary or in groups of 3-6	present as broad concentric band which alternates with parenchymatous bands
Phelloderm and stone cells	phelloderm of 2-3 continuous bands of stone cells	phelloderm as outer and inner regions with groups of 6-7 stone cells	stone cells present throughout phelloderm
Medullary rays	uni-biseriate	uniseriate - triseriate	multi-seriate
Mucilage canals and oil cells	absent	absent	present

Table 2: Contradistinctive traits for identification between two species of Saraca and Polyalthia longifolia (Adapted from Devan & Warrier 2021)



Figure 2: Chemical structures of some of the phytoconstituents of *Saraca asoca* (Adapted from Ahmad *et. al.* 2016, Swar *et. al.* 2017).

cytotoxic and anti-tumor activity (Asokan & Thangavel 2014). The MTT and SRB assays showed significant inhibition of HT-29 colon cancer lines showing in vitro cytotoxicity of the bark extract (Jeenathunisa & Rajan 2021). The presence of natural flavonoids (catechin) in methanolic bark extract has shown promising results against acute myeloid leukemia (Mukhopadhyay et al., 2017) in benzene induced swiss albino mice models. The mice treated with catechin showed reduced DNA fragmentation, up-regulation of p53, p21 and bc12, CYP2EI, caspase 11 and myeloperoxidase. Catechin showed modulatory influence on cell cycle deregulation and hematological abnormalities which are associated with benzene induced leukemia. Thus, indicating that catechin is similar to doxorubicin (chemotherapeutic drug) to give stability and protection towards cancer (Mukhopadhyay et al., 2017). Baneriee et al., (2022) synthesized phyto-fabricated silver nanoparticles from an aqueous bark extract of S. asoca. They checked the effectiveness of the nanoparticles on acute myeloid leukemia (AML) and compared to doxorubicin, a chemotherapeutic drug involving in vivo studies on AML mice along with in vitro studies on lymphocytes and leukemia cells to check the cytotoxic and genotoxic effects. The nanoparticles were selectively cytotoxic towards the leukemia lymphocytes cells through oxidative damage, suggesting potential avenues of phyto-fabricated nanoparticles.

Anti-microbial (anti-bacterial and anti-fungal)

Numerous investigations have demonstrated the antibacterial activity in aqueous, alcoholic (methanol and ethanol) and acetone extracts of the bark, leaves and dried flower buds against a number of bacteria such as Escherichia coli, Klebsiella pnemone, Klebsiella aerogenes, Salmonella typhosa, Salmonella typhimurium, Salmonella typhii, Salmonella enteritis, Bacillus subtilis, B. cereus, Pseudomonas vulgaris, Pseudomonas aeruginosa, Vibrio chlorae, Proteus vulgaris, Staphylococcus aureus (Shirolkar et al., 2013, Mohan et al., 2016). The anti-fungal activity of aqueous and methanol extracts of various parts of the plant have been reported against a number of fungi like Alternaria alternate, Alternaria cajani, Colletotrichum gloeosporioides, Helminthosporium sp, Curvularia lunata, Bipolaris sp., Fusarium sp., Candida albicans, Aspergillus flavus, Aspergillus niger and Aspergillus fumigatus(Shirolkar et al., 2013, Chakraborty et al., 2014). Due to its antibacterial efficacy this tree may be a beneficial resource in the future for a single molecular antibiotic against increasing antibiotic resistance of a number of human pathogenic bacteria.

Anti-depressant

Methanolic bark extract was used to evaluate the antidepressant activity in rats by inducing Chronic Unpredictable Mild Stress (CUMS) and estimating the associated antioxidant enzymes such as catalyase, superoxide dismutase activities, glutathione malondialdehyde levels in the brain. The extract showed decrease in the stress parameters along with an increase in the associated antioxidant enzyme levels. This improved the behavioral and neurochemical response, modulating neurotransmitter activity associated with depression, thus showed potential anti-depressant characteristics (Gill *et al.*, 2018).

Anti-epileptic

Ethanol extracts of *S. asoca* flower were used to investigate anti-epileptic activity in swiss albino mice by using Maximum electroshock (MES) and pentylenetetraxole (PTZ) induced seizure models. Decrease in tonic hind limb extension (THLE) phase and increase in the latency period of seizures were measured against the induced convulsions. The ethanol
 Table 3: Biological activities in different plant organs.

Plant parts	Chemical class	Biological activity	Reference
Bark	Flavonoids (catechin)	anti-cancer	Mukhopadhyay et. al., 2017
		cardioprotective	Swami <i>et. al.,</i> 2013
	polyphenols	antioxidant	Singh <i>et. al.,</i> 2015
		anti-depressant regulation of menstrual cycle anti-keratinizing	Gill <i>et. al.,</i> 2018 Hazarika <i>et. al.,</i> 2021
		-	Shahid <i>et. al.,</i> 2015
_eaf	tannins, triterpenoids, saponins, flavonoids and glycosides	analgesic activity	Verma <i>et. al.,</i> 2010
	alkaloids, tannins, flavonoids and saponnins	anti-helmintic	Sharma <i>et. al.,</i> 2011
		anti-ulcer	Verma <i>et. al.,</i> 2020
Flower	phytoesterols	estrogenic stimulatory	Swar <i>et. al.,</i> 2017
	polyphenols, phytoesterols, flavonoids, and saponins	anti-epileptic anti-hyperglycemic and anti- hyperlipidemic	Srinivasa 2018 Mishra and Vijaykumar 2014
	flavonoids	anti-diabetic	Thilagam <i>et. al.,</i> 2021
Seed	phenolic compounds, flavonoids and terpenoids	anti-arthritic and anti- inflammatory	Gupta <i>et. al.,</i> 2014
	flavonoids, tannins and saponins.	anti-pyretic	Sasmal <i>et. al.,</i> 2012

extract of the flowers significantly decreased the THLE and seizure latency in MES and PTZ models, thereby exhibiting the activity (Srinivasa *et al.*, 2018).

Anti-hyperglycemic and hypolipidemic

The methanol, petroleum ether and chloroform extract of S. asoca leaves have exhibited a significant potential for anti-hyperglycemic and antioxidant ability in Streptozotocin induced diabetic rat model, as theoral administration of the extracts significantly reduced the blood glucose levels; the methanol extract was most potential for its antihyperglycemic activity (Kumar et al., 2012). In another study, the ethanolic extract of Ashoka flowers showed remarkable capability for its anti-hyperglycemic and antioxidant activities in Streptozotocin-nicotinamide induced diabetic rats (Mishra & Vijaykumar 2014). The flower extract is rich in phenolic compounds phytoesterol, flavonoids and saponins which lowers the blood glucose levels and those of the antioxidant enzymes including catalyst glutathione peroxidase, superoxide dismutase, glutathione reductase to near normal. The flavonoids in the flower extract, known for their antioxidant properties scavenged the free radicals produced during diabetes (Mishra & Vijaykumar 2014). Thilagamet al., (2021) reported ethanolic extracts of S. asoca flowers containing polyphenols, flavonoids phytoesterols and saponins significantly increased serum insulin levels along with a decrease in blood glucose levels, alkaline phosphatase, HbA1C, serum triglyceride and total cholesterol levels in alloxan-induced diabetic rats suggesting anti-diabetic along with anti-hyperlipidemic activity.

Anti-inflammatory and anti-arthritic

Treatment with *S. asoca* methanol extract to adjuvantinduced arthritis in female wistar rats decreased the paw thickness and raised the body weight, along with significant lowering of the plasma and liver lysosomal enzyme levels, urinary collagen, pro-inflammatory cytokines levels, protein bound carbohydrates, and restored the histological architecture of the joints, thus decreasing the rheumatoid arthritis in rats (Saravanan *et al.*, 2011).

In another study with Freund's adjuvant-induced arthritis in wistar albino rat model, the seed acetone extract with high concentration of phenolic compounds, flavonoids and terpenoids were shown to exhibit anti-arthritic and anti-inflammatory potential by significantly reducing glucosamine and hydroxy proline concentration in the urine, ankle joint inflammation and paw edema along with normal radiological images of the joint (Gupta *et al.*, 2014).

Cardioprotective

Cardioprotective activity was demonstrated by Swami *et al.*, (2013) using cyclophosphamide induced cardiotoxicity in rats. Cyclophosphamide caused changes in ECG, increased biomarker enzymes such as creatine kinase, lactate dehydrogenase, alkaline phosphatase, and increased cholesterol and triglyceride levels. Treatment with ethanol bark extract revoked the changes caused by cyclophosphamide by restoring ECG, cardiac biomarkers, oxidative enzymes, lipid profile and lowered the severity of myocardium cellular damage.

Analgesic

Saraca asoca leaf extracts in water, methanol, chloroform and petroleum ether were examined for phytoconstituents and analgesic activity. The analgesic activity was carried out in albino rats employing tail immersion method and formalin induced methods. Formalin test is an effective analytic method to correlate with clinical pain. The methanol leaf extracts showed highest analgesic activity, relieved pain in the animals injected with formalin in a dose dependent manner by acting on both peripheral and central nervous system. The phytoconstituents in the leaf extract such as tannins, triterpenoids, saponins, flavonoids and glycosides may be involved in the analgesic activity (Verma *et al.*, 2010).

Anti-ulcer

The ethanol extracts of *S. asoca* leaves produced remarkable protection against ulcers which were induced by ethanol, pylorus ligation and indomethacin in albino rats, by reducing ulcer index, increasing gastric pH and reducing total acidity gastric volume showing anti-ulcer activity (Verma *et al.*, 2020).

Anti-pyretic

Anti-pyretic action of acetone seed extract of *S. asoca* was demonstrated in the rat models with artificially induced pyrexia using Brewer's yeast (Sasmal *et al.*, 2012). The anti-pyretic activity could be attributed to the phyto-constituents such as flavonoids, tannins and saponins.

Anti-helmintic

Parasitic worms are common infections in man leading to anaemia, malnutrition, eosinophilla and pneumonia (Sharma *et al.*, 2011). The *S. asoca* leaf extracts in aqueous, methanoland chloroform are reported to show antihelmintic activity due to paralysis and death of adult earthworm, *Pheretima posthuman*, which resembles the human round worm parasite. The methanol extract was most effective as compared to standard drug albendazole. The phytoconstituents of the methanol extract such as alkaloids, tannins, flavonoids and saponins may be responsible for the anti-helmintic activity (Sharma *et al.*, 2011).

Genetic Studies

Karyomorphological studies using root tipshave showed somatic cells with 17 sets of homologous pair of chromosomes (Deepa *et al.,* 2013).

The complete chloroplast (cp) genome/plastome was isolated and sequenced, and showed a total of 5,206,216,851 paired-end filtered reads with 151 base pairs. The total plastome length was 137,743 base pairs with GC content of 35.26% and 126 coding genes. This molecular signature can be employed to distinguish the genuine *Saraca asoca* samples from its adulterants (Ali *et al.*, 2020). The chloroplast (cp) genome/ plastome contains helpful genetic information due to its conservative rate of evolution, as it has uniparental

inheritance. The rbc*L* and mat*K* genes were established for molecular typing of *S. asoca* in molecular phylogenetic analysis (Ali *et al.*, 2021).

The phylogenetic analysis and evolutionary interrelationships of *S. asoca* was investigated with other members of the Detarieae tribe belonging to the sub family Caesalpiniodeae, using the mat*K* chloroplast gene. The analysis implied that *S. asoca* may have undergone synonymous substitutions during evolution and it showed close relationship with *Saraca palembanica*, *S. declinata*, *Endertia spectabilis* and *Lysidice rhodostegia* (Saha *et al.*, 2013).

Genetic Diversity

Molecular marker-based methods can be employed for characterization of the DNA and identification of genetic diversity analysis.

Inter-simple Sequence Repeat (ISSR) markers have been employed to evaluate the genetic variation between 11 populations of the tree species with 106 accessions gathered from Western Ghats (Hedge et al., 2018). Using UPGMA and Bayesian model, 2 clusters of comparable grouping styles were constructed. UPGMA dissimilarity analysis revealed close association of genotypes based on the geographical locations (Hedge et al., 2018). These genetic diversity and genetic relationships were further supported using ISSR and found genotype collected from different geographical locations showed high genetic variability among them (Hedge et al., 2019). In another study involving ISSR, 15 genotypes were collected from different geographical locations, from a total of 40 ISSR markers analyzed 28 markers showed polymorphic bands. A total of 149 bands were amplified, of which 133 were seen to be polymorphic. This showed higher genetic variability in the genotypes which were collected from different geographical locations (Yadav et al., 2020).

Random Amplified Polymorphic DNA (RAPD) profiling was conducted in a total of 165 individuals from 5 natural populations in Odisha, which showed a comparatively high genetic diversity and polymorphism among the population, hence recommending *in situ* conservation (Senapati *et al.*, 2012). Florescent labeled RAPD primer analysis based on 6 accessions from Hyderabad exhibited high polymorphism, indicating even a small population size can show high genetic diversity within the species (Mohan *et al.*, 2017). RAPD fingerprinting analysis on 7 wild and 3 cultivated populations from Western Ghats showed substantial genetic diversity amongst the natural populations (Saini *et al.*, 2018).

DNA fingerprinting and genetic profiling by Amplified Fragment Length Polymorphism (AFLP) technique was employed to compare and differentiate *Saraca asoca* from adulterants like *Polyalthia longifolia* bark samples. *Polyalthia longifolia* showed high level of genetic diversity as compared to *Saraca asoca* (Sharma *et al.*, 2018). Sumangala *et al.*, (2013) identified 10 novel microsatellite markers which could be used for studying population genetics of the species.

DNA Barcoding

DNA barcoding technique has been used to differentiate adulterants in 25 samples available in the market. The market and *S. asoca* specimens were barcoded employing 2 chloroplastic regions rbc*L* and psbA-trnH to develop Biological Reference Material (BRM) i.e. sample with the unique identification number. From the amplified DNA barcoded region of BRM sequences only 3 matched with the rbc*L* region and only one matched with the psbA-trnH region, thereby indicating prevalence of widespread adulteration (Urumarudappa *et al.*, 2016).

Conclusions

Saraca asoca offers many promising prospects in traditional herbal therapy and modern medicine. The tree is apparently a universal panacea for its usefulness in many different ailments. The versatile plant has a variety of phytochemical constituents which are responsible for its many fold activities. However, further studies and endeavors are required, especially to evaluate the in vivo biological activities for better validation of the efficacy of its different constituents. Further, detailed phytochemical studies on leaves and flowers could help to identify them as prospective substitutes for the bark and also help to develop a probable phytomarker which may lead to the identification of genuine Saraca asoca and prevent adulteration. Owing to its varied uses, the demand for genuine bark is constantly on the rise. Overexploitation, indiscriminate use along with habitat destruction has led tosharp reduction of its population. Molecular characterization using genetic markers may help in identifying different genotype pools which may be useful for future conservation, improvement and management of the species. Transcriptome analysis may help to identify genes involved in seed viability and could also help in improving and increasing its shelf life. Thus, the conservation practice will require innovative biotechnology tools. Lastly government initiatives, experts support and creating awareness among the public may be beneficial to the already dwindling population. In summary, S. asoca has been extensively studied with reference to its profile, phytochemistry and various biological activities. However, newerstudies and findings would enhance the existing therapeutic significance of S. asoca and encourage its future use in modern medicine, whereas novel biotechnological and genetic approaches may help in its conservation.

Acknowledgement

The authors are thankful to Prof. S. R. Yadav, Department of Botany, Shivaji University, Kolhapur, Maharashtra for providing the images of *Saraca asoca*. The authors are grateful for constructive suggestions by the reviewer.

Declarations

Authors declre no conflict of interest

References

- Ahmad F, Misra L, Tewari R, Gupta P, Gupta V K and Darokar MP (2016) Isolation and HPLC profiling of chemical constituents of *Saracaasoca* stem bark. *Ind. J. Chem* **55B:** 353-361.
- Ahmad S R and Ghosh P (2022) A systematic investigation on flavonoids, catechin, β-sitosterol and lignin glycosides from *Saracaasoca* (ashoka) having anti-cancer and antioxidant properties with no side effects. *J. Ind. Chem. Soc.* **99(1)**:100293. DOI:10.1016/j.jics.2021.100293
- Ali M A, Kim S-Y, Pan T K, Hemaid F A, Elshikh M S, Elangbam M S, Lee J, Farah M A and Al-Anazi K M (2020)Complete chloroplast genome of vulnerable medicinal plant *Saracaasoca* (Fabaceca). *Mitochondrial DNA Part B* **5(1):** 754-755.
- Ali M A, Pan T K, Gurung A B, Farah M A, Al-Hemaid F, Alanazi K M, Elangbam M, Lee J, Pandey S K, OliurRehman M and Kim S-Y (2021) Plastome of *Saracaasoca* (Detarioideae, Fabaceae): annotation, comparison among subspecies and molecular typing. *Saudi J. Biol.Sci.* **28(2):** 1487–1493. doi: 10.1016/j. sjbs.2020.12.008
- Asokan A and Thangavel M (2014) Invitro cytotoxic studies of crude methanolic extract of *Saracaasoca* bark extract. *ISOR J. Pharma. Biol. Sci.* **4(9):** 26-30.
- Banerjee P, Roy P and Nath D (2021) Chemopreventive potential of a phytofabricated silver nanoformulation using bark extract of *Saracaasoca* in AML mice *in vivo* and *in vitro*. *J. Nanomed*. **4(1):** 1036.
- Baranwal V (2014) Asoka: Herbal boon to gynecological problems-an overview of current research. *Ayurveda J. Health.* **12(4):** 58-62.
- Begum S N, Ravikumar K and Ved D K (2014) 'Asoka'- an important medicinal plant, its market scenario and conservation measures in india. *Curr. Scie.***107(1):** 26-28.
- Borokar A A and Pansare T A (2017) Plant Profile, Phytochemistry and Pharmacology of Ashoka (*Saracaasoca* (Roxb.)De Wilde) - A comprehensive Review. *Int. J. Ayurvedic Herb Med.* **7(2)**: 2524-2541.
- Chakraborty R, Sen S, Deka M K, Rekha B and Sachan S (2014) Antimicrobial evaluation of *Saracaasoca* leaves extract by disk diffusion method. *J.Pharma.Chem.Biol.Sci.* **1(1):** 1-5.
- Chauhan S (2019) Floral and pollination biology of *Saracaasoca* (Roxb.) De Wilde (Caesalpinioideae). *Int. J.Plant Repro.Biol.* **11(1):** 31-37.
- Deepa K S, Santoshkumar A V, Rekha K and Joseph J (2013) Karyotype analysis in *Saracaasoca* (Roxb.) de Wilde. *J. Trop. Agric.* **51(1):** 135-139.
- Devan A S and Warrier R R(2021) *Saracaasoca* morphology and diversity across its natural distribution in India. *Int. J. Complement. Alt. Med.* **14(6):** 317-323.
- Ghosh S, Majumder M, Majumder S, Ganguly N K and Chatterjee B P (1999) Saracin: a lectin from *Saracaasoca* seed integument induces apoptosis in human T-lymphocytes. *Arch. Biochem. Biphy.* **371(2):** 163-168.
- Gill M, Kinra M, Rai A, Chamallamudi M R and Kumar N (2018) Evaluation of antidepressant activity of methanolic extract of *Saracaasoca* bark in a chronic unpredictable mild stress model. *Neuroreport* **29(2):** 134-140.
- Gupta M, Sasmal S and Mukherjee A (2014) Therapeutic effects of acetone extract of *Saracaasoca* seeds on rats with adjuvant-induced arthritis via attenuating inflammatory responses. *ISRN Rheumatol*.:959687. doi: 10.1155/2014/959687. eCollection 2014

- Hazarika P, Kalita S and Choudhary K (2021) Restorative effects of *Saracaasoca*methanolic bark extract on levonorgestrel affected reproductive cycle and hormone levels in female albino rats. *U. P. J. Zool.***3**: 15-22.
- Hedge S, Pai S R and Roy S (2017) Combination of DNA isolation and RP-HPLC analysis method for bark samples of *Saracaasoca* and its adulterant. *3 Biotech*. 2017 Jul;7(3):208. doi: 10.1007/ s13205-017-0791-9. Epub 2017 Jun 30. PMID: 28667648; PMCID: PMC5493565.
- Hedge S, Pai S R, Bhagwat R M, Saini A, Rathore P K, Jalapure S S, Hedge H V, Sugunan A P, Gupta V S, Kholkute S D and Roy S (2018) Genetic and phytochemical investigations for understanding population variability of the medicinal important tree *Saracaasoca* to help develop conservation strategies. *Phytochemistry* **156**: 43-54.
- Hegde S, Pai S R, Bhagwat R M, Saini A, Rathore P K, Jalapure S S, Hedge H V, Sugunan A P, Gupta V S, Kholkute S D and Roy S (2019) Population genetic and phytochemical dataset of *Saracaasoca*: A traditionally important medicinal tree. *Data Brief* **25**: 104173.
- IUCN https://www.iucnredlist.org/species/ 34623/9879360# population

IUCN https://www.iucnredlist.org/species/34623/9879360

- Jeenathunisa N and Ranjan S (2021)*In vitro* cytotoxic studies of *Saracaasoca* bark extracts on HT-29 cancer cell line. *Research J. Pham. Tech.* **14(1):** 42-46.
- Khatoon S, Singh N, Kumar S, Srivastava N, Rathi A and Mehrotra S (2009) Authentication and quality evaluation of an important Ayurvedic drug - Ashoka bark. *J. Sci. Ind. Res.* **68**: 393-400.
- Kumar S, Narwal S, Kumar D, Singh G, Narwal S and Arya R (2012) Evaluation of antihyperglycemic and antioxidant activities of *Saracaasoca* (Roxb.) De Wild leaves in streptozotocin induced diabetic mice. *Asian Pacific J Tropical Disease* **2(3)** :170-176.
- Kumar V (2016) Rapid molecular authentication of Ashoka (*Saracaasoca* (Roxb.) de Wild), vulnerable medicinal plant species and its adulterant *Polyalthialongifolia*Benth by the development of SCAR markes and multiplex-PCR. *Res. J. Biotech.* **10:** 59-68.
- Mishra S B and Vijaykumar M (2014) Anti-hyperglycemic and antioxidant effect of *Saraca asoca* (Roxb. De Wilde) flowers in Streptozotocin-Nicotimomuide induced diabetic rats: A therapeutic study. *J Bioanal Boomed* **S 12**: 003. doi:10.4172/1948-593X.S12-003
- Mishra S B and Vijaykumar M (2015) Floral morphology and HPLC fingerprint profile of *Saracaasoca* (Roxb.) De Wilde (Caesalpinaceae): A consecrated tree of India. *Int. Lett. Nat. Sci.* **43:** 54-62.
- Mohan Ch, Kistamma S, Vani P and Reddy A N (2016) Biological activities of different parts of *Saraca asoca* an endangered valuable medicinal plant. *Int. J. Curr. Microbiol. App. Sci.* **5(3)** : 300-308.
- Mohan Ch, Reddy M S, Kumar S M, Manzelat S F and Cherku P D (2017) RAPD studies of *Saracaasoca* by florescent-labeled primers and development of micropropagation protocol for its conservation. *Int. J. App. Agric.Res.***12(2)**: 137-151.
- Mukhopadhyay M K, Shaw M, Nath D (2017) Chemopreventive potential of major flavonoid compound of methanolic bark extract of *Saracaasoca* (Roxb.) in benzene induced toxicity of acute myeloid leukemia mice. *Pharmacogn Mag.* **13(Suppl 2): S**216-223. doi: 10.4103/pm.pm_326_15. Epub 2017 Jul 11.
- Patwardhan A, PimputkarM ,Mhaskar M, Prerna A, Barve N, Gunaga R, Mirgal A, Salunkhe C and Vasudeva R (2016)

Distribution and population status of threatened medicinal tree *Saracaasoca* (Roxb.) De Wilde from Sahyadri-Konkan ecological corridor. *Curr. Sci.* **111(9)** 1500-1506.

- Patwardhan A, Pimputkar M, Mhaskar M, Prerna A, Barve N and Vasudeva R (2014) Status of *Saraca asoca*: An endangered medicinal plant species of conservation concern from Northern Western Ghats biodiversity Hotspot. *Biodiversity Watch* **3**: 90-102.
- Pradhan P, Joseph L, Gupta V, Chulet R, Arya H, Verma R and Bajpai A (2009) *Saraca asoca* (Ashoka): a review. *J. Chem. Pharm. Res.* **1(1):** 62-71.
- Preeti B, Bharti A, Sharma A N and Singh V (2012) A review on Saraca indica plant. Int. Res. J. Pharm. **3(4)**:80-84.
- Sadhu S K, Khatun A, Phattanawasin, P, Ohtsuki T and Ishibashi M (2007)Lignan glycosides and flavonoids from *Saraca asoca* with antioxidant activity. *J. Nat. Med* **61:** 480-482.
- Saha J, Gupta K and Gupta B (2013) Phylogenetic analysis and evolutionary relationships of *Saracaasoca* with their allied taxa (Tribe-Detarieae) based on the chloroplast mat*K* gene. *J. Plant Biochem. Biotechnol.* **24(1):** 65-74.
- Saha J, Mitra T, Gupta and Mukherjee S (2012) Phytoconstituents and HPTLC analysis in *Saraca asoca* (Roxb.) Wilde. *Int. J. Pharm. Pharm. Sci.* **4:** 96-99.
- Saha J, Mukherjee S, Gupta K and Gupta B (2013) High-performance thin-layer chromatographic analysis of antioxidants present in different parts of *Saraca asoca* (Roxb.) de Wilde. *J. Pharm. Res.* **7:** 798-803.
- Saini A, Hedge S, Hedge H V, Kholkute S D and Roy S (2018) Assessment of genetic diversity of *Saraca asoca* (Roxb.) De Wilde: a commercially important, but endangered forest tree species in Western Ghats, India. *New Zealand J. Forest. Sci.* **48:** 17-28.
- Santapau H and Henry A N (1998) A dictionary of the flowering plants in India, National Institute of Science publication (CSIR), New Delhi. pp152
- Saravanan S, Babu N P, Pandikumar P and Ignacimuthu S (2011) Therapeutic effect of *Saraca asoca* (Roxb.) Wilde on lysosomal enzymes and collagen metabolism in adjuvant induced arthritis. *Inflammopharmacology* **19(6):** 317-325.
- Sasmal S, Majumdar S, Gupta M, Mukherjee A and Mukherjee P K (2012) Pharmacognostical, phytochemical and pharmacological evaluation for the antipyretic effect of the seeds of *Saraca asoca* Roxb. *Asian Pac. J. Trop. Biomed.* 2(10): 782-786.
- Senapati S K, Das G K, Aprajita S and Rout G R (2012) Assessment of genetic variability in the Asoka tree in India. *Biodiversity* **13(1):** 16-23.
- Shahid A P, Salini S, Sasidharan N, Padikkala J, Raghavamenon A C and Babu T D (2015) Effect of *Saracaasoca* (Asoka) on estradiol-induced keratinizing metaplasia in rat uterus. *J. Basic Clin. Physiol.Pharmacol.* **26(5):** 509-515.
- Sharma A Gupta S, Sachan S Mishra A and Banerji A (2011) Antihelmintic activity of the leaf of *Saraca indica* Linn. *Asian J. Pharm. Life Sci.* **1(4):** 391-395.
- Sharma P C, Yelne M B, Dennis T J, Joshi A and Billore K V (2000) Database on medicinal plants used in Ayurveda. National Agricultural Library, US Department of Agriculture.
- Sharma P, Sharma S, YadavS, Srivastava A, Varma A, Luhana K and Shrivastava N (2018) Categorization of *Saraca asoca* and *Polyalthia longifolia*, using chemical and genetic fingerprinting associated with multivariate statistical analysis. J. Appl. Res. Med. Arom. Plants **9:** 91-96.

- Shirolkar A, Gahlaut A, Chhillar A K and Dabur R (2013) Quantitative analysis of catechins in *Saraca asoca* and correlation with antimicrobial activity. *J. Pham. Anal.* **3**: 421-428.
- Sil S, De K K and Ghosh A (2021) Phylogenetic analysis of six different species of *Saraca* L. (Fabaceae, Caesalpinioideae) based on chloroplast mat*K* gene. *Biodiversitas* **22(9):** 3880-3889.
- Sil S, Mallick T, De K K, Pramanik A and Ghosh A (2018) Comparative morphological study of three species of *Saraca* L.(Fabaceae) by the statistical approach to find out the logic of potential morphological markers. *Beni-Suef Univ. J. Basic Appl. Sci.* **7(4):** 612-619.
- Singh S, Ananta Krishna T H, Kamalraj S, Kuriakose G C, Valayil J M and Jayabaskaran (2015) Phytomedicinal medicinal importance of *Saracaasoca* (Ashoka): an exciting past, an emerging present and a promising future. *Curr. Sci.* **109(10)**: 1790-1801.
- Smitha G R and Thondaiman(2016) Reproductive biology and breeding systems of *Saraca asoca* (Roxb.) De Wilde: vulnerable medicinal plant. *Springer Plus* **5**:2025. https://doi. org/10.1186/s40064-016-3709-9
- Srinivasa K, Kumara S (and)Neetha C S 2018 Evaluation of anticonvulsant activity of *Saracaasoca* (Roxb.) Wilde flower in Swiss albino mice. *Nat. J.Physiol. Pharmacy Pharmacology* 8(6): 858-862.
- Sumangala R C, Shaanker R U, Dayanandan S, Vasudeva R and Ravikanth G (2016) Identification of novel microsatellite markers for *Saraca asoca*, a medicinal important tree species in India. *J. Genet.* **92(Suppl 2):** 93. https://doi.org/10.1007/ s12041-013-0277-4
- Swamy A H M V, Patel U M, Koti B C, Gadad P C, Patel N L and Thippeswamy A H M (2013) Cardioprotective effect of *Saraca indica* against cyclophosphamide induced cardiotoxicity in rats: a biochemical, electrocardiographic

and histopathological study. *Indian J. pharmacol.* **45(1):** 44-48.

- Swar G, Shailajan S and Menon S (2017) Activity based evaluation of a traditional ayurvedic medicinal plant: *Saraca asoca* (Roxb.) de Wilde flowers as estrogenic agents using ovariectomized rat model. *J. Ethanopharmcol.* **195:** 324-333.
- Thilagam E, Chidambaram K, Raviteja C, Vahana T and Vasudevan P (2021) Anti-hyperglycemic and hypolipidemic effects of *Saraca asoca* (Roxb.) Wilde. Flowers in alloxan-treated diabetic rats. *J. Pharm. Pharmacogn. Res.* **9(1):** 58-68.
- Urumarudappa S K J, Gogna N, Newmaster S G, Venkatarangaiah K, Subramanyam R, Saroja S G, Gudasalamani R, Dorai K and Ramanan U S (2016) DNA barcoding and NMR spectroscopybased assessment of species adulteration in the raw herbal trade of *Saracaasoca* (Roxb.) Willd, an important medicinal plant. *Int. J. Legal Med.* **130(6):** 1457-1470.
- Verma A, Jana G K, Chakraborty R, Sen S, Sachan S and Mishra A (2010) Analgesic activity of various leaf extracts of *Saracaindica* Linn. *Der Pharmacia* Let. **2(3):** 352-357.
- Verma P, Paswan S K, Vishwakarma V K, Saxena P Rao C V and Shrivastava S (2020) Evaluation of the antiulcer activity of ethanolic leave extract of *Saracaasoca* against ethanol, pylorus ligature and indomethacin induced ulcer in albino rats. *Curr. Bioactive Compounds* **16(8)** : 1191-1196.
- WHO (2005) World Health Organization, National policy on traditional medicine and regulation of herbal medicines: Report of WHO global survey.
- Yadav S S, Waoo A A and Tiwari S K (2020) Identification of genetic relationship among *Saracaasoca* genotypes using Inter Simple Sequence Repeat markers. *Biosc. Biotech. Res. Comm.* **13(4):** 2072-2076.
- Yadav S S, Woo AA and Choure K (2019) Review of medicinally important plant species *Saraca asoca* (Roxb.). *Int. J. Adv. Res.***7(9)**: 154-166.