

# STRATEGIES FOR GROWING LOW GRAIN ARSENIC RICE IN ARSENIC CONTAMINATED SOILS

## SUDHAKAR SRIVASTAVA<sup>1</sup>, SURABHI AWASTHI<sup>2</sup>, RESHU CHAUHAN<sup>2</sup>, SANJAY DWIVEDI<sup>2</sup> AND R.D. TRIPATHI<sup>2</sup>\*

<sup>1</sup>Plant Stress Biology Laboratory, Institute of Environment and Sustainable Development, Banaras Hindu University, Varanasi-221005, U.P., India, <sup>2</sup>Plant Ecology & Environmental Science Division, CSIR-National Botanical Research Institute, Lucknow-226 001, U.P., India \*Corresponding author email: tripathird@gmail.com DOI:

Arsenic is a carcinogenic element whose widespread contamination in India and other parts of the world threatens crop cultivation and public health. Rice is one of the major crops affected due to As contamination in field since rice is known to accumulate As most effectively among crop plants. The problem of As in rice grains has been shown to affect few months old children to elderly people. A number of diseases and ailments arise due to As toxicity. Hence, there is a need to grow rice in As contaminated fields sustainably with high yields and with low As in grains. In Indian perspectives, the solutions must be low-cost and feasible for local application considering the poor income status of Indian farmers and small fields. The research efforts have been put towards for the past about one and half decades and various agronomic, biological and transgenic approaches have been tested. The present review article discusses the problem of As in rice and sheds light on prospective solutions to tackle the problem.

Keywords: Arsenic, elemental interaction, microbial inoculation, transgenic rice

Arsenic (As) is present naturally in all soils in several minerals (Smedley & Kinniburgh 2002). Arsenic exists in several inorganic [As(III) and As(V)] and organic [monomethylarsonic acid (MMAV), dimethylarsinic acid (DMAV)] (Planer-Friedrich et al. 2017, Awasthi et al. 2017). It is recognized as a carcinogenic element by the International Agency for Research on Cancer (IARC). Arsenic exposure leads to several ailments including skin lesions, hyperkeratosis, hyper-/hypo-pigmentation, cardiovascular diseases, diabetes and cancer etc. (Mitra et al. 2020). Arsenic finds application in agriculture sector as insecticides, pesticides and herbicides, such as calcium arsenite and dimethylarsenic acid (Bencko & Foong 2017). Arsenic is also used in glass. leather preservatives, poisonous baits, and semiconductors. It also finds application in the treatment of several diseases (Torka et al. 2016).

Over the years, As contamination of the environment has continuously increased in several parts of the world owing to natural biogeochemical processes as well as various anthropogenic activities (Ravenscroft et al., 2013). Arsenic contamination is a global menace threatening the health of millions of people (Majumdar and Banik 2019, Ishikawa et al. 2019). Arsenic contaminated groundwater is used for drinking and irrigation purposes and therefore it enters into the food chain through crop plants (Tripathi et al. 2007; Meharg et al. 2009). Arsenic accumulation beyond maximum permissible levels has been reported in a number of crop plants, vegetables, fruits, fishes, mushrooms, sea animals, and a number of food products (Upadhyay et al. 2019). Rice is the most affected crop due to As contamination (Upadhyay et al. 2019) and As accumulation in rice is expected to increase further in future in the conditions of predicted climate change (Muehe et al. 2019). Rice is widely cultivated throughout the world. In South- and Southeast Asia, rice is commonly cultivated in flooded conditions with rainwater and groundwater. Since the groundwater has become As contaminated, its use has resulted into As build up in soil in past about 30 years (Upadhyay et al. 2019). The continuous

(Upadhyay *et al.* 2019). The continuous flooding generates reducing conditions in paddy fields which favourably increase the concentration of As(III) form of As since the speciation of As depends on pH and Eh of soil (Kerl *et al.* 2018, Majumdar *et al.* 2018). Further, the physiological (oxgen release through roots, silica hyperaccumulator plant) and molecular (high expression of silicic acid transporters involved in As(III) uptake and transport) features contribute to high As accumulation in rice (Awasthi *et al.* 2017). Arsenic levels in soil are also influenced by presence of other elements like phosphate, manganese and sulfur and also by organic matter content (Srivastava *et al.* 2016; Zafeiriou *et al.* 2019, Maguffin *et al.* 2020). The factors affecting As concentration and speciation in paddy fields can also be managed to regulate As accumulation in rice plants (Upadhyay *et al.* 2020).

The As accumulating behavior of rice ensues from a number of transporters involved in the uptake, translocation and sequestration of As (Awasthi et al. 2017). The aquaglyceroporins mediate As(III) uptake and transport (Ma et al., 2008; Lindsay and Maathuis, 2017) while phosphate transporters perform that for As(V) (Catarecha et al. 2007: Wang et al. 2016). The vacuolar sequestration of As and hence, tissue specific localization, is mediated by ATPbinding cassette (ABC)-type transporters (Song et al. 2010, 2014). However, ABC-type transporters mediate As transport only when it is complexed with phytochelatins (PCs), the oligomers of glutathione (GSH) involved in metal complexation in plants (Grill et al. 1987). Thus, enzymes of thiol metabolism and increased synthesis of cysteine, GSH and PCs also play crucial role in As accumulation and localization (Srivastava et al. 2010; Shukla et al. 2012). The complexation of As by PCs also helps in toxicity alleviation as As becomes bound and free As ion is not available for reaction with biological molecules (Srivastava et al. 2007). One of the important enzymes in As metabolism is arsenate reductase (AR) that mediates the reduction of As(V) to As(III). A few ARs like high As content 1;1/1;2/4 (HAC1;1/1;2/4) have been identified in rice (Shi et al. 2016; Xu et al. 2017). A number of antioxidants also help in the alleviation of As toxicity through regulation of reactive oxygen species (ROS) and by avoiding oxidative stress (Srivastava *et al.* 2011, Chauhan *et al.* 2020). Various metabolic processes are regulated by molecules (GABA), hormones (jasmonates, abscissic acid (ABA)), microRNAs, kinases and transcription factors (Pathare *et al.* 2013; Srivastava *et al.* 2013, Castrillo *et al.* 2013, Yu *et al.* 2012). The available knowledge about mechanism of As tolerance can be used to develop transgenic rice plants having low grain As (Tripathi *et al.* 2007). The present article focuses on agronomic and molecular strategies that can be implemented to grow safely and to have low As level in rice grains.

Screening of Low grain arsenic accumulating cultivars: After screening of more than 500 rice germplasms, including a collection of popularly cultivated cultivars in India and entries for initial evaluation trails (IETs), were conducted for three years in both seasons (Boro, the summer and Aman, the Rainfed) in collaboration with Rice Research Station (RRS), Chinsurah, West Bengal, On the basis of yield and grain As level, 25 rice germplasms were selected for multi-locational field trials. The trials were conducted at six sites of West Bengal viz., Gaighata, Durgapur, Beldanga, Chinsurah, Purbosthali and Birnagar having different levels of As contamination (10-25mg As/Kg) in soil for three years (2012- 2014). From the multilocational field trials total nine rice cultivars viz., Nayanmoni, Gotrabhog CN1646-2, CN1643-3, and IET-4786 and popular rice genotypes IR-36, Satya, IR72406-B-R-3-2-2-1 and Pusa Basmati-1 were identified as low grain As accumulating cultivars (Table -1). These low grain As accumulating cultivars has good yield/hectare and will be suitable for cultivation in almost all As affected districts of The benefits of these cultivars to the WB. farmers will be derived from increased income in As contaminated areas due to resistance to As induced yield losses. Some of these variety can also be used by the farmers of other As contaminated states of India including Indo-Gangetic plain (West Bengal, Chattisgarh, Bihar, Uttar Pradesh, Uttarakhand, Punjab etc.)

Table 1: Grain arsenic accumulation in widely cultivated rice cultivars of West Bengal, India

|    | Rice Genotynes                                   | District/ Zone in which cultivated   | Grain arsenic   | Vield/ hectare |
|----|--|--|---|----------------|
|    | Rice Genotypes                                   | District Zone in which cultivated  | level ( $\mu g kg^{-1} dw$ )<br>during field trials<br>at three sites*<br>2008-2010 | Ticlu/ nectare |
| А. | Popular rice genotypes                           |  |   |                |
|    | IR-36  | Nadia, Murshidabad, Malda,<br>Bardhaman, Howrah, Hooghly, Bankura  | 120-911   | 4.50 - 5.00    |
|    | Satya  | Different parts of Hooghly   | 116-336   | 5.00 - 5.50    |
|    | IR72406-B-R-3-2-2-1                              | Different parts of Hooghly   | 345-1267  | 3.00 - 3.50    |
|    | Pusa Bas-I                                       | Hooghly, Bardhaman, Nadia, Bankura<br>and some Basmati growing areas of India  | 129-487   | 4.00 - 4.50    |
| В. | NBRI screened low As accumulating rice genotypes |  |   |                |
|    | Nayanmoni  | Mainly in Nadia district   | 46-202  | 3.50 - 4.00    |
|    | Gotrabhog (IET 19226)                            | Nadia, Hooghly and Bardhaman   | 64-174  | 3.00 - 3.50    |
|    | CN1646-2   | Recently developed by RRS and now<br>popularly grown in Hooghly, Bardhaman,<br>Nadia, Bankura, Malda   | 43-156  | 5.50 - 6.00    |
|    | CN1643-3   | Recently developed by RRS and now<br>popularly grown in Hooghly, Bardhaman,<br>Nadia, Bankura.   | 73-186  | 4.00 - 4.50    |
|    | IET-4786   | Almost all rice growing districts of West<br>Bengal like Hooghly, Malda,<br>Murshidabad, Bardhaman, Howrah, 24<br>Parganas (N&S), Bankura, Midnapore<br>(East & West). | 99-241  | 4.50 - 5.00    |

without any yield penalty as they are well adopted for different agro-climatic zones.

Agronomic strategies for As reduction in rice: Agronomic strategies constitute methods that can be implemented by farmers during rice cultivation so as to reduce As accumulation in rice plants. Arsenic level in soil and its speciation in soil changes according to the cultivation practice and a number of physical, chemical and biological factors of soil (Li *et al.* 2014, Kumarathilaka *et al.* 2018, Muehe *et al.* 2019). Therefore, the appropriate management of rice cultivation practices can reduce As bioavailability and its accumulation by rice plants. A number of agronomic management practices have been tested by the researchers in

this direction, such as irrigation management (aerobic irrigation, sprinkler irrigation), elemental supplementation (nitrogen, selenium, silica, etc.), hormonal /chemical supply (salicylic acid, GABA, thiourea etc.), soil inversion, and inoculation of microorganisms (Srivastava *et al.* 2019; Huhmann *et al.* 2019; Seyfferth *et al.* 2019, Li *et al.* 2019, Awasthi *et al.* 2018, Chauhan et al., 2017; Moreno-Jimenez *et al.*, 2014). The following discussion presents various strategies and positive and negative points associated to each one.

#### Abiotic arsenic reduction strategies Elemental supplementation affects As

bioavailability and reduces As accumulation in rice: From the rhizosphere in soil to cytoplasm of cell, elements interact and therefore, influence the bioavailability and toxicity of each other in a dynamic manner. The interaction of As with several elements like silica (Si), phosphorus (P), selenium (Se), sulfur (S), and nitrogen (N) has been studied (Tripathi et al. 2013, Dixit et al. 2015, Chauhan et al. 2017, Srivastava et al. 2019). Silica has been the most studied as As(III) and Si use the same pathway for uptake and transport in plants and hence it is expected that Si may compete well with As(III) and reduce its accumulation in rice. Further, Si is a beneficial element for rice plants (Tripathi et al. 2013). So additional Si supply improves growth of plants and its potential to tolerate As toxicity. Li et al. (2009) observed significant reduction in the level of straw and grain total As when rice plants were supplemented with silica gel in a pot experiment. However, an increase in DMA level in grains was noticed. The effect of two different silica containing minerals namely diatomaceous earth and silica gel was tested by Seyfferth and Fendorf (2012). They found that diatomaceous earth did not cause any change or further increased the level of As in grains. However, Si gel addition decreased As in grains. Hence, variable mineralogy and available Si influences As accumulation differently. Tripathi et al. (2013) conducted laboratory experiments and reported that Si supplementation reduced As accumulation and improved As tolerance in rice seedlings. Two contrasting rice cultivars i.e. As tolerant (Triguna) and As sensitive (IET-4786) were exposed to two concentrations of AsIII (10 and  $25 \,\mu\text{M}$ ) and silicic acid (0.5 and 1 mM Si). The higher concentration of Si (1 mM) during As stress showed significant reduction of As accumulation in shoot in both the cultivars. Supplementation of Si (1 mM) during As stress ameliorated the As induced oxidative stress in tolerant rice cultivar not only due to lowering of As accumulation but also through improved thiol and antioxidative system of Triguna compared to IET-4786 (Tripathi et al. 2013).

However, in a field study, Lee et al. (2014) observed that Si supply led to increase in both As and Si concentrations in rice plants. Syu et al. (2016) reported that application of Si significantly reduced the accumulation of Si in rice shoots due to competition in uptake of As and Si. A study by Khan and Gupta in 2018 reported that priming of rice seed with Si along with As reduced the accumulation of As. Recently, Cui et al. (2020) reported that foliar application of nanoparticle of Si provided tolerance against As stress by maintaining the integrity the cell wall and by reducing the oxidative stress and As accumulation. The supplementation of Si rice husk to the soil also helped to reduce As accumulation in rice without hampering the nutrient availability (Seyfferth et al. 2016). A recent study by Leksungnoen et al. (2019) reported that biochar produced from silica rich rice husk plays an important role in reduction of As accumulation in rice grain (Leksungnoen et al., 2019).

Sulfur (S) is an important element in the context of As detoxification as this is a constituent of GSH and PCs involved in As complexation. It has been noticed that external GSH supply reduces As toxicity and accumulation in rice seedlings (Shri et al. 2009). Further, an inhibitor of GSH biosynthesis, buthionine-L-sulfoximine (BSO) was supplied to rice plants. It was found that BSO reduced GSH and PC levels and thus increased As transport to flag leaves and grains (Vezza et al. 2019). Dixit et al. (2015a) also observed in hydroponics study that supply of high S level reduced As translocation to shoot as this restricted As to roots itself. An opposite result was seen when low S supply was given. The mechanism of S induced As stress tolerance in rice was also evaluated by Dixit et al. (2015b). They showed that S alleviated As toxicity by modulating several protein levels, amino acids and thiol metabolism. Zhang et al. (2016) found that sulfur application decreased As concentration in rice grains by 44%. The response was found to be correlated to changes in expression of a number of genes including

transporters. Srivastava et al. (2016) conducted a laboratory experiment where they supplied low and zero S level to rice seedlings in presence of As. It was observed that even in condition of zero S supply, plants tend to accumulate PCs to tackle As toxicity. The combined toxicity of As and S depleted condition showed loss of photosynthetic pigments and increased accumulation of anthocyanins. However, even under depleted S supply, increased synthesis and consumption of S-containing thiols was observed upon As exposure. The study inferred that rice plant combated As induced stress and S depleted condition by reduction of As accumulation and by altering subcellular distribution of As (Srivastava et al. 2016). Kumar et al. (2019) reported that the over-expression of highaffinity sulphate transporters. OsSultr1;1 provides tolerance against different abiotic stresses under limiting sulphur condition. An increased formation of iron plaque on S supplementation is also reported, which act as a barrier for As and leads to reduction in As accumulation in plants (Shakoor et al. 2019).

Phosphate is known to compete with As(V) for uptake and transport in plants. Although As exists mostly in the form of As(III) in paddy fields, there is expected conversion of As(III) back to As(V) around rice roots. This is due to the fact that rice roots release oxygen and therefore just close to roots, oxygenic conditions are promoted. The changes in microbial community around roots are also reported (Srivastava et al. 2018). Hence, despite the presence of As mostly in the form of As(III), rice rhizospheric As chemistry might be completely different. Considering these facts in mind, phosphate supplementation has also been tested by researchers. However, in most of the studies, phosphate addition caused an increase in As accumulation in rice plants (Hossain et al. 2009, Talukder et al. 2011, Wu et al. 2011). This was considered to be due to replacement of As(V) adsorbed on to soil particles and iron plaque on rice roots and thus making As(V) more bioavailable to plants.

Luan et al. (2018) reported a major vacuolar phosphate transporter (VPT1) that is involved in vacuolar inorganic phosphate (Pi) homeostasis and thus play an important role in As(V) tolerance in Arabidopsis. This study reported that mutant VPT1 elevated the levels of Pi in the cytosol, thus suppressed the expression of PHT1-type transporters leading to reduced accumulation of As. Another study by Xie et al. (2019) reported the role of OsNLA1 in As uptake and tolerance. The increased expression of OsNLA1 is reported during AsV stress and authors revealed that OsNLA1 play an important role in regulation of phosphate transporters and thus provide As tolerance.

Selenium is an important element which acts as an antioxidant and is a constituent of several enzymes (Chauhan et al. 2019). Kumar et al. (2014) demonstrated improvement in growth and tolerance of rice seedlings to As stress upon Se supplementation. Chauhan et al. (2017) also reported improved plant growth and nutrient status of rice plants on Se supplementation. A marked reduction in As accumulation as well as increased accumulation of Se in rice plant was observed. The As toxicity amelioration in presence of Se was achieved via increase in enzymatic and non-enzymatic antioxidant, such as phenolic compounds (Gallic acid, protocatechuic acid, caffeic and Rutin) (Chauhan et al. 2017) Transcriptome and proteome profiling of rice plant also suggested that supplementation of Se reduced the As induced oxidative stress and toxicity in rice plant by up-regulation of expression of key As transporters and defense responsive genes as well as their respective proteins (Chauhan et al. 2020).

Hence, elemental fertilization and optimum management of elemental nutrition of plants can help improve growth of rice plants in As contaminated fields. The strategy can also assist in reducing As level in rice grains and lead to higher yields of better quality rice grains.

Water influences As dynamics in soil and As levels in rice plants: The irrigation of rice fields with groundwater is the major source of As for rice fields. The water is itself an important regulator of As biogeochemistry in soil owing to its influence of soil physicial and chemical properties and microbial activity (Upadhyay et al. 2020). Rice is a semi-aquatic plant and sufficient water availability plays a key role in achieving proper rice growth and productivity (Islam et al. 2019). Conventionally, the fields are flooded throughout the growth period of rice plants. However, there are other cultivation methods of rice aimed to reduce water use such as aerobic cultivation, alternate wetting and drying and sprinkler irrigation. These approaches were designed to reduce excessive water consumption by rice crops. These also hold worth in the light of increasing water shortage which is expected to worsen further in future. Further, such water conservation approaches might also allow rice cultivation in more aerobic conditions and reduce As accumulation in rice grains.

Duxbury and Panaullah (2007) compared Boro rice grown either in flooded conditions or raised bed cultivation in Faridpur, Bangladesh. They observed decrease in rice yields in both flooded and raised bed conditions. However, raised bed cultivation caused lower decline in rice yields and also reduced grain As level. Talukder et al. (2011) demonstrated increase in grain yield (13% increase) and grain As (62% decrease) in raised bed cultivation. Moreno-Jimenez et al. (2014) found that sprinkler irrigation application to rice plants reduced grain total As by one-third in only one application as compared to traditional irrigation. When the sprinkler irrigation was used for longer duration, an even greater reduction in As accumulation in rice grains was achieved. However, sprinkler irrigation generates aerobic conditions that increase Cd bioavailability to rice plants. Hence, an increase in Cd concentration in rice grains was observed (Moreno-Jimenez et al. 2014). In the experiments of Arao et al. (2009) and Li et al.

(2009) also aerobic rice cultivation resulted in significant increase in Cd concentration while a decline in Se and Fe levels. Spanu *et al.* (2012) reported that sprinkler irrigation is helpful to reduce the As accumulation by 50%. Yang *et al.* (2019) reported alternate wetting and drying of rice during cultivation plays an important role in significant reduction in As accumulation. A study by Islam *et al.* (2019) also reported that water management via alternate wetting and drying is an effective way to reduce As accumulation.

Supplementation of chemical and biomolecules enhances As tolerance and reduces As accumulation of rice : Srivastava *et al.* (2014) conducted a lab study where they supplemented As exposed rice seedlings with a redox active molecule, thiourea (TU). Thiourea supplementation was found not only ameliorate As stress in rice seedlings but also to reduce As concentrations in shoot parts. The responses could be linked to regulation of As transporters (Lsi1 and Lsi2) in redox dependent manner along with other transporters as well as biochemical modifications.

Singh *et al.* (2015) applied salicylic acid (SA) to rice seedlings and noticed improved growth and stress in rice plants. SA application also decreased As translocation from root to shoot. The positive results were attributable to a number of biochemical and molecular changes. Singh et al. (2017) reported that nitric oxide and SA are the signaling molecules and provide protection against As toxicity by modulating signaling response against As. Kumar *et al.* (2017) showed the role of  $\gamma$ aminobutyric acid (GABA) in As tolerance by modulating expression of As transporters viz., LSi-1 and LSi-2. GABA has been suggested to act as a central regulator of As stress responses of plants in Indian mustard plants (Pathare et al. 2013). In rice plants also, Pathare et al. (2016) suggested involvement of GABA and 14-3-3 protein interaction in As stress responses.

#### Biotic arsenic reduction strategies Bacterial inoculants

Microbes constitute a diverse group which includes bacteria, algae, fungi that are associated with improvement of plant growth as well as increase nutrient status. They also play an important role to provide tolerance against As toxicity by As methylation and conversion to non-toxic forms, siderophore procution and by reduction of the bioavailability of As. Dolphen and Thiravetyan (2019) reported that microbes are efficient tool to reduce As accumulation. Inoculation of B. pumilus (an endophytic bacterium) with leonardite reduced the accumulation of As in rice by down-regulating of Lsi-1 and Lsi-2. Moens et al. (2020) reported that inoculation of a hyperaccumulator rhizobacterial strain, Ochrobactrum tritici to the rice during As stress decreased the As induced toxicity in rice and also improved the yield of rice plant. In some studies, the use of more than one bacterium in the form of a consortium was done. A study by Nookongbut et al. (2018) reported that microbial consortium of bacteria (Rhodopseudomonas palustris C1 and Rubrivivax benzoatilyticus C31) not only enhanced the rice growth but also reduced As accumulation in rice by producing siderophores, indole acetic acid and exopolymeric substances. Thus it can be concluded that microbial consortium ameliorates As toxicity and reduces As uptake in rice and also increases the yield.

## Fungal inoculants

Arbuscular mycorrhizal fungi impart tolerance against various stresses including heavy metals. Li *et al.* (2016) reported that *Rhizophagus intraradices* significantly reduced As concentrations in rice grains by transforming inorganic As into less toxic organic form in rice. Zhang *et al.* (2016) found that AMF inoculation decreased the proportion of inorganic As in grains of a As tolerant rice variety. Li *et al.* (2011) also demonstrated that *Rhizophagus irregularis* inoculation to lowland rice and *Glomus geosporum*. inoculation to upland rice resulted in an increase in grain yield and grain P/As ratio. Li et al. (2011, 2013) suggested that the response of an inoculum of AMF varies different rice varieties and therefore the best candidate or the best combination of AMF inoculants should be used. Wu et al. (2015) reported that inoculation of micorrhizal fungi Glomus geosporum BGC HUN02C, G. versiforme BGC GD01B and G. mosseae BGC GD01A increased the vield of rice plant as well as mitigated As toxicity in rice. Mohd et al. (2017) reported that an endophytic fungus Piriformospora indica colonized the rice roots during As stress by modulating antioxidant enzyme activities. Thie led to balanced redox status of the cell which and protected photosynthetic machinery of the plants. The supplementation of fungus ameliorated As toxicity by different mechanisms such as by reducing the bioavailability of As to the plant, via biotransformation of the toxic form of As into insoluble particulate matter and by modulating the antioxidative system of the plant. Several studies also reported that filamentous fungus viz., Penicillium sp., Aspergillus sp., Trichoderma sp., Cladosporium sp., Rhizopus sp. and Westerdykella sp. are potent candidate to mitigate As toxicity due to their biotransforming potential but among these Penicillium sp. and Aspergillus sp. are most As tolerant fungal species (Batista et al. 2016, Segura et al. 2018). Thus the application of fungal sp. in paddy fields might be a good alternative for bioremediation in Ascontaminated soils and amelioration of As toxicity in rice as well as for reducing As in rice grains.

## **Transgenic approaches**

For the sustainable solution of the problem and for achieving desired objectives like growth and yield improvement and As level reduction in grains, transgenic variety development has been considered a viable objective. The targeted strategies were either focused to enhance tolerance of rice plants to As stress through changes in the level of antioxidants, GSH and PCs or to reduce As uptake and transport via changes in transporter proteins (reviewed in Srivastava et al. 2012). Since, As has some organic species which are volatile, one strategy has been to express bacterial genes in rice plants that can increase As volatilization to the environment and thus reduce As in grains. Meng et al. (2011) developed transgenic rice expressing arsM gene (Sadenosylmethyltransferase) of Rhodopseudomonas palustris. They found induced As methylation and eventual volatilization of As (10-fold higher volatile As) in transgenic rice. Duan et al. (2012) utilized ACR3 (Arsenic Compounds Resistance 3) gene, responsible for As extrusion from cells to the medium, from Sachharomyces cerevisiae for the development of transgenic rice. In the transgenic lines, As concentrations in shoot and root were 30% lower than WT. Importantly, there was a decline in husk and grain As in transgenic lines (about 20-30%) even when grown under continuous flooding. Oryza sativa C-type ATP-binding cassette (ABC) transporter (OsABCC1) was chosen to develop transgenic plants and the gene was knocked down. The mutant lines accumulated less As in shoots than wild type. However, mutants showed more As accumulation in flag leaf, husk and brown rice (13-18-fold higher) than WT (Song et al. 2014). This could be due to the fact that in absence of OsABCC1, lower amounts of As-PC complexes would be sequestered to vacuole. Shri et al. (2014) developed transgenic rice lines expressing phytochelatin synthase gene isolated from Ceratophyllum demersum for achieving higher As complexation with PCs. In the obtained transgenic lines, significant increase in PCS activity and in the level of PCs was noticed. This led to higher As accumulation in root and shoot owing to greater ability of transgenic rice to complex As. It positively resulted in significant decline in grain and husk As concentration as compared to WT. Hence, altering either the complexation of As or the

sequestration of As-PC complexes can be

potential strategies for developing low grain As

accumulating cultivar in near future. Tiwari et al. (2014) has demonstrated involvement of NRAMP1 transporter of rice (OsNRAMP1) in As transport. The transporter was found to be localized in plasma membrane of endodermis and percicyle cells and is involved in xylem loading of As. Expression of OsNRAMP1 in Arabidopsis although provided tolerance to plants but resulted in greater As accumulation n both root and shoot as compared to WT. The transporter there provides an opportunity for low grain As rice development through alteration of its expression. Kumar et al. (2013) expressed a rice lambda class of glutathione-Stransferase gene (OsGSTL2) in Arabidopsis and found that it could provide tolerance to heavy metals including As. Glutathione-Stransferases are involved in GSH-mediated detoxification of xenobiotics including metals and hence, this gene can be helpful in the development of rice with improved tolerance to As. The expression of WaarsM gene of Westerdykella aurantiaca in rice was able to convert toxic As into less toxic methylated As and reduced As accumulation in rice by its volatilization and methylation (Verma et al. 2018). Shi et al. (2016) utilized arsenic reductase genes from rice plants (OsHAC1;1 and OsHAC1:2) and developed knock out and overexpressing lines. The overexpressing rice lines showed greater reduction of As(V) to As(III), higher efflux of As(III) to external medium while reduced As accumulation in shoots. The knock out lines showed an opposite response. Similar results were observed in case of another arsenic reductase gene from rice, OsHAC4 (Xu et al. 2017). Thus, increased expression of arsenic reductase gene in roots allows plants to rapidly get rid of As through reduction to As(III) and subsequent efflux. Deng et al. (2018) generated transgenic rice plants with the use of two genes. One Saccharomyces cerevisiae yeast cadmium factor (ScYCF1) and other OsABCC1. Both genes were expressed in the root cortical and internode phloem cells. Further, another gene from Escherichia coli, y-glutamylcysteine synthetase ( $\gamma$ -ECS) was also expressed. The

transgenic rice plants exhibited significantly reduced translocation of As firstly from rootto-shoot and also from internode-to-grain during grain filling stage. An eventual reduction in grains was found to be up to 70% without any negative impact on rice plant growth and development.

#### **Concluding remarks**

A number of approaches have been studied at laboratory and field level and mixed responses have been found in terms of As stress mitigation and As reduction in rice. It seems reasonable to cultivate screened and selected low grain As accumulating cultivars for As prone areas of different agro-climatic zones. However, future studies need to be more focused and accurate to provide a novel, feasible and cost-effective solution for growing rice in As contaminated areas. Future research also needs to evaluate the impact of changing climatic conditions and to assess the effects of multiple stressors to come out with sustainable solutions to tackle As problem in rice.

## REFERENCES

Arao T, Kawasaki A, Baba K, Mori S and Matsumoto S 2009 Effects of water management on cadmium and arsenic accumulation and dimethylarsinic acid concentrations in Japanese rice. *Environmental Science & Technology*, **43**(24), pp.9361-9367.

Awasthi S, Chauhan R, Dwivedi S, Srivastava S, Srivastava S and Tripathi R D, 2018. A consortium of alga (Chlorella vulgaris) and bacterium (Pseudomonas putida) for amelioration of arsenic toxicity in rice: A promising and feasible approach. Environmental and Experimental Botany, 150, pp.115-126.

Awasthi, S., Chauhan, R., Srivastava, S. and Tripathi, R.D., 2017. The journey of arsenic from soil to grain in rice. Frontiers in Plant Science, 8, p.1007. Batista, B.L., Barião, C.V., Souza, J.M.O., Paulelli, A.C.C., Rocha, B.A., de Oliveira, A.R.M., Segura, F.R., Braga, G.Ú.L., Tonani, L., von Zeska-Kress, M.R. and Barbosa Jr, F., 2016. A low-cost and environmentallyfriendly potential procedure for inorganic-As remediation based on the use of fungi isolated from rice rhizosphere. Journal of environmental chemical engineering, 4(1), pp.891-898.

Bencko, V. and Foong, F.Y.L., 2017. The history of arsenical pesticides and health risks related to the use of Agent Blue. Annals of agricultural and environmental medicine,

## 24(2), pp.312-316.

Castrillo, G., Sánchez-Bermejo, E., de Lorenzo, L., Crevillén, P., Fraile-Escanciano, A., Mohan, T.C., Mouriz, A., Catarecha, P., Sobrino-Plata, J., Olsson, S. and del Puerto, Y.L., 2013. WRKY6 transcription factor restricts arsenate uptake and transposon activation in Arabidopsis. The Plant Cell, 25(8), pp.2944-2957.

Catarecha, P., Segura, M.D., Franco-Zorrilla, J.M., García-Ponce, B., Lanza, M., Solano, R., Paz-Ares, J. and Leyva, A., 2007. A mutant of the Arabidopsis phosphate transporter PHT1; 1 displays enhanced arsenic accumulation. The Plant Cell, 19(3), pp.1123-1133. Chauhan, R., Awasthi, S., Indoliya, Y., Chauhan, A.S., Mishra, S., Agrawal, L., Srivastava, S., Dwivedi, S., Singh, P.C., Mallick, S. and Chauhan, P.S., 2020. Transcriptome and proteome analyses reveal selenium mediated amelioration of arsenic toxicity in rice (Oryza sativa L.). Journal of Hazardous Materials, 390, p.122122.

Chauhan, R., Awasthi, S., Srivastava, S., Dwivedi, S., Pilon-Smits, E.A., Dhankher, O.P. and Tripathi, R.D., 2019. Understanding selenium metabolism in plants and its role as a beneficial element. Critical Reviews in

Environmental Science and Technology, 49(21), pp.1937-1958.

Chauhan, R., Awasthi, S., Tripathi, P., Mishra, S., Dwivedi, S., Niranjan, A., Mallick, S., Tripathi, P., Pande, V. and Tripathi, R.D., 2017. Selenite modulates the level of phenolics and nutrient element to alleviate the toxicity of arsenite in rice (Oryza sativa L.). Ecotoxicology and environmental safety, 138, pp.47-55.

Cui, J., Li, Y., Jin, Q. and Li, F., 2020. Silica nanoparticles inhibit arsenic uptake into rice suspension cells via improving pectin synthesis and the mechanical force of the cell wall. Environmental Science: Nano, 7(1), pp.162-171.

Deng, F., Yamaji, N., Ma, J.F., Lee, S.K., Jeon, J.S., Martinoia, E., Lee, Y. and Song, W.Y., 2018. Engineering rice with lower grain arsenic. Plant biotechnology journal, 16(10), pp.1691-1699.

Dixit, G., Singh, A.P., Kumar, A., Dwivedi, S., Deeba, F., Kumar, S., Suman, S., Adhikari, B., Shukla, Y., Trivedi, P.K. and Pandey, V., 2015. Sulfur alleviates arsenic toxicity by reducing its accumulation and modulating proteome, amino acids and thiol metabolism in rice leaves. Scientific reports, 5(1), pp.1-16.

Dixit, G., Singh, A.P., Kumar, A., Mishra, S., Dwivedi, S., Kumar, S., Trivedi, P.K., Pandey, V. and Tripathi, R.D., 2016. Reduced arsenic accumulation in rice (Oryza sativa L.) shoot involves sulfur mediated improved thiol metabolism, antioxidant system and altered arsenic transporters. Plant Physiology and Biochemistry, 99, pp.86-96

Dixit, G., Singh, A.P., Kumar, A., Singh, P.K., Kumar, S., Dwivedi, S., Trivedi, P.K., Pandey, V., Norton, G.J., Dhankher, O.P. and Tripathi, R.D., 2015. Sulfur mediated reduction of arsenic toxicity involves efficient thiol metabolism and the antioxidant defense J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:100

system in rice. Journal of Hazardous Materials, 298, pp.241-251.

Dolphen, R. and Thiravetyan, P., 2019. Reducing arsenic in rice grains by leonardite and arsenic–resistant endophytic bacteria. Chemosphere, 223, pp.448-454.

Duan, G., Kamiya, T., Ishikawa, S., Arao, T. and Fujiwara, T., 2012. Expressing ScACR3 in rice enhanced arsenite efflux and reduced arsenic accumulation in rice grains. Plant and Cell Physiology, 53(1), pp.154-163.

Duxbury, J.M. and Panaullah, G.M., 2007. Remediation of Arsenic for Agriculture Sustainability. Food Security and Health in Bangladesh, pp.1-28.

Grill, E., Winnacker, E.L. and Zenk, M.H., 1987. Phytochelatins, a class of heavy-metalbinding peptides from plants, are functionally analogous to metallothioneins. Proceedings of the National Academy of Sciences, 84(2), pp.439-443.

Hossain, M.B., Jahiruddin, M., Loeppert, R.H., Panaullah, G.M., Islam, M.R. and Duxbury, J.M., 2009. The effects of iron plaque and phosphorus on yield and arsenic accumulation in rice. Plant and Soil, 317(1-2), pp.167-176.

Huhmann, B., Harvey, C.F., Uddin, A., Choudhury, I., Ahmed, K.M., Duxbury, J.M., Ellis, T. and van Geen, A., 2019. Inversion of high-arsenic soil for improved rice yield in Bangladesh. Environmental science & technology, 53(7), pp.3410-3418.

Ishikawa, S., Arao, T. and Makino, T., 2019. Agronomic Strategies for Reducing Arsenic Risk in Rice. In Arsenic Contamination in Asia (pp. 181-198). Springer, Singapore.

Islam, S., Rahman, M.M. and Naidu, R., 2019. Impact of water and fertilizer management on arsenic bioaccumulation and speciation in rice

plants grown under greenhouse conditions. Chemosphere, 214, pp.606-613.

Kerl, C.F., Rafferty, C., Clemens, S. and Planer-Friedrich, B., 2018. Monothioarsenate uptake, transformation, and translocation in rice plants. Environmental science & technology, 52(16), pp.9154-9161.

Khan, E. and Gupta, M., 2018. Arsenic–silicon priming of rice (Oryza sativa L.) seeds influence mineral nutrient uptake and biochemical responses through modulation of Lsi-1, Lsi-2, Lsi-6 and nutrient transporter genes. Scientific reports, 8(1), pp.1-16.

Kumar, A., Singh, R.P., Singh, P.K., Awasthi, S., Chakrabarty, D., Trivedi, P.K. and Tripathi, R.D., 2014. Selenium ameliorates arsenic induced oxidative stress through modulation of antioxidant enzymes and thiols in rice (Oryza sativa L.). Ecotoxicology, 23(7), pp.1153-1163.

Kumar, N., Dubey, A.K., Upadhyay, A.K., Gautam, A., Ranjan, R., Srikishna, S., Sahu, N., Behera, S.K. and Mallick, S., 2017. GABA accretion reduces Lsi-1 and Lsi-2 gene expressions and modulates physiological responses in Oryza sativa to provide tolerance towards arsenic. Scientific reports, 7(1), pp.1-11.

Kumar, S., Asif, M.H., Chakrabarty, D., Tripathi, R.D., Dubey, R.S. and Trivedi, P.K., 2013. Expression of a rice Lambda class of glutathione S-transferase, OsGSTL2, in Arabidopsis provides tolerance to heavy metal and other abiotic stresses. Journal of hazardous materials, 248, pp.228-237.

Kumar, S., Khare, R. and Trivedi, P.K., 2019. Arsenic-responsive high-affinity rice sulphate transporter, OsSultr1; 1, provides abiotic stress tolerance under limiting sulphur condition. Journal of hazardous materials, 373, pp.753-762. J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:101

Kumarathilaka, P., Seneweera, S., Meharg, A. and Bundschuh, J., 2018. Arsenic speciation dynamics in paddy rice soil-water environment: sources, physico-chemical, and biological factors-a review. Water research, 140, pp.403-414.

Lee, C.H., Huang, H.H., Syu, C.H., Lin, T.H. and Lee, D.Y., 2014. Increase of As release and phytotoxicity to rice seedlings in Ascontaminated paddy soils by Si fertilizer application. Journal of hazardous materials, 276, pp.253-261.

Leksungnoen, P., Wisawapipat, W., Ketrot, D., Aramrak, S., Nookabkaew, S., Rangkadilok, N. and Satayavivad, J., 2019. Biochar and ash derived from silicon-rich rice husk decrease inorganic arsenic species in rice grain. Science of The Total Environment, 684, pp.360-370.

Li, B., Zhou, S., Wei, D., Long, J., Peng, L., Tie, B., Williams, P.N. and Lei, M., 2019. Mitigating arsenic accumulation in rice (Oryza sativa L.) from typical arsenic contaminated paddy soil of southern China using nanostructured  $\alpha$ -MnO2: Pot experiment and field application. Science of the Total Environment, 650, pp.546-556.

Li, H., Chen, X.W. and Wong, M.H., 2016. Arbuscular mycorrhizal fungi reduced the ratios of inorganic/organic arsenic in rice grains. Chemosphere, 145, pp.224-230.

Li, H., Man, Y.B., Ye, Z.H., Wu, C., Wu, S.C. and Wong, M.H., 2013. Do arbuscular mycorrhizal fungi affect arsenic accumulation and speciation in rice with different radial oxygen loss?. Journal of hazardous materials, 262, pp.1098-1104.

Li, H., Wu, C., Ye, Z.H., Wu, S.C., Wu, F.Y. and Wong, M.H., 2011. Uptake kinetics of different arsenic species in lowland and upland rice colonized with Glomus intraradices. Journal of hazardous materials, 194, pp.414-421.

Li, J., Dong, F., Lu, Y., Yan, Q. and Shim, H., 2014. Mechanisms controlling arsenic uptake in rice grown in mining impacted regions in South China. PLoS One, 9(9), p.e108300.

Li, R.Y., Ago, Y., Liu, W.J., Mitani, N., Feldmann, J., McGrath, S.P., Ma, J.F. and Zhao, F.J., 2009. The rice aquaporin Lsi1 mediates uptake of methylated arsenic species. Plant Physiology, 150(4), pp.2071-2080.

Li, R.Y., Stroud, J.L., Ma, J.F., McGrath, S.P. and Zhao, F.J., 2009. Mitigation of arsenic accumulation in rice with water management and silicon fertilization. Environmental Science & Technology, 43(10), pp.3778-3783.

Lindsay, E.R. and Maathuis, F.J., 2017. New molecular mechanisms to reduce arsenic in crops. Trends in plant science, 22(12), pp.1016-1026.

Luan, M., Liu, J., Liu, Y., Han, X., Sun, G., Lan, W. and Luan, S., 2018. Vacuolar Phosphate transporter 1 (VPT1) affects arsenate tolerance by regulating phosphate homeostasis in Arabidopsis. Plant and Cell Physiology, 59(7), pp.1345-1352.

Ma, J.F., Yamaji, N., Mitani, N., Xu, X.Y., Su, Y.H., McGrath, S.P. and Zhao, F.J., 2008. Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. Proceedings of the National Academy of Sciences, 105(29), pp.9931-9935.

Maguffin, S.C., Abu-Ali, L., Tappero, R.V., Pena, J., Rohila, J.S., McClung, A.M. and Reid, M.C., 2020. Influence of manganese abundances on iron and arsenic solubility in rice paddy soils. Geochimica et Cosmochimica Acta, 276, pp.50-69.

Majumdar, A., Barla, A., Upadhyay, M.K., Ghosh, D., Chaudhuri, P., Srivastava, S. and Bose, S., 2018. Vermiremediation of metal (loid) s via Eichornia crassipes phytomass extraction: a sustainable technique for plant J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:102

amelioration. Journal of environmental management, 220, pp.118-125.

Majumder, S. and Banik, P., 2019. Geographical variation of arsenic distribution in paddy soil, rice and rice-based products: a meta-analytic approach and implications to human health. Journal of environmental management, 233, pp.184-199.

Meharg, A.A., Williams, P.N., Adomako, E., Lawgali, Y.Y., Deacon, C., Villada, A., Cambell, R.C., Sun, G., Zhu, Y.G., Feldmann, J. and Raab, A., 2009. Geographical variation in total and inorganic arsenic content of polished (white) rice. Environmental science & technology, 43(5), pp.1612-1617.

Meng, X.Y., Qin, J., Wang, L.H., Duan, G.L., Sun, G.X., Wu, H.L., Chu, C.C., Ling, H.Q., Rosen, B.P. and Zhu, Y.G., 2011. Arsenic biotransformation and volatilization in transgenic rice. New Phytologist, 191(1), pp.49-56.

Mitra, A., Chatterjee, S. and Gupta, D.K., 2020. Environmental arsenic exposure and human health risk. In Arsenic Water Resources Contamination (pp. 103-129). Springer, Cham.

Moens, M., Branco, R. and Morais, P.V., 2020. Arsenic accumulation by a rhizosphere bacterial strain Ochrobactrum tritici reduces rice plant arsenic levels. World Journal of Microbiology and Biotechnology, 36(2), p.23.

Mohd, S., Shukla, J., Kushwaha, A.S., Mandrah, K., Shankar, J., Arjaria, N., Saxena, P.N., Narayan, R., Roy, S.K. and Kumar, M., 2017. Endophytic fungi Piriformospora indica mediated protection of host from arsenic toxicity. Frontiers in microbiology, 8, p.754.

Moreno-Jiménez, E., Meharg, A.A., Smolders, E., Manzano, R., Becerra, D., Sánchez-Llerena, J., Albarrán, Á. and López-Piñero, A., 2014. Sprinkler irrigation of rice fields reduces

grain arsenic but enhances cadmium. Science of the total environment, 485, pp.468-473.

Muehe, E.M., Wang, T., Kerl, C.F., Planer-Friedrich, B. and Fendorf, S., 2019. Rice production threatened by coupled stresses of climate and soil arsenic. Nature communications, 10(1), pp.1-10.

Nookongbut, P., Kantachote, D., Megharaj, M. and Naidu, R., 2018. Reduction in arsenic toxicity and uptake in rice (Oryza sativa L.) by As-resistant purple nonsulfur bacteria. Environmental Science and Pollution Research, 25(36), pp.36530-36544.

Pathare, V., Srivastava, S. and Suprasanna, P., 2013. Evaluation of effects of arsenic on carbon, nitrogen, and sulfur metabolism in two contrasting varieties of Brassica juncea. Acta physiologiae plantarum, 35(12), pp.3377-3389.

Pathare, V., Srivastava, S., Sonawane, B.V. and Suprasanna, P., 2016. Arsenic stress affects the expression profile of genes of 14-3-3 proteins in the shoot of mycorrhiza colonized rice. Physiology and Molecular Biology of Plants, 22(4), pp.515-522.

Planer-Friedrich, B., Kuhnlenz, T., Halder, D., Lohmayer, R., Wilson, N., Rafferty, C. and Clemens, S., 2017. Thioarsenate toxicity and tolerance in the model system Arabidopsis thaliana. Environmental Science & Technology, 51(12), pp.7187-7196.

Ravenscroft, P., McArthur, J.M. and Hoque, M.A., 2013. Stable groundwater quality in deep aquifers of Southern Bangladesh: The case against sustainable abstraction. Science of the Total Environment, 454, pp.627-638.

Segura, F.R., Paulelli, A.C.C., Braga, G.Ú.L., dos Reis Pedreira Filho, W., Silva, F.F. and Batista, B.L., 2018. Promising filamentous native fungi isolated from paddy soils for J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:103

arsenic mitigation in rice grains cultivated under flooded conditions. Journal of environmental chemical engineering, 6(4), pp.3926-3932.

Seyfferth, A.L. and Fendorf, S., 2012. Silicate mineral impacts on the uptake and storage of arsenic and plant nutrients in rice (Oryza sativa L.). Environmental science & technology, 46(24), pp.13176-13183.

Seyfferth, A.L., Amaral, D., Limmer, M.A. and Guilherme, L.R., 2019. Combined impacts of Si-rich rice residues and flooding extent on grain As and Cd in rice. Environment international, 128, pp.301-309.

Seyfferth, A.L., Morris, A.H., Gill, R., Kearns, K.A., Mann, J.N., Paukett, M. and Leskanic, C., 2016. Soil incorporation of silica-rich rice husk decreases inorganic arsenic in rice grain. Journal of Agricultural and Food Chemistry, 64(19), pp.3760-3766.

Shakoor, M.B., Riaz, M., Niazi, N.K., Ali, S., Rizwan, M., Arif, M.S. and Arif, M., 2019. Recent advances in arsenic accumulation in rice. In Advances in Rice Research for Abiotic Stress Tolerance (pp. 385-398). Woodhead Publishing.

Shi, S., Wang, T., Chen, Z., Tang, Z., Wu, Z., Salt, D.E., Chao, D.Y. and Zhao, F.J., 2016. OsHAC1; 1 and OsHAC1; 2 function as arsenate reductases and regulate arsenic accumulation. Plant Physiology, 172(3), pp.1708-1719.

Shri, M., Dave, R., Diwedi, S., Shukla, D., Kesari, R., Tripathi, R.D., Trivedi, P.K. and Chakrabarty, D., 2014. Heterologous expression of Ceratophyllum demersum phytochelatin synthase, CdPCS1, in rice leads to lower arsenic accumulation in grain. Scientific reports, 4, p.5784.

Shri, M., Kumar, S., Chakrabarty, D., Trivedi,

P.K., Mallick, S., Misra, P., Shukla, D., Mishra, S., Srivastava, S., Tripathi, R.D. and Tuli, R., 2009. Effect of arsenic on growth, oxidative stress, and antioxidant system in rice seedlings. Ecotoxicology and environmental safety, 72(4), pp.1102-1110.

Shukla, D., Kesari, R., Mishra, S., Dwivedi, S., Tripathi, R.D., Nath, P. and Trivedi, P.K., 2012. Expression of phytochelatin synthase from aquatic macrophyte Ceratophyllum demersum L. enhances cadmium and arsenic accumulation in tobacco. Plant cell reports, 31(9), pp.1687-1699.

Singh, A.P., Dixit, G., Kumar, A., Mishra, S., Kumar, N., Dixit, S., Singh, P.K., Dwivedi, S., Trivedi, P.K., Pandey, V. and Dhankher, O.P., 2017. A protective role for nitric oxide and salicylic acid for arsenite phytotoxicity in rice (Oryza sativa L.). Plant Physiology and Biochemistry, 115, pp.163-173.

Singh, A.P., Dixit, G., Mishra, S., Dwivedi, S., Tiwari, M., Mallick, S., Pandey, V., Trivedi, P.K., Chakrabarty, D. and Tripathi, R.D., 2015. Salicylic acid modulates arsenic toxicity by reducing its root to shoot translocation in rice (Oryza sativa L.). Frontiers in Plant Science, 6, p.340.

Smedley, P.L. and Kinniburgh, D.G., 2002. A review of the source, behaviour and distribution of arsenic in natural waters. *Applied geochemistry*, *17*(5), pp.517-568.

Song, W.Y., Park, J., Mendoza-Cózatl, D.G., Suter-Grotemeyer, M., Shim, D., Hörtensteiner, S., Geisler, M., Weder, B., Rea, P.A., Rentsch, D. and Schroeder, J.I., 2010. Arsenic tolerance in Arabidopsis is mediated by two ABCC-type phytochelatin transporters. Proceedings of the National Academy of Sciences, 107(49), pp.21187-21192.

Song, W.Y., Yamaki, T., Yamaji, N., Ko, D., Jung, K.H., Fujii-Kashino, M., An, G., Martinoia, E., Lee, Y. and Ma, J.F., 2014. A rice J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:104

ABC transporter, OsABCC1, reduces arsenic accumulation in the grain. Proceedings of the National Academy of Sciences, 111(44), pp.15699-15704.

Spanu, A., Daga, L., Orlandoni, A.M. and Sanna, G., 2012. The role of irrigation techniques in arsenic bioaccumulation in rice (Oryza sativa L.). Environmental science & technology, 46(15), pp.8333-8340.

Srivastava, A.K., Srivastava, S., Mishra, S., D'Souza, S.F. and Suprasanna, P., 2014. Identification of redox-regulated components of arsenate (As V) tolerance through thiourea supplementation in rice. Metallomics, 6(9), pp.1718-1730.

Srivastava, S., Akkarakaran, J.J., Sounderajan, S., Shrivastava, M. and Suprasanna, P., 2016. Arsenic toxicity in rice (Oryza sativa L.) is influenced by sulfur supply: impact on the expression of transporters and thiol metabolism. Geoderma, 270, pp.33-42.

Srivastava, S., Mishra, S., Tripathi, R.D., Dwivedi, S., Trivedi, P.K. and Tandon, P.K., 2007. Phytochelatins and antioxidant systems respond differentially during arsenite and arsenate stress in Hydrilla verticillata (Lf) Royle. Environmental science & technology, 41(8), pp.2930-2936.

Srivastava, S., Pathare, V.S., Sounderajan, S. and Suprasanna, P., 2019. Nitrogen supply influences arsenic accumulation and stress responses of rice (Oryza sativa L.) seedlings. Journal of hazardous materials, 367, pp.599-606.

Srivastava, S., Srivastava, A.K., Suprasanna, P. and D'Souza, S.F., 2010. Comparative antioxidant profiling of tolerant and sensitive varieties of Brassica juncea L. to arsenate and arsenite exposure. Bulletin of environmental contamination and toxicology, 84(3), pp.342-346.

Srivastava, S., Srivastava, A.K., Suprasanna, P. and D'souza, S.F., 2013. Identification and

profiling of arsenic stress-induced microRNAs in Brassica juncea. Journal of experimental botany, 64(1), pp.303-315.

Srivastava, S., Srivastava, S., Bist, V., Awasthi, S., Chauhan, R., Chaudhry, V., Singh, P.C., Dwivedi, S., Niranjan, A., Agrawal, L. and Chauhan, P.S., 2018. Chlorella vulgaris and Pseudomonas putida interaction modulates phosphate trafficking for reduced arsenic uptake in rice (Oryza sativa L.). Journal of hazardous materials, 351, pp.177-187.

Srivastava, S., Suprasanna, P. and D'Souza, S.F., 2011. Redox state and energetic equilibrium determine the magnitude of stress in Hydrilla verticillata upon exposure to arsenate. Protoplasma, 248(4), pp.805-815.

Srivastava, S., Suprasanna, P. and D'souza, S.F., 2012. Mechanisms of arsenic tolerance and detoxification in plants and their application in transgenic technology: a critical appraisal. International Journal of Phytoremediation, 14(5), pp.506-517.

Syu, C.H., Huang, C.C., Jiang, P.Y., Chien, P.H., Wang, H.Y., Su, J.Y. and Lee, D.Y., 2016. Effects of foliar and soil application of sodium silicate on arsenic toxicity and accumulation in rice (Oryza sativa L.) seedlings grown in Ascontaminated paddy soils. Soil Science and Plant Nutrition, 62(4), pp.357-366.

Talukder, A.S.M.H.M., Meisner, C.A., Sarkar, M.A.R. and Islam, M.S., 2011. Effect of water management, tillage options and phosphorus status on arsenic uptake in rice. Ecotoxicology and environmental safety, 74(4), pp.834-839.

Talukder, A.S.M.H.M., Meisner, C.A., Sarkar, M.A.R. and Islam, M.S., 2011. Effect of water management, tillage options and phosphorus status on arsenic uptake in rice. Ecotoxicology and environmental safety, 74(4), pp.834-839.

Torka, P., Al Ustwani, O., Wetzler, M., Wang, E.S. and Griffiths, E.A., 2016. Swallowing a

J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:105

bitter pill–oral arsenic trioxide for acute promyelocytic leukemia. Blood reviews, 30(3), pp.201-211.

Tripathi, P., Tripathi, R.D., Singh, R.P., Dwivedi, S., Goutam, D., Shri, M., Trivedi, P.K. and Chakrabarty, D., 2013. Silicon mediates arsenic tolerance in rice (Oryza sativa L.) through lowering of arsenic uptake and improved antioxidant defence system. Ecological engineering, 52, pp.96-103.

Tripathi, R.D., Srivastava, S., Mishra, S., Singh, N., Tuli, R., Gupta, D.K. and Maathuis, F.J., 2007. Arsenic hazards: strategies for tolerance and remediation by plants. Trends in biotechnology, 25(4), pp.158-165.

Upadhyay, M.K., Majumdar, A., Suresh Kumar, J. and Srivastava, S., 2020. Arsenic in rice agro-ecosystem: Solutions for safe and sustainable rice production. Frontiers in Sustainable Food Systems, 4, p.53.

Upadhyay, M.K., Shukla, A., Yadav, P. and Srivastava, S., 2019. A review of arsenic in crops, vegetables, animals and food products. Food chemistry, 276, pp.608-618.

Verma, S., Verma, P.K., Meher, A.K., Bansiwal, A.K.,

Tripathi, R.D. and Chakrabarty, D., 2018. A novel fungal arsenic methyltransferase, WaarsM reduces grain arsenic accumulation in transgenic rice (Oryza sativa L.). Journal of hazardous materials, 344, pp.626-634.

Vezza, M.E., Luna, D.F., Agostini, E. and Talano, M.A., 2019. Glutathione, a key compound for As accumulation and tolerance in soybean plants treated with AsV and AsIII. Environmental and Experimental Botany, 162, pp.272-282.

Wang, P., Zhang, W., Mao, C., Xu, G. and Zhao, F.J., 2016. The role of OsPT8 in arsenate uptake and varietal difference in arsenate tolerance in rice. Journal of experimental botany, 67(21), pp.6051-6059.

Wu, F., Hu, J., Wu, S. and Wong, M.H., 2015. Grain yield and arsenic uptake of upland rice inoculated with arbuscular mycorrhizal fungi in As-spiked soils. Environmental Science and Pollution Research, 22(12), pp.8919-8926.

Wu, Z., Ren, H., McGrath, S.P., Wu, P. and Zhao, F.J., 2011. Investigating the contribution of the phosphate transport pathway to arsenic accumulation in rice. Plant Physiology, 157(1), pp.498-508.

Xie, M.Y., Tian, Z.H., Yang, X.L., Liu, B.H., Yang, J. and Lin, H.H., 2019. The role of OsNLA1 in regulating arsenate uptake and tolerance in rice. Journal of plant physiology, 236, pp.15-22.

Xu, J., Shi, S., Wang, L., Tang, Z., Lv, T., Zhu, X., Ding, X., Wang, Y., Zhao, F.J. and Wu, Z., 2017. OsHAC4 is critical for arsenate tolerance and regulates arsenic accumulation in rice. New Phytologist, 215(3), pp.1090-1101.

Yang, Y., Hu, H., Fu, Q., Zhu, J. and Huang, G., 2019. Water management of alternate wetting and drying reduces the accumulation of arsenic in brown rice-as dynamic study from rhizosphere soil to rice. Ecotoxicology and Environmental Safety, 185, p.109711. J. Indian bot. Soc. Sp. Issue Vol. 100(A) 2020:106

Yu, F., Qian, L., Nibau, C., Duan, Q., Kita, D., Levasseur, K., Li, X., Lu, C., Li, H., Hou, C. and Li, L., 2012. FERONIA receptor kinase pathway suppresses abscisic acid signaling in Arabidopsis by activating ABI2 phosphatase. Proceedings of the National Academy of Sciences, 109(36), pp.14693-14698.

Zafeiriou, I., Gasparatos, D., Kalyvas, G., Ioannou, D. and Massas, I., 2019. Desorption of Arsenic from Calcareous Mine Affected Soils by Phosphate Fertilizers Application in Relation to Soil Properties and As Partitioning. Soil Systems, 3(3), p.54.

Zhang, J., Zhao, C.Y., Liu, J., Song, R., Du, Y.X., Li, J.Z., Sun, H.Z., Duan, G.L. and Zhao, Q.Z., 2016. Influence of sulfur on transcription of genes involved in arsenic accumulation in rice grains. Plant Molecular Biology Reporter, 34(3), pp.556-565.

Zhang, X., Wu, S., Ren, B. and Chen, B., 2016. Water management, rice varieties and mycorrhizal inoculation influence arsenic concentration and speciation in rice grains. Mycorrhiza, 26(4), pp.299-309.