



RESEARCH ARTICLE

A Survey of Traditional Medicinal Plants in District Hapur, Uttar Pradesh: Treatment of Gynecological Disorder

Amita Sharma and Soniya Rani*

Abstract

Gynecological disorders are recognized as a major worldwide issue impacting the health of women. During the survey of district Hapur, a total of 52 ethnomedicinal plant species were recorded, along with local names, families, habits, and plant parts used. These 52 medicinal plant species that provide crude drugs pertain to 47 genera and 33 families. These plants were found to cure 12 kinds of gynecological disorders. During documentation of plants used to cure gynecological disorders, the most frequently prominent families were Fabaceae (9), Lamiaceae, Moraceae (4), Apiaceae, Apocynaceae, Amaryllidaceae, Brassicaceae, and Zingiberaceae (2). During the survey, it was reported that twenty-five families had one species each. Among the genera, *Ficus* (4 spp.), *Ocimum* (2 spp.), and *Allium* (2 spp.) were the dominant genera. Based on their ethnomedicinal uses, the most significant species were *Curcuma longa*, *Allium cepa*, *Asparagus racemosus*, and *Vitis vinifera*. This is first-hand information related to ethnomedicinal plant species that are used by the local people to treat gynecological disorders in district Hapur. This research aims to preserve the ancient treasure collected through folklore and herbal healers to treat gynecological problems. Further pharmacological survey of these plants may provide some important drugs for the treatment of common gynecological disorders that may be used as powerful medications.

Keywords: Ethnomedicinal, folklore, gynecological disorders, plant species, traditional.

Introduction

Gynecological disorders are problems of females such as abortion, infertility, menopause, leucorrhoea, gonorrhoea, delivery problems, menstrual trouble, uterine prolapse, cervical cancer, breast cancer, endometrial cancer, PCOS, etc. Gynecological disorders are regarded as an important global problem for women's health as they often include conditions in the female reproductive system. Menstrual disorders are prevalent among women. Due to their hesitation and lack of knowledge, many women decide not to consult with physicians. So, women rely primarily on the usage of indigenous herbs to treat various ailments in

rural areas (Das *et al.* 2015). In modern times, non-steroidal anti-inflammatory medications, an operation, and allopathic treatment are often used to treat gynecological diseases. Even though these treatments are very effective, there are also a lot of common potential side effects. These include nausea and vomiting from surgery or anesthesia, breast enlargement, post-hysterectomy sexual dysfunction, skin rashes or digestive issues from drug use, or more serious kidney, liver, and heart impairment from drug use, especially when used for a long time (Lawal *et al.* 2013, Panda *et al.* 2018). Furthermore, several medicines during pregnancy have a risk of harming the embryo. Due to socioeconomic factors, many women are compelled to seek abortions. Women have the option to induce abortions themselves in nations where abortion is illegal or if the healthcare system is unable to adequately care for the population. Abortion is defined as "causing miscarriage," which is a punishable offense under the Indian Penal Code (1860).

The WHO states that "women's health care is crucial." Women who reside in hamlets are very impoverished, both financially and educationally (WHO). In general, rural pregnant women would rather have their babies delivered by a trained village midwife than by a gynecologist. They are unable to visit the medical facilities and multispecialty clinics due to financial constraints and distance. In many

Department of Botany, Raghunath Girls' Post Graduate College, Meerut, Uttar Pradesh, India

***Corresponding Author:** Soniya Rani, Department of Botany, Raghunath Girls' Post Graduate College, Meerut, Uttar Pradesh, India, E-Mail: Soniyasaxena1186@gmail.com

How to cite this article: Sharma A and Rani S (2024). A Survey of Traditional Medicinal Plants in District Hapur, Uttar Pradesh: Treatment of Gynecological Disorder. J. Indian bot. Soc., 104 (3): 136-147. Doi: [10.61289/jibs2024.05.15.160](https://doi.org/10.61289/jibs2024.05.15.160)

Source of support: Nil

Conflict of interest: None.

underdeveloped nations, primary maternity care is given by traditional birth attendants, or dais. From ancient times in India, traditional birth attendants, or dais, have provided prenatal and postpartum treatment, support, and guidance. Traditional birth attendants, or dais, acquired their knowledge and experience through practice and traditions. The major source of healthcare for almost 80% of the world's population is traditional herbal medicine, while many modern pharmaceutical medications are derived from plants for many disorders (Gupta & Solanki 2013). In India, the ancient medical systems of Unani, Ayurveda, Homeopathy, and Siddha rely heavily on wild-harvested plants for over 95% of their plant-based medicines (Satyavati *et al.* 1987). Many medicinal plants have been identified in several Ayurvedic scriptures, viz. Charaka Samhita, Sushruta Samhita, Ashtanga Hridaya, etc. As earlier as we know, different types of medicinal plants and their preparations are used in different gynecological disorders, i.e. Aartavkshaya, Aartavivrridhi, Asrigdara, Phirang, Raktagulma, Somaroga, Upadansha, Yonivyapada, Yonikanda, etc., which are treated with different herbs having different Ayurvedic properties, i.e. Ashok, Amalaki, Banyan tree (Vat), Sesamum (Til), Triphala, etc. and Tandulodaka is used as a main Anupana for the treatment of Asrigdara. Honey, Milk, and Ushnodaka (lukewarm water) are also used in the treatment of different gynecological disorders (Gautam *et al.* 2017).

Due to their safety, effectiveness, and affordability, herbal medications are gaining popularity among people in both urban and rural regions in emerging nations such as China and India. Approximately 8,000 plant species are recognized for their ethnomedicinal value (Joshi 1995), and the development of innovative plant-based therapeutic products has been aided by traditional

knowledge-based formulations or indigenous traditional medicine (Katewa 2009). Very little quantitative research has been conducted on the use of traditional medicines for gynecological care, although many ethnomedicinal studies have been conducted on herbal plants. Thankfully, there is still a strong history of treating female health issues using herbal remedies. These conventional treatments, however, lack extensive documentation. There is a lack of traditional knowledge about traditional medicines due to several factors, including the rise of nuclear families, where grandmothers are missing, migration to cities, and easy access to synthetic pharmaceuticals. For women of all ages, gynecological problems constitute a major contributor to morbidity and medical expenses. Those who are having genitourinary instrumentation or catheterization, as well as young women who are sexually active and elderly, are the most in danger. Over 40% of women get a urinary tract infection at some point in their lives (Adeniyi *et al.* 2006, Tripathi *et al.* 2011, Mishra *et al.* 2013).

Because sexual disorders are socially stigmatized, the majority of women in rural areas seek treatment from traditional medical healers. Through documentation, the current research aims to preserve the priceless legacy of the local people and traditional healers on the use of herbal treatments against gynecological disorders and disseminate it to the rest of the globe. We gathered traditional knowledge about medicinal plants used to cure gynecological problems in this survey.

Materials And Methods

Investigation Area

The present survey was carried out in the district of Hapur of Uttar Pradesh State, India (Figure 1). The district is located at




Figure 1: Map of the survey area A: India B: Uttar Pradesh C: Hapur district

Table 1: Enumeration of ethnomedicinal plant species used by local people for gynecological disorders in district Hapur, Uttar Pradesh

S. No.	Diseases Name	Plant part used	Botanical Name of Plant	Common Name of Plant	Family	Formulation & application of plant part used	Active Phytochemical
1.	Amenorrhea, Hypomenorrhea & Oligomenorrhea	Root	<i>Asparagus racemosus</i> Willd.	Satmul	Asparagaceae	5 gram. root powder with warm water is consumed for 15 days.	Shatavarin 1 (Thakur <i>et al.</i> 2021)
		Leaves	<i>Gymnema sylvestris</i> R. Br.	Gurmar	Apocynaceae	Decoction of leaves is consumed.	Lupeol, Betaine, Stigmasterol (Vora <i>et al.</i> 2023)
		Fruit	<i>Piper nigrum</i> L.	Kali mirch	Piperaceae	Powdered fruit is consumed.	Piperine, Catechin, (Takooree <i>et al.</i> 2019, Afroze <i>et al.</i> 2022)
		Seed	<i>Sesamum indicum</i> L.	Til	Pedaliaceae	Seeds are consumed on an empty stomach.	Sesamin, Sesamolin, Sesaminol (Dachtler <i>et al.</i> 2003, Yavari <i>et al.</i> 2014)
2.	Breast lump (Tumor & Cancer)	Bulb	<i>Allium cepa</i> L.	Pyaz	Amaryllidaceae	Bulb is consumed orally daily.	Quercetin, Allicin (Chakraborty <i>et al.</i> 2022)
		Bark	<i>Bauhinia variegata</i> L.	Kachnar	Fabaceae	Decoction (10-20 milliliter) of the bark powder (20-30 gram) of Kachnar (<i>Bauhinia purpurea</i>) and Gorakhmundi (<i>Sphaeranthus indicus</i>) (5-10 gram) is given twice a day.	Kaempferol (Sharma <i>et al.</i> 2019)
		Bark	<i>Ficus religiosa</i> L.	Pipal	Moraceae	Bark extract is consumed.	Rutin, 3-caffeoylquinic acid, Luteolin 7-O-rutinoside, 6-C-glucosyl-8-C-arabinosylapigenin, Kaempferol-3-O-rutinoside (Murugesu <i>et al.</i> 2021)
		Root	<i>Mimosa pudica</i> L.	Chui Mui	Fabaceae	Root paste is externally applied to the breast.	L-mimosine (Xu <i>et al.</i> 2018)
		Leaves	<i>Moringa oleifera</i> Lam.	Shahjan	Moringaceae	Leaves powder with warm water is consumed twice a day.	Rutin, Niaziminin, Kaempferol (Pareek <i>et al.</i> 2023, Purwal <i>et al.</i> 2010)
		Leaves	<i>Ocimum sp.</i>	Tulsi	Lamiaceae	Decoction of leaves is consumed.	Apigenin (Way <i>et al.</i> 2004, Chorocho <i>et al.</i> 2023)
		Leaves	<i>Ricinus communis</i> L.	Arand	Euphorbiaceae	Warmed leaves with castor oil (<i>Ricinus communis</i>) are externally applied to the breast.	Coumaric acid, Epigallocatechin, Ricinoleic acid, Ricinine (Majumder <i>et al.</i> 2019)
		Seed	<i>Sesamum indicum</i> L.	Til	Pedaliaceae	Seeds are chewed on an empty stomach.	β -sitosterol (Downie <i>et al.</i> 1999, Shantha <i>et al.</i> 2014)
		Seed	<i>Trigonella foenum-graecum</i> L.	Methi	Fabaceae	Powder of seeds with cow's milk is given twice a day	Diosgenin (Goyal <i>et al.</i> 2016)
	Fruit	<i>Vitis vinifera</i> L.	Angur	Vitaceae	Fruit is consumed.	Resveratrol, Catechins, Epicatechin, Gallic acid, p-coumaric (Tsantila <i>et al.</i> 2024)	

3.	Cervical Cancer	Bulb	<i>Allium cepa</i> L.	Pyaz	Amaryllidaceae	Bulb is consumed orally daily.	Convercetin, Allicin (Wozniak <i>et al.</i> 2021)
		Clove	<i>Allium sativum</i> L.	Lehsun	Amaryllidaceae	Garlic paste is consumed orally.	Allicin (Wozniak <i>et al.</i> 2021)
		Root	<i>Asparagus racemosus</i> Willd.	Satmul	Asparagaceae	The root powder is consumed.	Rutin, Kaempferol (Wozniak <i>et al.</i> 2021)
		Root	<i>Beta vulgaris</i> L.	Chukander	Amaranthaceae	The root is consumed.	Betaines (Wozniak <i>et al.</i> 2021)
		Leaves	<i>Camellia sinensis</i> L. Ktze.	Chai	Theaceae	Leaves extract is consumed.	Epigallocatechin-3-gallate (Wang <i>et al.</i> 2018, Wozniak <i>et al.</i> 2021)
		Fruit	<i>Citrus sp.</i>	Nimbu, Santra	Rutaceae	Fruit juice is given orally.	Naringin (Ramesh & Alshatwi 2013, Wozniak <i>et al.</i> 2021)
		Rhizome	<i>Curcuma longa</i> L.	Haldi	Zingiberaceae	Powder is consumed orally.	Curcumin (Wozniak <i>et al.</i> 2021)
		Root	<i>Daucus carota</i> L.	Gajar	Apiaceae	Root is consumed.	β -carotene, α -carotene, Lutein, Lycopene, Zeaxanthin, Caffeic acid (Wozniak <i>et al.</i> 2021)
		Bark	<i>Ficus religiosa</i> L.	Pipal	Moraceae	Bark extract is consumed.	Rutin, 3-caffeoylquinic acid, Luteolin 7-O-rutinoside, 6-C-glucosyl-8-C-arabinosylapigenin, Kaempferol-3-O-rutinoside (Murugesu <i>et al.</i> 2021)
		Leaves, Fruit	<i>Ginkgo biloba</i> L.	Ginkgo	Ginkgoaceae	Decoction of leaves and fruit is given orally.	Ginkgolide B (Xu <i>et al.</i> 2020, Wozniak <i>et al.</i> 2021)
Leaves	<i>Ocimum sp.</i>	Tulsi	Lamiaceae	Decoction of leaves is given twice a day.	Apigenin (Choroshu <i>et al.</i> 2023)		
Fruit	<i>Vitis vinifera</i> L.	Angur	Vitaceae	Fruit is given orally.	Rutin, Gallic acid, Caffeic acid, Ferulic acid, Naringin (Wozniak <i>et al.</i> 2021)		
4.	Dysmenorrhea & Menorrhagia	Bark	<i>Senegalia catechu</i> (L.f.) P.J.H. Hurter & Mabb.	Khair	Fabaceae	Extract of bark is consumed orally.	(+)- Catechin, Epigallocatechin, Epigallocatechin gallate, Epicatechin gallate (Afroze <i>et al.</i> 2022)
		Bark, Root	<i>Phyllanthus reticulatus</i> Poir	Makhi	Phyllanthaceae	Root/Bark decoction is given orally twice a day for dysmenorrhea.	Lupeol, Lupeol acetate, Stigmasterol, Scopoletin (Afroze <i>et al.</i> 2022)
		Fruit	<i>Piper nigrum</i> L.	Kali mirch	Piperaceae	Powdered fruit is consumed.	Piperine, Catechin, (Afroze <i>et al.</i> 2022)
5.	Endometrial cancer	Bulb	<i>Allium cepa</i> L.	Pyaz	Amaryllidaceae	Bulb is consumed orally daily.	Quercetin, Convercetin, Allicin (Wozniak <i>et al.</i> 2021)
		Root	<i>Asparagus racemosus</i> Willd.	Satmul	Asparagaceae	The root powder is consumed.	Asparanin A (Wozniak <i>et al.</i> 2021)
		Whole plant	<i>Berberis sp.</i>	Berberberry	Berberidaceae	Extract of the whole plant is given twice a day.	Berberin (Wang <i>et al.</i> 2018)
		Leaves	<i>Camellia sinensis</i> L. Ktze.	Chai	Theaceae	Extract of leaves is consumed.	Epigallocatechin gallate (Wang <i>et al.</i> 2018, Wozniak <i>et al.</i> 2021)
		Rhizome	<i>Curcuma longa</i> L.	Haldi	Zingiberaceae	Powder is consumed orally.	Curcumin (El <i>et al.</i> 2019)

	Leaves	<i>Ocimum sp.</i>	Tulsi	Lamiaceae	Decoction of leaves is consumed.	Apigenin (Chorosh et al. 2023)	
	Seed	<i>Sida cordifolia</i> L.	Khareti	Malvaceae	The decoction of seeds is taken.	β -Sitosterole, Stigmasterol (Khurana et al. 2016)	
6.	Hypogalactia	Root	<i>Asparagus racemosus</i> Willd.	Satmul	Asparagaceae	Root powder with milk is consumed.	Shatavarin-1 (Thakur et al. 2021)
	Seed	<i>Foeniculum vulgare</i> (Fennel)	Sauf	Apiaceae	Seed is given orally.	Anethole, Estragole (Tabares et al. 2014)	
	Seed	<i>Moringa oleifera</i> Lam.	Shahjan	Moringaceae	Seed powder with cow's milk is consumed twice a day.	Arachidic acid, Docosaheptaenoic acid (Sari et al. 2020)	
	Whole plant	<i>Silybum marianum</i> (L.) Gaertn.	Doodh Patra	Asteraceae	Extract of the whole plant is consumed.	Silybin B (Dietz et al. 2016)	
	Seed	<i>Trigonella foenum-graecum</i> L.	Methi	Fabaceae	Powder of seeds with cow's milk is given twice a day	Diosgenin, Apigenin, Luteolin (Tabares et al. 2014)	
7.	Induce abortion	Gum, Bark, Flower	<i>Butea monosperma</i> (Lam.) Kuntze	Dhak	Fabaceae	Powder of bark, flower, and gum is given with warm water on an empty stomach.	Butin (Bhargava 1986)
	Root	<i>Calotropis procera</i> (Aiton) W. T. Aiton	Aak	Apocynaceae	Powder of root is consumed.	Calotropin (Wadhvani et al. 2021)	
	Rhizome	<i>Curcuma longa</i> L.	Haldi	Zingiberaceae	Powder with curd is externally applied & consumed orally.	Curcumin (Tossetta et al. 2021)	
	Whole plant	<i>Cynodon dactylon</i> Pers.	Durva	Poaceae	Juice of whole plant is given twice a day.	Apigenin (Malpani et al. 2020)	
	Seed	<i>Trigonella foenum-graecum</i> L.	Methi	Fabaceae	Powder of seeds with cow's milk is given twice a day	Gitogenin, Trigonelline (Aafi et al. 2020)	
	Leaves	<i>Justicia adhatoda</i> L.	Adusa	Acanthaceae	Decoction of leaves is given.	Vasicine (Gupta et al. 1978)	
	Root, Bark,	<i>Moringa oleifera</i> Lam.	Shahjan	Moringaceae	Root powder with warm water is consumed twice a day.	Morintides, β -sitosterol (Pareek et al. 2023, Attah et al. 2020)	
8.	Infertility in female	Root	<i>Asparagus racemosus</i> Willd.	Satmul	Asparagaceae	Root powder with warm water is consumed for 15 days.	Shatavarin 1 (Thakur et al. 2021)
	Whole plant	<i>Ocimum basilicum</i> L.	Tulsi	Lamiaceae	Decoction of the whole plant is consumed.	Eugenol (Chorosh et al. 2023)	
	Fruit	<i>Vitis vinifera</i> L.	Angur	Vitaceae	Fruit is consumed.	Resveratrol (Chorosh et al. 2023)	
9.	Leucorrhoea	Seed	<i>Albizia lebeck</i> L. Benth	Siris	Fabaceae	Seed powder with honey kept in an earthen pot in the sun for 15 days. Paste is given twice a day.	D-Catechin, Albizziahexoside A (Behera et al. 2021)
	Root	<i>Justicia adhatoda</i> L.	Adusa	Acanthaceae	Decoction of root is given.	Vasicine (Hossain et al. 2016)	
	Gum, Bark, Flower	<i>Butea monosperma</i> (Lam.) Kuntze	Dhak	Fabaceae	Powder of bark, flower, and gum (each in equal amount) is given with warm water on an empty stomach.	Butin (Sharma et al. 2011)	

	Leaves, Bark	<i>Ficus benghalensis</i> L.	Bargad	Moraceae	Leaves and bark powder with honey are given for 15 days.	Rutin, Friedelin, Leucoanthocyanin (Behera <i>et al.</i> 2021)
	Fruit, Leaves	<i>Ficus religiosa</i> L.	Pipal	Moraceae	The powder of fruit and tender leaves in equal amounts with jaggery is given twice a day for Uterine diseases.	Coumarin (Behera <i>et al.</i> 2021)
	Fruit	<i>Ficus racemose</i> L.	Gular	Moraceae	Fresh fruit is given daily.	Stigmasterol, Lupeol, Lupeol acetate, Taraxasterol ester, Hentriacontane (Behera <i>et al.</i> 2021)
	Bark	<i>Ficus virens</i> L.	Pilkhan	Moraceae	Bark extract is consumed.	Stigmasterol, Lupeol, Scutellarein (Behera <i>et al.</i> 2021)
10. Ovarian Cancer	Bulb	<i>Allium cepa</i> L.	Pyaz	Amaryllidaceae	Bulb is consumed orally daily.	Quercetin, Allicin (Wozniak <i>et al.</i> 2021)
	Leaves	<i>Brassica oleracea</i>	Gobhi	Brassicaceae	The cooked vegetable is consumed orally.	Isothiocyanates Indoles (Zhang <i>et al.</i> 2006)
	Leaves	<i>Camellia sinensis</i> L. Ktze.	Chai	Theaceae	Extract of leaves is consumed.	Epigallocatechin gallate (Wang <i>et al.</i> 2018, Wozniak <i>et al.</i> 2021)
	Rhizome	<i>Curcuma longa</i> L.	Haldi	Zingiberaceae	Powder is consumed orally.	Curcumin (Wozniak <i>et al.</i> 2021)
	Root	<i>Daucus carota</i> L.	Gajar	Apiaceae	Root is consumed.	β -carotene, α -carotene, Lycopene, Zeaxanthin (Wozniak <i>et al.</i> 2021)
	Leaves, Fruit	<i>Ginkgo biloba</i> L.	Ginkgo	Ginkgoaceae	Decoction of leaves and fruit is given orally.	Quercetin, Ginkgolide A & B (Ye <i>et al.</i> 2007, Wozniak <i>et al.</i> 2021)
	Leaves	<i>Ocimum sp.</i>	Tulsi	Lamiaceae	Extract of leaves is consumed.	Apigenin (Suh <i>et al.</i> 2015, Choroshko <i>et al.</i> 2023)
	Fruit	<i>Vitis vinifera</i> L.	Angur	Vitaceae	Fruit is consumed.	Resveratrol (Wozniak <i>et al.</i> 2021)
11. PCOS (polycystic ovary syndrome)	Leaves	<i>Aloe vera</i> (L.) Burm. f.	Gvarpatha	Asphodelaceae	Leaves juice is given on an empty stomach.	Aloe-emodin, Barbaloin (Lakshmi <i>et al.</i> 2023)
	Whole plant	<i>Berberis sp.</i>	Burberry	Berberidaceae	Extract of the whole plant is given twice a day.	Berberin (Li <i>et al.</i> 2015, Mishra <i>et al.</i> 2022)
	Fruit	<i>Cocos nucifera</i> L.	Nariyal	Arecaceae	Coconut water is consumed orally.	Skimmiwallin, Isoskimmiwallin (Bhandary <i>et al.</i> 1995, Lakshmi <i>et al.</i> 2023)
	Stigma	<i>Crocus sativa</i> L.	Kesar	Iridaceae	Saffron water is given twice a day.	Crocin (Yasmin <i>et al.</i> 2019)
	Rhizome	<i>Curcuma longa</i> L.	Haldi	Zingiberaceae	Powder is consumed orally.	Curcumin (Lakshmi <i>et al.</i> 2023)
	Bark	<i>Ficus religiosa</i> L.	Pipal	Moraceae	Bark extract is consumed daily.	β -sitosterol-D-glycoside (Behera <i>et al.</i> 2021)
	Seed	<i>Foeniculum vulgare</i> Mill.	Sauf	Apiaceae	Seed is consumed.	Anethole, Ascorbic acid, α -tocopherol, β -tocopherol, γ -tocopherol, δ -tocopherol (Lakshmi <i>et al.</i> 2023)
	Whole plant	<i>Fumaria parviflora</i> Lam.	Pittapapada	Papaveraceae	Decoction of the whole plant is consumed daily in the morning.	Protopine (Mubeen <i>et al.</i> 2022)

	Root	<i>Glycyrrhiza glabra</i> L.	Mulethi	Fabaceae	Root extract is used twice a day.	Liquiritigenin, Isoliquiritigenin, Liquiritin, Isoliquiritin, Glabridin, Glabrene (Yang <i>et al.</i> 2018, Lakshmi <i>et al.</i> 2023)
	Leaves	<i>Gymnema sylvestris</i> R. Br.	Gurmar	Apocynaceae	Decoction of leaves is consumed.	Lupeol, Betaine, Stigmasterol (Vora <i>et al.</i> 2023)
	Seed	<i>Linum usitatissimum</i> L.	Alsi	Linaceae	Extract of seed is consumed twice a day.	Secoisolariciresinol, Secoisolariciresinol diglycoside, Linolenic acid (Lakshmi <i>et al.</i> 2023)
	Whole plant	<i>Mentha spicata</i> L.	Pudina	Lamiaceae	Juice of the whole plant is consumed daily.	Lutein, Rutin, Rosmarinic acid, Caffeic acid (Lakshmi <i>et al.</i> 2023)
	Leaves	<i>Ocimum sp.</i>	Tulsi	Lamiaceae	Decoction of leaves is consumed.	Apigenin (Choroshko <i>et al.</i> 2023)
	Fruit	<i>Punica granatum</i> L.	Anar	Lythraceae	Juice of fruit is consumed daily.	Catechin, Epicatechin, Gallocatechin, Quercetine, Kaempferol (Yadav <i>et al.</i> 2020, Lakshmi <i>et al.</i> 2023)
	Fruit, Root	<i>Tribulus terrestris</i> L.	Chota gokhru	Zygophyllaceae	Powder of fruit and root with warm water is consumed.	Kaempferol, Kaempferol-3-glucoside, Kaempferol-3-rutinoside, Tribuloside, Furostanol, Diosgenin (Lakshmi <i>et al.</i> 2023)
	Seed	<i>Trigonella foenum-graecum</i> L.	Methi	Fabaceae	Powder of seed with cow's milk is given twice a day.	β -pinene, β -caryophyllene, Camphor, Neryl acetate (Lakshmi <i>et al.</i> 2023)
	Whole plant	<i>Vitex agnus-castus</i> L.	Nirgundi	Lamiaceae	Juice of whole plant is given twice a day orally.	Apigenin, 3-methylkaempfero, Luteolin, Casticin (Lakshmi <i>et al.</i> 2023)
	Rhizome	<i>Zingiber officinalis</i> Roscoe	Adrak	Zingiberaceae	Rhizome powder is given twice a day.	Gingerol, Shogaol Zingerone (Atashpour <i>et al.</i> 2017, Lakshmi <i>et al.</i> 2023)
12. Uterine fibroids	Bulb	<i>Allium cepa</i> L.	Pyaz	Amaryllidaceae	Bulb is consumed orally daily.	Quercetin (Block 1985, Islam <i>et al.</i> 2014)
	Clove	<i>Allium sativum</i> L.	Lehsun	Amaryllidaceae	Garlic paste is consumed orally.	Allicin (Block 1985, Islam <i>et al.</i> 2014)
	Leaves	<i>Camellia sinensis</i> L. Ktze.	Chai	Theaceae	Leaves extract is consumed.	Epigallocatechin gallate, Quercetin (Singh <i>et al.</i> 2011, Islam <i>et al.</i> 2014)
	Fruit	<i>Citrus sp.</i>	Nimbu, Santra	Rutaceae	Fruit juice is given orally.	Quercetin (Hertog <i>et al.</i> 1993)
	Rhizome	<i>Curcuma longa</i> L.	Haldi	Zingiberaceae	Powder of rhizome is consumed orally.	Curcumin (Wilken <i>et al.</i> 2011)
	Root	<i>Daucus carota</i> L.	Gajar	Apiaceae	The root is consumed.	Indole-3-carbinol (Aggarwal and Ichikawa 2005)
	Seed	<i>Glycine max</i> (L.) Merr.	Soybean	Fabaceae	Seed is consumed.	Isoliquiritigenin, Genistein (Cuendet <i>et al.</i> 2010, Islam <i>et al.</i> 2014)

Root	<i>Glycyrrhiza glabra</i> L.	Mulethi	Fabaceae	Root extract is used twice a day.	Isoliquiritigenin (Cuendet <i>et al.</i> 2010, Islam <i>et al.</i> 2014)
Fruit	<i>Psidium guajava</i> L.	Amrud	Myrtaceae	Fruit is consumed.	Lycopene (Holzapfel <i>et al.</i> 2013)
Fruit	<i>Punica granatum</i> L.	Anar	Lythraceae	Fruit is consumed.	Ellagic acid (Vattem <i>et al.</i> 2005, Islam <i>et al.</i> 2014)
Seed	<i>Raphanus sativus</i> L.	Muli	Brassicaceae	The powdered seeds of Muli (<i>Raphanus sativus</i>), seeds of carrot (<i>Daucus carota</i> subsp. <i>Sativus</i>), Vijayasar (<i>Pterocarpus marsupium</i>) (100 gram each), and Munnaka (<i>Vitis vinifera</i>) (200 gram) dried in shade and made into a paste. This paste (10 gram amount) with jaggery is given twice a day.	Indole-3-carbinol (Aggarwal and Ichikawa 2005)
Fruit	<i>Solanum lycopersicum</i> L.	Tamatar	Solanaceae	Fruit is consumed.	Quercetin (Hertog <i>et al.</i> 1993)
Seed	<i>Vicia faba</i> L.	Bakla	Fabaceae	Cooked beans/seeds are consumed.	Genistein (Islam <i>et al.</i> 2014)
Fruit	<i>Vitis vinifera</i> L.	Angoor	Vitaceae	Fruit is consumed.	Resveratrol, Ellagic acid (Labinskyy <i>et al.</i> 2006, Islam <i>et al.</i> 2014)

28.72° N, 77.78° E. Hapur district is a part of Meerut division. It covers an area of about 660 sq. km. in the north-west part of Uttar Pradesh. It falls within the National Capital Region. The district is bounded on the north by Meerut district, on the south by Bulandshahar district, on the east by Amroha district, and on the west by Ghaziabad district. The district consists of three tehsils: Hapur, Garhmukteshwar, and Dhaulana. Hapur is a part of the Indus-Gangetic Plain and the river Ganga forms the eastern boundary of Garhmukteshwar where people come for pilgrimage. Hapur district is the poorest in forest resources as the total area is reported to be 846 hectares which is 0.74% of the total geographical area (114276 hectares) of the district.

Data Collection, Identification & Documentation

The survey was carried out in different regions of the district from 2021 to 2023. Meanwhile, a questionnaire was used for an ethnomedicinal survey on the use of native medicinal plants for curing several kinds of gynecological disorders (Jain, 1989 Martin, 2010). Ethno-gynecological information was gathered through interviews and questionnaires with 12 herbal informants such as women, midwives, dais, and vaidyas of the Hapur district. The local language was used to gather ethnomedicinal information. The ethnic-gynecological data was recorded along with the vernacular name, the botanical name of the plant, family name, plant part(s) used, and the mode of preparing the ethnomedicine to cure women's gynecological disorders (Table 1).

Standard methods of collection, drying, mounting, preservation, and maintenance of plant specimens were followed according to Jain and Rao 1977. The identification and nomenclature of the listed plants were based on the online website "Plants of the World Online (POWO)" (<https://powo.science.kew.org>) taxonomic database. All the preserved herbarium specimens were deposited at the Herbarium of Raghunath Girls' Post Graduate College, Meerut (Uttar Pradesh).

Results And Discussions

Diversity of ethnomedicinal plant species

An ethnomedicinal survey related to gynecological disorders was conducted in district Hapur of Uttar Pradesh and 52 plant species belonging to 47 genera were reported. Among the 52 plant species, 51 plants were angiosperm and one plant species was gymnosperm. Among the angiosperm plants, 13% were monocot and 87% were dicot. Most cited families of medicinal plants documented were Fabaceae (17%), Lamiaceae, and Moraceae (8%), Apiaceae, Apocynaceae, Amaryllidaceae, Brassicaceae, and Zingiberaceae (4% each); twenty-five families had 2% each shown in Figure 2. Among 47 genera reported, *Ficus* (4 spp.), *Ocimum* (2 spp.), *Allium* (2 spp.) were dominant. Among 52 ethnomedicinal plant species, 26 (50%) plant species belong to herbs, followed by 13 (25%) trees, shrubs 10 (19%), climber 2 (4%), and grasses 1 (2%) shown in Figure 3. Among the

knowledge. Further clinical and pharmacological studies are necessary to explore the untapped potentials of the recorded plant species. Given the current state of knowledge on the chemical components of these plants, it is hoped that future research will identify the specific bioactive components that may be used to treat gynecological disorders. The current investigation indicates that further research is necessary to fully understand the extremely intriguing outcomes of gynecological problems. People from rural or tribal areas have a good understanding of the environment and the resources around them. To preserve this very effective traditional therapeutic technique, more research into this indigenous knowledge is vitally needed, as is its scientific confirmation, acceptance in the medical system, and popularization.

Acknowledgement

The traditional healers of district Hapur, to whom grateful acknowledgment is made for revealing the precious information on medicinal plant species related to gynecological disorders. Authors are thankful to Prof. R. K. Sharma, formerly Principal and Head of the Botany Department, S.S.V.P.G. College, Hapur, Mahant Ramprakash Muni, Vaidh Naresh Kumar, Dai Kallo Devi, Vaidh Omark Prasad Sharma, Shri Ashokanand ji, Vaidh Dhramveer Singh, Smt. Poonam Devi and other informants for their help and for sharing their treasure of knowledge about ethnomedicinal plants.

References

- Aafi E, Tabarrai M, Ardakani MM, Ardakani MRS and Lamardi SNS (2020). Evaluation of scientific evidence for abortifacient medicinal plants mentioned in traditional Persian medicine. *Traditional Medicine Research* **5(6)**: 449-463.
- Adeniyi BA, Ayeni FA and Ogunbanwo ST (2006). Antagonistic activities of lactic acid bacteria isolated from Nigerian fermented dairy food against organisms implicated in urinary tract infection. *Biotechnology* **5(2)**: 183-188.
- Afroze CA, Ahmed MN, Jahan R and Rahmatullah M (2022). Evaluation of herbal ingredients used in an ethno-polyherbal formulation for treating menorrhagia and dysmenorrhea in Bangladesh. *Phytomedicine Plus* **2(4)**: 100366.
- Aggarwal BB and Ichikawa H (2005). Molecular targets and anticancer potential of indole-3-carbinol and its derivatives. *Cell cycle* **4(9)**: 1201-1215.
- Atashpour S, Jahromi HK, Jahromi ZK and Maleknasab M (2017). Comparison of the effects of Ginger extract with clomiphene citrate on sex hormones in rats with polycystic ovarian syndrome. *Int. J. Rep. BioMed.* **15(9)**: 561-568.
- Attah AF, Moody JO, Sonibare MA, Salahdeen HH, Akindele OO, Nnamani PO and Raji Y (2020). Aqueous extract of Moringa oleifera leaf used in Nigerian ethnomedicine alters conception and some pregnancy outcomes in Wistar rats. *S. Afr. J. Botany* **129**: 255-262.
- Behera K, Sharma U, Mitra S and Sharma KC (2021). Review of modified Panchvalkal extract with special reference to Leucorrhoea. *WJPMR* **7(4)**: 194-197.
- Bhandary MJ, Chandrashekar KR and Kaveriappa KM (1995). Medical ethnobotany of the siddis of Uttara Kannada district, Karnataka, India. *J. Ethnopharm.* **47(3)**: 149-158.
- Bhargava SK (1986). Estrogenic and postcoital contraceptive activity in rats of butin isolated from Butea monosperma seed. *J. Ethnopharm.* **18(1)**: 95-101.
- Chakraborty AJ, Uddin TM, Zidan BRM, Mitra S, Das R, Nainu F and Emran TB (2022). Allium cepa: A treasure of bioactive phytochemicals with prospective health benefits. *Evidence-Based Complementary and Alternative Medicine: eCAM*, **2022**: 1-27.
- Choroshio SH, Malik N, Panesar G, Kumari P, Jangra S, Kaur R and Murthy HC (2023). Phytochemicals: an alternative for infertility treatment and associated conditions. *Oxidative Medicine and Cellular Longevity* **2023**: 1-21.
- Cuendet M, Guo J, Luo Y, Chen S, Oteham CP, Moon RC and Pezzuto JM (2010). Cancer chemopreventive activity and metabolism of isoliquiritigenin, a compound found in licorice. *Cancer prevention research* **3(2)**: 221-232.
- Dachtler M, Van de Put FH, Stijn F, Beindorff CM and Fritsche J (2003). On-line LC-NMR-MS characterization of sesame oil extracts and assessment of their antioxidant activity. *European journal of lipid science and technology* **105(9)**: 488-496.
- Das DC, Sinha NK and Das M (2015). The use of medicinal plants for the treatment of gynaecological disorders in the eastern parts of India. *I. J. Obst. Gynae.* **2 (1)**: 16-27.
- Dietz BM, Hajirahimkhan A, Dunlap TL and Bolton JL (2016). Botanicals and their bioactive phytochemicals for women's health. *Pharmacological reviews* **68(4)**: 1026-1073.
- Downie A, Fink CS and Awad AB (1999). Effect of phytosterols on MDA-MB-231 human breast cancer cell growth. *Faseb journal* **13(4)**: A333.
- El Khoury D, Matar R and Touma T (2019). Curcumin and endometrial carcinoma: an old spice as a novel agent. *Int. J. Women's Health* **16**: 249-256.
- Elham MN, Yasmin M, Mohammed M and Mohammed MS (2019). Effect of crocin on letrozole-induced polycystic ovarian syndrome. *The Medical Journal of Cairo University* **87**: 5237-5243.
- Gautam P, Bairwa O, Dwivedi D, Kumar S and Singh AK (2017). A Progressive step towards classical uses of Medicinal Plants in Gynecological Disorders. *J. Ayu. Int. Med. Sci.* **2(03)**: 225-236.
- Goyal S, Gupta N and Chatterjee S (2016). Investigating therapeutic potential of Trigonella foenum-graecum L. as our defence mechanism against several human diseases. *Journal of toxico.* **2016**: 1-10.
- Gupta U and Solanki H (2013). Herbal folk remedies used in treatment of Gynecological disorders by tribals of Simalwara Region, Dungarpur, Rajasthan. *Int. J. Pure App. Sci. Tech.* **17(1)**: Pp100.
- Gupta OP, Anand KK, Ghatak BR and Atal CK (1978). Vasicine, alkaloid of Adhatoda vasica, a promising uterotonic abortifacient. *Ind. J. Exp. Bio.* **16(10)**: 1075-1077.
- Hertog MG, Hollman PC and Van de Putte B (1993). Content of potentially anticarcinogenic flavonoids of tea infusions, wines, and fruit juices. *Journal of agricultural and food chemistry* **41(8)**: 1242-1246.
- Holzappel NP, Holzappel BM, Champ S, Feldthusen J, Clements J and Hutmacher DW (2013). The potential role of lycopene for the prevention and therapy of prostate cancer: from molecular mechanisms to clinical evidence. *Int. J. Mol. Sci.* **14(7)**: 14620-14646.

- Hossain MT and Hoq MO (2016). Therapeutic use of *Adhatoda vasica*. *A. J. Med. Bio Res.* **2(2)**: 156-163.
- Islam MS, Akhtar MM, Ciavattini A, Giannubilo SR, Protic O, Janjusevic M and Ciarmela P (2014). Use of dietary phytochemicals to target inflammation, fibrosis, proliferation, and angiogenesis in uterine tissues: promising options for prevention and treatment of uterine fibroids?. *Molecular nutrition & food research* **58(8)**: 1667-1684.
- Jain SK (1989). Methods and approaches in Ethnobotany. (Ed) Society of Ethnobotanists, CDRI. Lucknow.
- Jain SK and Rao RR (1977). A Handbook of Field and Herbarium Methods. Today and Tomorrow's Printers and Publishers, New Delhi.
- Joshi P (1995). Ethnobotany of the primitive tribes in Rajasthan. Printwell Jaipur.
- Katewa SS (2009). Indigenous people and forests: Perspectives of an ethnobotanical study from Rajasthan (India). *Herbal drugs: Ethnomedicine to modern medicine* Pp 33-56.
- Khurana N, Sharma N, Patil S and Gajbhiye A (2016). Phytopharmacological properties of *Sida cordifolia*: a review of folklore use and pharmacological activities. *Asian J Pharm Clin Res.* **2**: 52-58.
- Labinskyy N, Csiszar A, Veress G, Stef G, Pacher P, Oroszi G and Ungvari Z (2006). Vascular dysfunction in aging: potential effects of resveratrol, an anti-inflammatory phytoestrogen. *Current medicinal chemistry* **13(9)**: 989-996.
- Lakshmi JN, Babu AN, Kiran SM, Nori LP, Hassan N, Ashames A and Shaik AB (2023). Herbs as a source for the treatment of polycystic ovarian syndrome: A systematic review. *BioTech* **12(1)**: 1- 4.
- Lawal IO, Amao AO, Lawal KO, Alamu OT and Sowunmi IL (2013). Phytotherapy approach for the treatment of gynaecological disorder among women in Ido Local Government Area of Ibadan, Oyo State, Nigeria. *J. Adv. Sci. Res.* **4(03)**: 41-44.
- Li L, Li C, Pan P, Chen X, Wu X, Ng EHY and Yang D (2015). A single arm pilot study of effects of berberine on the menstrual pattern, ovulation rate, hormonal and metabolic profiles in anovulatory Chinese women with polycystic ovary syndrome. *PloS one* **10(12)**: e0144072.
- Malpani A, Mahurkar N and Aswar U (2020). Phytochemical analysis and antifertility potential of *Cynodon dactylon* in female Wistar rats: A herbal approach towards contraception. *Chinese Herbal Medicines* **12(3)**: 281-288.
- Martin GJ (2010). Ethnobotany: a methods manual. Routledge.
- Majumder M, Debnath S, Gajbhiye RL, Saikia R, Gogoi B, Samanta SK and Mukhopadhyay R (2019). *Ricinus communis* L. fruit extract inhibits migration/invasion, induces apoptosis in breast cancer cells and arrests tumor progression in vivo. *Scientific Reports* **9(1)**: 14493.
- Mishra D, Singh RK, Srivastava RK and Dubey SR (2013). Ethnomedicinal plants used to cure the gynaecological disorders by ethnic populace of Sitapur district, Uttar Pradesh, India. *Med. Plants-Int. Jou. Phytomed. Rel. Ind.* **5(4)**: 238-245.
- Mishra JN and Verma NK (2017). A brief study on *Catharanthus roseus*: A review. *Int. J. Res. Pharmacy Pharmaceut Sci* **2(2)**: 20-23.
- Mishra N, Verma R and Jadaun P (2022). Study on the effect of berberine, myoinositol, and metformin in women with polycystic ovary syndrome: a prospective randomised study. *Cureus* **14(1)**: e21781.
- Mubeen AM, Krishnan V, Mubeen Y, Jeyaprakash G (2022). Reliability of siddha systems of medicine on polycystic ovary syndrome with scientific justification -a review. *IJP* **13(1)**: 1-8.
- Murugesu S, Selamat J and Perumal V (2021). Phytochemistry, pharmacological properties, and recent applications of *Ficus benghalensis* and *Ficus religiosa*. *Plants* **10(12)**: p2749.
- Panda T, Mishra N, Rahimuddin S, Pradhan BK, Rout SD and Mohanty RB (2018). Folk medicine used for the treatment of gynaecological disorders in rural areas of Bhadrak district, Odisha, India. *Botanica Lithuanica* **24(2)**: 132-142.
- Penagos Tabares F, Bedoya Jaramillo JV and Ruiz-Cortés ZT (2014). Pharmacological overview of galactogogues. *Veterinary medicine international* **2014**: p20.
- Pareek A, Pant M, Gupta MM, Kashania P, Ratan Y, Jain V and Chuturgoon AA (2023) *Moringa oleifera*: An updated comprehensive review of its pharmacological activities, ethnomedicinal, phytopharmaceutical formulation, clinical, phytochemical, and toxicological aspects. *International journal of molecular sciences* **24(3)**: p2098.
- Sarih K, Siradjuddin S, Maddepungeng M, Hadju V, Saleh A and Tanzaha I (2020). *Moringa oleifera* intake during pregnancy and breastfeeding toward docosahexaenoic acid and arachidonic acid levels in breast milk. *J. Med. Sci.* **8(B)**: 757-761.
- Satyavati GV, Raina MK and Sharma M (1987). *Medicinal plants of India* (Vol. 2). New Delhi: Indian Council of Medical Research.
- Shantha TR, Shubhashree MN, Reddy MP and Venkateshwarlu G (2014). Pharmacognostic and Physicochemical Evaluation of the Different Varieties of *Sesamum indicum* L.-A Comparative Study. *Res. J. Pharm. Phytochem.* **6(2)**: 99-106.
- Sharma AK and Deshwal N (2011). An overview: on phytochemical studies of *Buta monosperma*. *IJPTR* **3(2)**: 864-871.
- Sharma N, Sharma A, Bhatia G, Landi M, Brestic M, Singh B and Bhardwaj R (2019). Isolation of phytochemicals from *Bauhinia variegata* L. Bark and their in vitro antioxidant and cytotoxic potential. *Antioxidants* **8(10)**: p 492.
- Singh BN, Shankar S and Srivastava RK (2011). Green tea catechin, epigallocatechin-3-gallate (EGCG): mechanisms, perspectives and clinical applications. *Biochemical pharmacology* **82(12)**: 1807-1821.
- Suh YA, Jo SY, Lee HY and Lee C (2015). Inhibition of IL-6/STAT3 axis and targeting Axl and Tyro3 receptor tyrosine kinases by apigenin circumvent taxol resistance in ovarian cancer cells. *Int. J. Oncology* **46(3)**: 1405-1411.
- Takoore H, Aumeeruddy MZ, Rengasamy KR, Venugopala KN, Jeewon R, Zengin G Mahomoodally MF (2019). A systematic review on black pepper (*Piper nigrum* L.): from folk uses to pharmacological applications. *Cri. rev. food science and nutrition* **59(1)**: S210-S243.
- Thakur S, Kaurav H and Chaudhary G (2021). Shatavari (*Asparagus Racemosus*)-the best female reproductive tonic. *Int. J. Res. Rev.* **8(5)**: 73-84.
- Tossetta G, Fantone S, Giannubilo SR and Marzioni D (2021). The multifaceted actions of curcumin in pregnancy outcome. *Antioxidants* **10(1)**: p126.
- Tripathi R, Mishra RP, Singh AR and Dwivedi SN (2011). Folklore use of some medicinal plants in the treatment of UTI infections. *Int. J. Drug Dis. Her. Res.* **1**: 58-60.
- Tsantila EM, Esslinger N, Christou M, Papageorgis P and Neophytou CM (2024). Antioxidant and Anticancer Activity of *Vitis*

- vinifera Extracts in Breast Cell Lines. *Life* **14(2)**: p228.
- Vattem DA and Shetty K (2005). Biological functionality of ellagic acid: a review. *J. Food bio.* **29(3)**: 234-266.
- Vora D, Kapadia H, Dinesh S, Sharma S and Manjegowda DS (2023). Gymnema sylvestre as a potential therapeutic agent for PCOS: insights from mRNA differential gene expression and molecular docking analysis. *Fut. J. Pharm. Sci.* **9(1)**: p76.
- Wadhvani BD, Mali D, Vyas P, Nair R and Khandelwal P (2021). A review on phytochemical constituents and pharmacological potential of Calotropis procera. *RSC advances* **11(57)**: 35854-35878.
- Wang Y and Zhang S (2018). Berberine suppresses growth and metastasis of endometrial cancer cells via miR-101/COX-2. *Biomedicine & Pharmacotherapy* **103**: 1287-1293.
- Way TD, Kao MC and Lin JK (2004). Apigenin induces apoptosis through proteasomal degradation of HER2/neu in HER2/neu-overexpressing breast cancer cells via the phosphatidylinositol 3-kinase/Akt-dependent pathway. *J. Bio. Che.* **279(6)**: 4479-4489.
- Wilken R, Veena MS, Wang MB and Srivatsan ES (2011). Curcumin: A review of anti-cancer properties and therapeutic activity in head and neck squamous cell carcinoma. *Mol. cancer* **10**: 1-19.
- Woźniak M, Krajewski R, Makuch S and Agrawal S (2021). Phytochemicals in gynecological cancer prevention. *Int. J. Mol. Sci.* **22(3)**: p1219.
- Xu Y, Ma Q, Chen D, Yin Q and Zhang W (2020). Effects of Ginkgolide B on the Proliferation and Apoptosis of Cervical Cancer Cells. *Cur. Top. Nut. Res.* **18(3)**: p227.
- Xu Y and Cai L (2018). L-mimosine induces caspase-9-mediated apoptosis in human osteosarcoma cells. *Mol. Med. Rep.* **17(3)**: 4695-4701.
- Yadav K, Ghadge P, Langeh A, Kalbhare S, Phadtare P and Bhoite R (2020). A review on herbal medicinal plant for treatment of polycystic ovarian syndrome (PCOS). *A. J. Ph. Res. Dev.* **8(4)**: 83-87.
- Yang H, Kim HJ, Pyun BJ and Lee HW (2018). Licorice ethanol extract improves symptoms of polycytic ovary syndrome in Letrozole-induced female rats. *Int. Med. Res.* **7(3)**: 264-270.
- Yavari M, Rouholamin S, Tansaz M, Bioos S and Esmaeili S (2014). Sesame a treatment of menstrual bleeding cessation in iranian traditional medicine: Results from a pilot study. *Shiraz E-Medical Journal* **15(3)**: e21893
- Ye B, Aponte M, Dai Y, Li L, Ho MCD, Vitonis A and Cramer, DW (2007). Ginkgo biloba and ovarian cancer prevention: epidemiological and biological evidence. *Cancer letters* **251(1)**: 43-52.
- Zhang Y, Munday R and Jobson HE (2006). Cruciferous vegetables, Derived ITC, Chemical structure of ITC. *J. Agr. F. Che.* **54**: 9370-9376.